


Male to Female Transgender AIDS Cases


Cumulative AIDS cases among male to female transgendered persons (MTF), 1981 - 2000



This map shows the geographic distribution of cumulative AIDS cases among MTF from 1981 through 2000 by US Census Block Groups in San Francisco City and County. The highest concentration of MTF AIDS cases have been in the low-income, inner-city Tenderloin neighborhood and adjacent areas of Nob Hill and SOMA. Scattered cases have occurred among residents of Mission, Western Addition, Duboce Triangle, Mission Dolores, Bayview, and Ingleside.

Data include persons who were residents of San Francisco at the time of their diagnosis. Data do not include persons with HIV infection who have not progressed to an AIDS diagnosis. MTF may be under-reported in surveillance data if their sex is recorded as female or male without indication of their gender reassignment.


Male to female transgendered persons (MTF) living with AIDS, 2000



This map shows the geographic distribution of MTF living with AIDS as of 2000 by US Census Block Groups in San Francisco City and County. MTF living with AIDS tend to reside in the low-income, inner-city Tenderloin neighborhood and adjacent areas of Nob Hill and SOMA. Mission, Mission Dolores, Western Addition, Bayview and low-income areas in the south of the city are also home to many MTF living with AIDS.

Data include persons who were residents of San Francisco at the time of their diagnosis and who were known to be alive on 31 December, 2000. Data do not include persons with HIV infection who have not progressed to an AIDS diagnosis. MTF may be under-reported in surveillance data if their sex is recorded as female or male without indication of their gender reassignment.


Male to female transgendered persons (MTF) living with AIDS by race/ethnicity, 2000



White male to female transgendered persons (MTF) living with AIDS, 2000

This map shows the geographic distribution of white MTF living with AIDS as of 2000 by US Census Tracts in San Francisco City and County. White MTF living with AIDS reside in the Tenderloin, Western Addition, Mission, and Potrero Hill.


Data include persons who were residents of San Francisco at the time of their diagnosis and who were known to be alive on 31 December, 2000. Data do not include persons with HIV infection who have not progressed to an AIDS diagnosis. MTF may be under-reported in surveillance data if their sex is recorded as female or male without indication of their gender reassignment.



African Americans male to female transgendered persons (MTF) living with AIDS, 2000

This map shows the geographic distribution of African American MTF living with AIDS as of 2000 by US Census Tracts in San Francisco City and County. African American MTF living with AIDS reside in low-income, inner-city area of the Tenderloin, SOMA, Western Addition, Mission, and Bayview.


Data include persons who were residents of San Francisco at the time of their diagnosis and who were known to be alive on 31 December, 2000. Data do not include persons with HIV infection who have not progressed to an AIDS diagnosis. MTF may be under-reported in surveillance data if their sex is recorded as female or male without indication of their gender reassignment.



Latino male to female transgendered persons (MTF) living with AIDS, 2000

This map shows the geographic distribution of Latino MTF living with AIDS as of 2000 by US Census Tracts in San Francisco City and County. Latino Transgendered persons living with AIDS reside in an inner-city band extending from Nob Hill in the northeast through Mission to Bernal Heights in the south.

Data include persons who were residents of San Francisco at the time of their diagnosis and who were known to be alive on 31 December, 2000. Data do not include persons with HIV infection who have not progressed to an AIDS diagnosis. MTF may be under-reported in surveillance data if their sex is recorded as female or male without indication of their gender reassignment.




Asian/Pacific Islander male to female transgendered persons (MTF) living with AIDS, 2000

This map shows the geographic distribution of Asian/Pacific Islander MTF living with AIDS as of 2000 by US Census Tracts in San Francisco City and County. Asian/Pacific Islander MTF living with AIDS reside in a few areas of Tenderloin, SOMA, Mission, Haight-Ashbury, and Outer Mission.

Data include persons who were residents of San Francisco at the time of their diagnosis and who were known to be alive on 31 December, 2000. Data do not include persons with HIV infection who have not progressed to an AIDS diagnosis. MTF may be under-reported in surveillance data if their sex is recorded as female or male without indication of their gender reassignment.

AIDS Cases among Youth (13 to 24 years old), Children (under 13 years old), and Transfusion/Blood Products Recipients


Cumulative AIDS cases among youth, 1981-2000



This map shows the geographic distribution of cumulative AIDS cases among youth age 13 to 24 years from 1981 through 2000 by US Census Block Groups in San Francisco City and County. The highest concentration of AIDS cases among youth have been in the low-income, inner-city neighborhoods of the Tenderloin, Mission, and Western Addition. A concentration of youth with AIDS also reside in Duboce Triangle and Mission Dolores, a transitional area between the more affluent Castro and lower-income Haight, Mission, and Western Addition.

Data include persons who were residents of San Francisco and age 13 to 24 years at the time of their diagnosis. In some cases, HIV transmission may have occurred prior to or around the time of delivery. Data do not include persons with HIV infection who have not progressed to an AIDS diagnosis.


Youth living with AIDS, 2000



This map shows the geographic distribution of youth age 13 to 24 years living with AIDS as of 2000 by US Census Block Groups in San Francisco City and County. Youth living with AIDS tend to reside in the low-income, inner-city neighborhoods of Tenderloin, Mission, and Western Addition. A concentration of youth with AIDS also reside in Duboce Triangle and Mission Dolores, a transitional area between the more affluent Castro and lower-income Haight, Mission, and Western Addition.

Data include persons who were residents of San Francisco and age 13 to 24 years at the time of their diagnosis and who were known to be alive on 31 December, 2000. Data do not include persons with HIV infection who have not progressed to an AIDS diagnosis. MTF may be under-reported in surveillance data if their sex is recorded as female or male without indication of their gender reassignment.


Cumulative pediatric AIDS cases, 1981-2000



This map shows the geographic distribution of cumulative AIDS cases among children under the age of 13 years diagnosed from 1981 through 2000 by US Census Block Groups in San Francisco City and County. Only a few scattered neighborhoods record pediatric cases. These tend to be in low-income Block Groups in Bayview, Potrero Hill, Mission, Tenderloin, SOMA, and Western Addition. Pediatric cases are also found in the scattered blocks in Sunset and southern tier of the city.

Data include persons who were residents of San Francisco and under 13 years old at the time of their diagnosis. We assume that most cases resulted from infection prior to or at the time of delivery. Data do not include persons with HIV infection who have not progressed to an AIDS diagnosis.

Cumulative transfusion/blood product AIDS cases, 1981-2000



This map shows the geographic distribution of cumulative AIDS cases resulting from transfusion of blood and blood products from 1981 through 2000 by US Census Block Groups in San Francisco City and County. No concentration of transfusion-associated HIV/AIDS is evident - cases are scattered throughout the city. The lack of pattern likely results from the diversity of conditions that necessitate transfusion unrelated to sexual or drug use behavior, including hemophilia, major surgery, severe anemia.

Data include persons who were residents of San Francisco at the time of their diagnosis. In some cases, HIV transmission may have occurred prior to or around the time of delivery. Data do not include persons with HIV infection who have not progressed to an AIDS diagnosis.

PAGE INTENTIONALLY LEFT BLANK