


**San Francisco  
Sexually Transmitted Disease  
Annual Summary, 2007**



**San Francisco Department of Public Health  
Population Health and Prevention Division  
Sexually Transmitted Disease Prevention and Control Services  
San Francisco, California USA**

September, 2008

## Contents

Preface.....	iii
I. Reported Morbidity .....	1
Chlamydia .....	4
Gonorrhea .....	8
Syphilis.....	11
Gender .....	18
Race and Ethnicity .....	21
Age.....	29
Geography .....	36
Adolescents.....	51
Congenital Syphilis.....	70
Rectal and pharyngeal infections in men .....	72
Other STDs .....	79
II. Prevalence Data.....	81
Screening program.....	81
Sentinel Surveillance.....	86
Detention facilities .....	91
III. City Clinic .....	96
Patient demographics .....	96
STDs .....	105
HIV Testing .....	110
Appendix I. Demographic Breakdowns of STD Morbidity.....	114
Appendix II. Demographic Breakdowns for City Clinic .....	131

### Copyright Information

All material contained in this report is in the public domain and may be used and reprinted without special permission; citation to source, however, is appreciated.

Inquiries and requests for additional copies and future updates should be addressed to:

**Robert P. Kohn, MPH**  
**Sexually Transmitted Disease Prevention and Control Services**  
**1360 Mission Street, Suite 401**  
**San Francisco, CA 94103**  
**(415) 355-2000**

### Suggested Citation

**STD Control Section. San Francisco Sexually Transmitted Disease Annual Summary, 2007. San Francisco Department of Public Health, San Francisco, California. July 2007.**

### Note on totals for previous years

Numbers in this document listed for past years may not match totals in previous reports. Totals may increase due to late reports, and may decrease when duplicate reports are eliminated or cases are subsequently identified as out of our jurisdiction.

### Technical Note

All graphs, tables and maps in this report were produced using SAS for Windows version 9.1. The document itself was created with *Microsoft Word*.

## Preface

*Abbreviations used in this report include the following:*

OOJ.....	out of jurisdiction
CHN .....	Community Health Network
PMD .....	private medical doctor
SFGH .....	San Francisco General Hospital
CHPY .....	Community Health Programs for Youth
P&S.....	primary and secondary (syphilis)
GC.....	gonorrhea
CT .....	chlamydia
SYPH .....	early syphilis
MSM.....	men who have sex with men
IDU.....	injecting drug user



## I. Reported Morbidity

### Sources of data

Title 17 of the California Administrative Code requires all clinicians treating or knowing of a patient with a suspected or documented reportable sexually transmitted disease (STD) and all laboratories with a test result or isolate suggesting infection by a reportable agent of an STD to report their findings to the patient's local health department. In 2007 the list of reportable STDs included syphilis of any stage, gonorrhea, chlamydia, chancroid, pelvic inflammatory disease (PID), and non-gonococcal urethritis (NGU). Reports of morbidity must include the patient's name and address and demographic information (gender, age, and race or ethnicity). As recommended by the United States Centers for Disease Control and Prevention (CDC), gender of sex partners became a required reportable item in San Francisco, effective July 2004.

Recent reviews of the San Francisco STD surveillance system suggest that reporting is nearly complete for syphilis, gonorrhea and chlamydia when a diagnostic test was performed in a laboratory, because laboratories routinely report the positive test result. However, many men and women with STDs are never diagnosed, either because they do not develop symptoms, are not screened appropriately by their provider, or have no access to health care. This is particularly true for chlamydia, since infection is often asymptomatic and routine screening is not widespread. Furthermore, demographic data is often missing from reports; in 2007, about twenty percent of all STDs reported lacked information on race or ethnicity. For these reasons, reported totals should be considered minimal estimates of the true number of STDs in the community.

### Disease rates

Rates have been listed in most tables along with reporting totals. Rates are equal to the number of STD cases within the specified population per 100,000 San Francisco residents in that population per year. Population figures for rates are from the US Census Data, except for congenital syphilis, where live birth totals for each year are used instead.

Rates should be used when comparing STD levels among different populations, as differences in disease totals are affected by the size of the population as well as incidence.

Comparison rates for California, the United States and the New York-Newark-Edison, NY-NJ-PA and Los Angeles-Long Beach-Santa Ana, CA metropolitan statistical areas (MSAs) are from *Sexually Transmitted Disease Surveillance, 2005*, by the Centers for Disease Control and Prevention, US Department of Health and Human Services (November 2007).

### Census data

Denominators for all rates in this report are based on the 2000 US Census data.

Data on race or ethnicity of STD patients is typically reported as a single value, with "Hispanic" or "Latino" as a category exclusive of all others. In the 2000 US Census, however, race was collected as a multiple-choice item, with Hispanic ethnicity recorded independently of race. In order to make denominators from the census data match totals from case reports, totals for residents reporting more than one race in the census data were divided among totals for residents indicating only one race. Failure to do so would have artificially increased all race-specific rates, since there are no patients recorded as "multi-racial" among the STD case reports.

*Summary*

*Chlamydia*

Number of reported chlamydia cases (2007) = 3,939

Chlamydia rate (2007) = 507.1 per 100,000 San Francisco residents

Change in chlamydia rate from 2006 = 2.8% decline

*Gonorrhea*

Number of reported gonorrhea cases (2007) = 2,010

Gonorrhea rate (2007) = 258.8 per 100,000 San Francisco residents

Change in gonorrhea rate from 2006 = 18.5% decline

*Syphilis*

Number of reported early syphilis cases (2007) = 352

Number of reported primary and secondary (P&S) syphilis cases (2007) = 202

Early syphilis rate (2007) = 45.3 per 100,000 San Francisco residents

P&S syphilis rate (2007) = 26.0 per 100,000 San Francisco residents

Change in early syphilis rate from 2006 = 16.3% decline

Change in P&S syphilis rate from 2006 = 16.9% decline




Figure 1. Major reportable STDs, San Francisco, 2003-2007. Note difference in scale for early syphilis.

Table 1. Reportable STD cases and rates, San Francisco, 2003-2007. Rates equal cases per 100,000 residents per year, except for non-gonococcal urethritis (NGU) (rates equal cases per 100,000 men), pelvic inflammatory disease (PID) (cases per 100,000 women), and congenital syphilis (cases per 100,000 live births). Note: no cases of Granuloma inguinale or late symptomatic syphilis have been reported since 1992.

	Reported cases					Rate				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
CHLAMYDIA	3,350	3,663	3,710	4,053	3,939	431.3	471.6	477.6	521.8	507.1
GONORRHEA	1,795	2,153	2,411	2,468	2,010	231.1	277.2	310.4	317.7	258.8
SYPHILIS (TOTAL)	642	682	511	534	469	82.7	87.8	65.8	68.7	60.4
---PRIMARY	108	125	82	83	63	13.9	16.1	10.6	10.7	8.1
---SECONDARY	228	224	167	160	139	29.4	28.8	21.5	20.6	17.9
---(TOTAL P&S)	336	349	249	243	202	43.3	44.9	32.1	31.3	26.0
---EARLY LATENT	191	203	179	177	150	24.6	26.1	23.0	22.8	19.3
---(TOTAL EARLY)	527	552	428	420	352	67.8	71.1	55.1	54.1	45.3
---UNKNOWN LATENT [1]	5	8	5	9	5	0.6	1.0	0.6	1.2	0.6
---LATE LATENT	110	122	78	105	112	14.2	15.7	10.0	13.5	14.4
---NEUROSYPHILIS	19	37	18	9	10	2.4	4.8	2.3	1.2	1.3
CONGENITAL SYPHILIS (TOTAL)	0	1	0	0	1	0.0	12.1	0.0	(N/A)	(N/A)
---BIRTHS	0	1	0	0	1	0.0	12.1	0.0	(N/A)	(N/A)
PID (ALL)	109	92	58	43	40	28.5	24.1	15.2	11.3	10.5
---PROBABLE PID [2]	51	50	32	29	29	13.4	13.1	8.4	7.6	7.6
---SUSPECT PID	58	42	26	14	11	15.2	11.0	6.8	3.7	2.9
NON-GONOCOCCAL URETHRITIS	985	954	899	859	725	249.5	241.6	227.7	217.6	183.6
LYMPHOGRANULOMA VENEREUM	0	22	5	18	4	0.0	2.8	0.6	2.3	0.5

<sup>1</sup> cases not known to be less than one year's duration where the patient is 40 years old or less and the initial titer is 1:32 or higher.

<sup>2</sup> PID cases meeting CDC case definition.

## Chlamydia

Number of reported chlamydia cases (2007) = 3,939

Chlamydia rate (2007) = 507.1 per 100,000 San Francisco residents

Change in chlamydia rate from 2006 = 2.8% decline

- Overall, chlamydia rates declined in San Francisco in 2007 after over a decade of increases.
- Rates of chlamydia declined 8.9% among females and increased 3.5% among males.
- Chlamydia was most often diagnosed by private medical doctors and clinics and private hospitals. San Francisco City Clinic accounted for 20.6% of the total chlamydia diagnoses.
- Chlamydia rates in San Francisco exceeded those for Los Angeles MSA, New York MSA, the state of California and the United States in 2006.




Figure 2. Reported chlamydia cases, San Francisco, 1985-2007. Note: chlamydia became a reportable disease in 1989; the Public Health Laboratory implemented nucleic acid amplification testing in Fall 1996.




Figure 3. Chlamydia rate, San Francisco, 2003-2007.




Figure 4. Chlamydia cases by health care provider for 2007. "Jails" includes cases from both adult and adolescent detention facilities. Not included: 107 cases reported by providers outside San Francisco jurisdiction.




Figure 5. Regional chlamydia rates compared for 2006, San Francisco vs. Los Angeles MSA, New York MSA, total California and total U.S.




Figure 6. Comparison of trends in chlamydia 2002-2006, San Francisco vs. Los Angeles MSA, New York MSA, total California and total U.S. (MSA data not available for 2001).

Table 2. Chlamydia cases by health care provider, San Francisco, 2003-2007.

	Reported cases					Percent of reports				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
OOJ PROVIDERS	111	117	146	107	126	3.3%	3.1%	3.9%	2.6%	3.1%
CITY CLINIC	847	783	848	794	812	25.2%	21.3%	22.8%	19.5%	20.6%
PUBLIC CLINIC (CHN)	124	176	143	167	134	3.7%	4.8%	3.8%	4.1%	3.4%
JAILS	250	314	297	337	238	7.4%	8.5%	8.0%	8.3%	6.0%
PRIVATE CLINIC/PMD	898	916	1,043	1,313	1,326	26.8%	25.0%	28.1%	32.3%	33.6%
PRIVATE HOSPITAL	688	724	626	835	799	20.5%	19.7%	16.8%	20.6%	20.2%
CHPY	79	97	116	125	137	2.3%	2.6%	3.1%	3.0%	3.4%
SFGH	274	251	228	140	112	8.1%	6.8%	6.1%	3.4%	2.8%
OUTREACH	21	15	22	23	28	0.6%	0.4%	0.5%	0.5%	0.7%
MAGNET	58	270	241	212	227	1.7%	7.3%	6.4%	5.2%	5.7%
(ALL PROVIDERS)	3,350	3,663	3,710	4,053	3,939	100%	100%	100%	100%	100%

## Gonorrhea

Number of reported gonorrhea cases (2007) = 2,010

Gonorrhea rate (2007) = 258.8 per 100,000 San Francisco residents

Change in gonorrhea rate from 2006 = 18.5% decline

- Overall, gonorrhea rates declined in San Francisco in 2007 after over a decade of increases.
- Rates of gonorrhea declined 18.6% among females and increased 18.7% among males.
- San Francisco City Clinic diagnosed 37.1% of total gonorrhea cases. Private medical doctors and clinics accounted for 20.1% of total diagnoses.
- Gonorrhea rates in San Francisco exceeded those for Los Angeles MSA, New York MSA, the state of California and the United States in 2006.







Figure 7. Gonorrhea rate, San Francisco, 2003-2007.



*Figure 8. Gonorrhea cases by health care provider for 2007. "Jails" includes cases from both adult and adolescent detention facilities. Not included: 34 cases reported by providers outside San Francisco jurisdiction.*



*Figure 9. Regional gonorrhea rates compared for 2006 San Francisco vs. Los Angeles MSA, New York MSA, total California and total U.S.*



**Table 3.** Gonorrhea cases by health care provider, San Francisco, 2003-2007. See Introduction for list of abbreviations used.

	Reported cases					Percent of reports				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
OOJ PROVIDERS	31	52	60	34	31	1.7%	2.4%	2.4%	1.3%	1.5%
CITY CLINIC	869	932	930	961	746	48.4%	43.2%	38.5%	38.9%	37.1%
PUBLIC CLINIC (CHN)	72	82	86	88	77	4.0%	3.8%	3.5%	3.5%	3.8%
JAILS	58	73	107	103	81	3.2%	3.3%	4.4%	4.1%	4.0%
PRIVATE CLINIC/PMD	339	250	346	420	405	18.8%	11.6%	14.3%	17.0%	20.1%
PRIVATE HOSPITAL	235	247	315	321	262	13.0%	11.4%	13.0%	13.0%	13.0%
CHPY	27	34	42	51	27	1.5%	1.5%	1.7%	2.0%	1.3%
SFGH	87	110	145	85	63	4.8%	5.1%	6.0%	3.4%	3.1%
OUTREACH	3	12	15	18	27	0.1%	0.5%	0.6%	0.7%	1.3%
MAGNET	74	361	365	387	291	4.1%	16.7%	15.1%	15.6%	14.4%
(ALL PROVIDERS)	1,795	2,153	2,411	2,468	2,010	100%	100%	100%	100%	100%

## Syphilis

Number of reported early syphilis cases (2007) = 352

Number of reported primary and secondary (P&S) syphilis cases (2007) = 202

Early syphilis rate (2007) = 45.3 per 100,000 San Francisco residents

P&S syphilis rate (2007) = 26.0 per 100,000 San Francisco residents

Change in early syphilis rate from 2006 = 16.3% decline

Change in P&S syphilis rate from 2006 = 16.9% decline

- Both total early and primary and secondary syphilis rates declined in San Francisco for the 3<sup>rd</sup> consecutive year.
- Nearly 90% of reported early syphilis occurred among men who have sex with men (MSM).
- San Francisco City Clinic diagnosed 24.1% of total early syphilis. Private medical doctors and clinics accounted for an additional 27.0% of total diagnoses. Private hospitals diagnosed 25% of the total reported early syphilis cases in 2007.
- Early syphilis rates in San Francisco well exceeded those for Los Angeles MSA, New York MSA, the state of California and the United States in 2006.




Figure 11. Reported early syphilis cases, San Francisco, 1955-2007.




Figure 12. Syphilis rates, San Francisco, 2003-2007.




Figure 13. Trends in primary and secondary (P&S) and early syphilis, San Francisco, 2003-2007.




Figure 14. *Syphilis cases by stage of disease, San Francisco, 2007.*




Figure 15. *Early syphilis cases by health care provider for 2007. "Jails" includes cases from both adult and adolescent detention facilities. Not included: 12 cases reported by providers outside San Francisco jurisdiction.*




Figure 16. Regional syphilis rates compared for 2006, San Francisco vs. Los Angeles MSA, New York MSA, total California and total U.S.




Figure 17. Trends in regional early syphilis rates compared for 2002-2006, San Francisco vs. Los Angeles MSA, New York MSA, total California and total U.S. (MSA data not available for 2001.)




Figure 18. Early syphilis cases by sexual orientation, San Francisco, 2003-2007.




Figure 19. Primary and secondary syphilis cases by sexual orientation, San Francisco, 2003-2007.




Figure 20. *Neurosyphilis cases by case status, San Francisco, 2003-2007.*

Table 4. *Syphilis cases and rates by stage of disease, San Francisco, 2003-2007.*

	Reported cases					Rate				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
P&S SYPHILIS	336	349	249	243	202	43.3	44.9	32.1	31.3	26.0
EARLY LATENT	191	203	179	177	150	24.6	26.1	23.0	22.8	19.3
(TOTAL EARLY SYPHILIS)	527	552	428	420	352	67.8	71.1	55.1	54.1	45.3
UNKNOWN LATENT	5	8	5	9	5	0.6	1.0	0.6	1.2	0.6
LATE LATENT	110	122	78	105	112	14.2	15.7	10.0	13.5	14.4
NEUROSYPHILIS	19	37	18	9	10	2.4	4.8	2.3	1.2	1.3

Table 5. *Early syphilis cases by health care provider, San Francisco, 2003-2007.*

	Reported cases					Percent of reports				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
OOJ PROVIDERS	12	14	9	12	8	2.2%	2.5%	2.1%	2.8%	2.2%
CITY CLINIC	168	181	100	110	85	31.8%	32.7%	23.3%	26.1%	24.1%
PUBLIC CLINIC (CHN)	23	24	14	15	10	4.3%	4.3%	3.2%	3.5%	2.8%
JAILS	5	4	3	2	0	0.9%	0.7%	0.7%	0.4%	0.0%
PRIVATE CLINIC/PMD	174	136	108	142	95	33.0%	24.6%	25.2%	33.8%	26.9%
PRIVATE HOSPITAL	85	104	97	85	88	16.1%	18.8%	22.6%	20.2%	25.0%
CHPY	0	1	2	1	1	0.0%	0.1%	0.4%	0.2%	0.2%
SFGH	54	40	49	26	24	10.2%	7.2%	11.4%	6.1%	6.8%
OUTREACH	1	2	2	0	1	0.1%	0.3%	0.4%	0.0%	0.2%
MAGNET	5	46	44	27	40	0.9%	8.3%	10.2%	6.4%	11.3%
(ALL PROVIDERS)	527	552	428	420	352	100%	100%	100%	100%	100%

*Table 6. Percentage of syphilis cases by stage of disease, San Francisco, 2003-2007.*

	2003	2004	2005	2006	2007
P&S OF EARLY	63.8%	63.2%	58.2%	57.9%	57.4%
EARLY OF TOTAL	82.1%	80.9%	83.8%	78.7%	75.1%
P&S OF TOTAL	52.3%	51.2%	48.7%	45.5%	43.1%
EARLY OF LATENT	62.4%	61.0%	68.3%	60.8%	56.2%

*Table 7. Syphilis cases by sexual orientation, San Francisco, 2003-2007.*

## Cases of P&amp;S SYPHILIS

	Cases					Percent				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
WOMEN	3	0	3	2	3	0.9%	0.0%	1.2%	0.8%	1.5%
GAY/BI MEN	318	323	217	217	181	94.6%	92.6%	87.1%	89.3%	89.6%
OTHER MEN	14	24	29	23	13	4.2%	6.9%	11.6%	9.5%	6.4%
UNKNOWN MEN	1	2	0	1	5	0.3%	0.6%	0.0%	0.4%	2.5%

## Cases of EARLY SYPHILIS


	Cases					Percent				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
WOMEN	10	3	8	4	5	1.9%	0.5%	1.9%	1.0%	1.4%
GAY/BI MEN	476	502	376	368	315	90.3%	90.9%	87.9%	87.6%	89.5%
OTHER MEN	40	45	43	45	24	7.6%	8.2%	10.0%	10.7%	6.8%
UNKNOWN MEN	1	2	1	3	8	0.2%	0.4%	0.2%	0.7%	2.3%



*Table 8. Neurosyphilis cases by case status, San Francisco, 2003-2007.*

	Reported cases				
	2003	2004	2005	2006	2007
CONFIRMED	12	15	9	6	9
PROBABLE	5	14	5	1	1
SUSPECTED	2	8	4	2	0
(TOTAL)	19	37	18	9	10

## Gender

- Chlamydia rates were nearly the same for males and females in 2007, consistent with trends in prior years. The high rate of chlamydia among men in San Francisco is largely due to the availability of nucleic acid amplification testing for chlamydia at rectal and pharyngeal sites for men who have sex with men.
- The gonorrhea rate among males in 2007 was 6.2 times higher than the rate among females, consistent with trends in prior years. The gonorrhea rates in San Francisco are largely driven by a high burden of disease among men who have sex with men.
- Over 98% of reported syphilis in San Francisco is among men, with over 90% of total cases among men who have sex with men.





*San Francisco Department of Public Health*

---

*Table 9. STD cases and rates by disease and gender, San Francisco, 2003-2007.*

Cases of CHLAMYDIA

	Reported cases					Incidence rate				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
(BOTH SEXES)	3,350	3,661	3,710	4,052	3,937	431.3	471.3	477.6	521.7	506.9
FEMALE	1,680	1,781	1,768	2,058	1,875	439.9	466.3	462.9	538.9	491.0
MALE	1,648	1,853	1,881	1,931	1,999	417.4	469.3	476.4	489.1	506.3

Cases of GONORRHEA

	Reported cases					Incidence rate				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
(BOTH SEXES)	1,795	2,151	2,411	2,465	2,009	231.1	276.9	310.4	317.4	258.6
FEMALE	250	233	349	333	271	65.5	61.0	91.4	87.2	71.0
MALE	1,539	1,912	2,054	2,120	1,723	389.8	484.3	520.2	536.9	436.4

Cases of EARLY SYPHILIS

	Reported cases					Incidence rate				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
(BOTH SEXES)	527	552	428	420	352	67.8	71.1	55.1	54.1	45.3
FEMALE	10	3	8	4	5	2.6	0.8	2.1	1.0	1.3
MALE	517	549	420	416	347	130.9	139.0	106.4	105.4	87.9

*Table 10. Male/female ratios by disease, San Francisco, 2003-2007.*

	Male/female ratio				
	2003	2004	2005	2006	2007
<b>Cases of</b>					
CHLAMYDIA	0.95	1.01	1.03	0.91	1.03
GONORRHEA	5.95	7.94	5.69	6.16	6.15
EARLY SYPHILIS	50.01	177.01	50.78	100.60	67.13

### Race and ethnicity

- Rates of reported STDs vary substantially by race/ethnicity, with African Americans disproportionately affected by chlamydia and gonorrhea.
- The rate of chlamydia among African Americans is 4.6 times the chlamydia rate among whites. When restricted to females, the African American rate is over 10 times higher than the rate among white females. Following an increase from 2005 to 2006, chlamydia rates among African Americans declined 11.2% in 2007.
- The rate of gonorrhea among African Americans in 2007 was 2.4 times higher than the rate among whites. When restricted to females, the African American rate was over 11 times higher than the rate among white females. Following an increase from 2005 to 2006, gonorrhea rates among African Americans declined 22.3% in 2007.
- In 2007, the early syphilis rate was highest among Native Americans; however, only 4 cases of syphilis were reported among this racial/ethnic group, making this rate unstable. The early syphilis rates among whites, African Americans, and Hispanics were similar. Asian/Pacific Islanders had the lowest early syphilis rate in 2007, consistent with trends seen in prior years.




Figure 24. Ratio of rates among African Americans to rates among whites by disease, San Francisco, 2003-2007.




Figure 25. Trends in chlamydia rates by race/ethnicity, San Francisco, 2003-2007.




Figure 26. Chlamydia rates by race/ethnicity, San Francisco, 2007.




Figure 27. Trends in gonorrhea rates by race/ethnicity, San Francisco, 2003-2007.




Figure 28. Gonorrhea rates by race/ethnicity, San Francisco, 2007.




Figure 29. Trends in early syphilis rates by race/ethnicity, San Francisco, 2003-2007.




Figure 30. Early syphilis rates by race/ethnicity, San Francisco, 2007.




Figure 31. Chlamydia rates compared for males and females by race/ethnicity, San Francisco, 2007.




Figure 32. Gonorrhea rates compared for males and females by race/ethnicity, San Francisco, 2007.




Figure 33. Ratio of chlamydia rate among African Americans to rate among whites by gender, San Francisco, 2003-2007.




Figure 34. Ratio of gonorrhea rate among African Americans to rate among whites by gender, San Francisco, 2003-2007.

Table 11. STD cases and rates by disease and race/ethnicity, San Francisco, 2003-2007.

## Cases of CHLAMYDIA

	Reported cases					Incidence rate				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
(ALL RACES)	3,350	3,661	3,710	4,052	3,937	431.3	471.3	477.6	521.7	506.9
ASIAN/PACIFIC ISLANDER	490	470	492	486	540	194.2	186.2	194.9	192.6	214.0
AFRICAN AMERICAN	829	884	854	1,007	894	1293.9	1379.7	1332.9	1571.7	1395.3
HISPANIC	534	582	608	646	586	487.7	531.5	555.2	589.9	535.1
NATIVE AMERICAN	28	16	26	15	24	614.2	351.0	570.3	329.0	526.4
WHITE	787	910	970	1,043	1,038	232.2	268.5	286.2	307.8	306.3

## Cases of GONORRHEA

	Reported cases					Incidence rate				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
(ALL RACES)	1,795	2,151	2,411	2,465	2,009	231.1	276.9	310.4	317.4	258.6
ASIAN/PACIFIC ISLANDER	129	144	161	172	134	51.1	57.1	63.8	68.2	53.1
AFRICAN AMERICAN	328	363	515	543	422	511.9	566.6	803.8	847.5	658.7
HISPANIC	222	267	335	340	280	202.7	243.8	305.9	310.5	255.7
NATIVE AMERICAN	9	19	24	10	11	197.4	416.8	526.4	219.3	241.3
WHITE	901	1,046	1,075	1,169	951	265.9	308.6	317.2	344.9	280.6

## Cases of EARLY SYPHILIS

	Reported cases					Incidence rate				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
(ALL RACES)	527	552	428	420	352	67.8	71.1	55.1	54.1	45.3
ASIAN/PACIFIC ISLANDER	40	48	35	27	24	15.8	19.0	13.9	10.7	9.5
AFRICAN AMERICAN	34	50	28	51	35	53.1	78.0	43.7	79.6	54.6
HISPANIC	114	113	89	74	56	104.1	103.2	81.3	67.6	51.1
NATIVE AMERICAN	2	6	7	6	4	43.9	131.6	153.5	131.6	87.7
WHITE	330	318	257	245	223	97.4	93.8	75.8	72.3	65.8

Table 12. Ratio of STD rates among African Americans to rates among whites, San Francisco, 2003-2007.

		Black/white ratio				
		2003	2004	2005	2006	2007
CHLAMYDIA	(ALL PATIENTS)	5.57	5.14	4.66	5.11	4.56
	WOMEN	13.55	9.76	9.42	11.29	10.41
	MEN	3.32	3.49	3.03	3.03	2.66
GONORRHEA	(ALL PATIENTS)	1.93	1.84	2.53	2.46	2.35
	WOMEN	14.93	10.52	12.65	20.28	11.01
	MEN	1.39	1.51	1.98	1.81	1.87
EARLY SYPHILIS	(ALL PATIENTS)	0.54	0.83	0.58	1.10	0.83
	WOMEN	4.93	4.93	1.64	9.87	(N/A)
	MEN	0.57	0.87	0.60	1.14	0.84

San Francisco Department of Public Health

Table 13. STD cases and rates by disease, gender, and race/ethnicity, San Francisco, 2003-2007.

Cases of CHLAMYDIA

	2003	2004	2005	2006	2007	Reported cases		Incidence rate			
						2003	2004	2005	2006	2007	
FEMALE (ALL RACES)	1,680	1,778	1,767	2,054	1,875	439.9	465.6	462.7	537.8	491.1	
ASIAN/PACIFIC ISLANDER	307	282	311	331	325	231.2	212.4	234.3	249.3	244.5	
AFRICAN AMERICAN	465	459	456	583	514	1437.4	1418.9	1409.6	1802.2	1589.6	
HISPANIC	268	303	303	325	262	518.9	586.7	586.7	629.2	507.6	
NATIVE AMERICAN	16	9	17	10	14	766.4	431.1	814.3	479.0	676.2	
WHITE	171	232	237	250	220	107.2	145.4	148.5	156.7	137.7	
MALE (ALL RACES)	1,648	1,854	1,877	1,929	1,974	417.4	469.6	475.4	488.6	500.0	
ASIAN/PACIFIC ISLANDER	178	185	174	147	222	148.8	154.7	145.5	122.9	185.5	
AFRICAN AMERICAN	360	415	385	415	347	1134.9	1308.3	1213.7	1308.3	1093.5	
HISPANIC	262	277	299	306	339	452.9	478.8	516.8	528.9	585.6	
NATIVE AMERICAN	12	7	9	5	10	485.6	283.2	364.2	202.3	408.0	
WHITE	611	664	716	774	720	340.7	370.3	399.3	431.6	401.5	

Cases of GONORRHEA


	2003	2004	2005	2006	2007	Reported cases		Incidence rate			
						2003	2004	2005	2006	2007	
FEMALE (ALL RACES)	250	233	351	333	284	65.5	61.0	91.9	87.2	74.5	
ASIAN/PACIFIC ISLANDER	22	17	18	29	30	16.6	12.8	13.6	21.8	22.8	
AFRICAN AMERICAN	114	95	162	183	135	352.4	293.7	500.8	565.7	417.7	
HISPANIC	20	26	35	27	30	38.7	50.3	67.8	52.3	58.6	
NATIVE AMERICAN	2	5	9	2	8	95.8	239.5	431.1	95.8	386.4	
WHITE	38	44	63	46	48	23.8	27.6	39.5	28.8	30.3	
MALE (ALL RACES)	1,539	1,912	2,054	2,121	1,805	389.8	484.3	520.2	537.2	457.1	
ASIAN/PACIFIC ISLANDER	105	129	143	142	125	87.8	107.9	119.6	118.7	104.5	
AFRICAN AMERICAN	210	268	354	355	319	662.0	844.9	1116.0	1119.1	1004.4	
HISPANIC	200	236	295	309	252	345.7	407.9	509.9	534.1	435.7	
NATIVE AMERICAN	7	14	15	8	6	283.2	566.5	606.9	323.7	244.8	
WHITE	858	987	1,007	1,112	901	478.5	550.4	561.5	620.1	502.7	

Cases of EARLY SYPHILIS

	2003	2004	2005	2006	2007	Reported cases		Incidence rate			
						2003	2004	2005	2006	2007	
FEMALE (ALL RACES)	10	3	8	4	8	2.6	0.8	2.1	1.0	2.1	
ASIAN/PACIFIC ISLANDER	1	0	2	0	0	0.8	0.0	1.5	0.0	0.0	
AFRICAN AMERICAN	1	1	1	2	4	3.1	3.1	3.1	6.2	12.5	
HISPANIC	7	1	2	1	4	13.6	1.9	3.9	1.9	7.8	
NATIVE AMERICAN	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	
WHITE	1	1	3	1	0	0.6	0.6	1.9	0.6	0.0	
MALE (ALL RACES)	517	548	419	416	363	130.9	138.8	106.1	105.4	91.9	
ASIAN/PACIFIC ISLANDER	39	48	34	27	22	32.6	40.1	28.4	22.6	18.5	
AFRICAN AMERICAN	33	48	27	49	38	104.0	151.3	85.1	154.5	120.8	
HISPANIC	107	110	86	73	58	184.9	190.1	148.6	126.2	101.1	
NATIVE AMERICAN	2	5	6	6	2	80.9	202.3	242.8	242.8	81.6	
WHITE	327	319	254	244	230	182.3	177.9	141.6	136.1	128.2	

## Age

- Rates of STDs were highly age-dependent and varied by gender.
- Rates of chlamydia were highest among adolescents and young adults (aged 15-24) and decreased dramatically among older age groups. Among females, chlamydia rates were highest for those ages 15-19; for males, rates were highest among the 20-24-year-old age group.
- Gonorrhea in San Francisco affected two key populations: adolescents and older men who have sex with men. As a result, the age distribution for males was bimodal with two age groups largely affected. For females, gonorrhea rates were highest in the 15-19-year-old age category. Among males, younger males aged 20-24 (representing more heterosexual transmission) and older males aged 35-44 (representing more MSM transmission) had the highest rates.
- Syphilis rates peaked among 40-44-year-old males, consistent with trends seen in prior years.






Figure 36. Age-specific chlamydia rates by gender, San Francisco, 2007.




Figure 37. Trends in chlamydia rates for selected age groups, San Francisco, 2003-2007.




Figure 38. Age-specific gonorrhea rates by gender, San Francisco, 2007




Figure 39. Trends in gonorrhea rates for selected age groups, San Francisco, 2003-2007.




Figure 40. Age-specific early syphilis rates by gender, San Francisco, 2007.




Figure 41. Trends in early syphilis rates for selected age groups, San Francisco, 2003-2007.




Figure 42. Trends in male age-specific rates by disease, San Francisco, 2003-2007.




Figure 43. Trends in female age-specific rates by disease, San Francisco, 2003-2007.

San Francisco Department of Public Health

Table 14. STD cases and rates by disease, gender and age group, San Francisco, 2003-2007.

Cases of CHLAMYDIA

	Reported cases					Incidence rate					
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007	
(BOTH GENDERS) (ALL)	3,350	3,663	3,710	4,053	3,939	431.3	471.6	477.6	521.8	507.1	
15-19 YRS	714	633	671	825	693	2142.0	1899.0	2013.0	2475.0	2079.0	
20-24 YRS	885	997	983	967	995	1578.8	1778.6	1753.7	1725.1	1775.1	
25-29 YRS	602	665	686	776	710	658.7	727.6	750.6	849.1	776.9	
30-34 YRS	421	487	438	435	454	472.9	547.0	492.0	488.6	510.0	
35-39 YRS	310	381	380	408	370	428.6	526.8	525.4	564.1	511.6	
40-44 YRS	196	247	243	302	286	318.8	401.8	395.3	491.2	465.2	
45-54 YRS	146	157	206	229	307	135.5	145.8	191.2	212.6	285.0	
55-64 YRS	25	39	52	44	65	38.3	59.7	79.7	67.4	99.6	
65+ YRS	6	3	6	8	5	5.7	2.8	5.7	7.5	4.7	
FEMALE	(ALL)	1,680	1,781	1,768	2,058	1,875	439.9	466.3	462.9	538.9	491.0
15-19 YRS	539	461	494	609	505	3304.7	2826.5	3028.8	3733.9	3096.3	
20-24 YRS	564	630	587	619	645	1989.6	2222.5	2070.8	2183.7	2275.4	
25-29 YRS	276	328	325	401	348	621.3	738.4	731.6	902.7	783.4	
30-34 YRS	136	155	155	169	153	335.7	382.6	382.6	417.1	377.6	
35-39 YRS	55	72	83	96	78	172.1	225.3	259.8	300.5	244.1	
40-44 YRS	39	47	44	62	39	139.5	168.2	157.4	221.8	139.5	
45-54 YRS	30	40	33	44	58	58.2	77.5	64.0	85.3	112.4	
55-64 YRS	7	9	14	10	13	21.1	27.1	42.2	30.1	39.2	
65+ YRS	0	0	3	1	1	0.0	0.0	4.9	1.6	1.6	
MALE	(ALL)	1,648	1,855	1,881	1,932	2,001	417.4	469.8	476.4	489.3	506.8
15-19 YRS	167	163	149	185	148	981.0	957.5	875.2	1086.7	869.4	
20-24 YRS	317	360	377	337	339	1144.1	1299.3	1360.7	1216.3	1223.5	
25-29 YRS	322	333	357	363	356	685.5	709.0	760.1	772.8	757.9	
30-34 YRS	281	331	279	262	301	579.3	682.3	575.1	540.1	620.5	
35-39 YRS	255	308	295	311	290	631.6	762.8	730.6	770.3	718.2	
40-44 YRS	157	200	199	240	247	468.3	596.6	593.6	715.9	736.8	
45-54 YRS	116	117	172	184	249	206.6	208.4	306.4	327.8	443.6	
55-64 YRS	18	29	38	34	52	56.1	90.4	118.4	106.0	162.0	
65+ YRS	6	3	3	7	4	13.5	6.8	6.8	15.8	9.0	

Cases of GONORRHEA

	Reported cases					Incidence rate					
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007	
(BOTH GENDERS) (ALL)	1,795	2,153	2,411	2,468	2,010	231.1	277.2	310.4	317.7	258.8	
15-19 YRS	106	106	170	167	120	318.0	318.0	510.0	501.0	360.0	
20-24 YRS	228	317	361	367	264	406.8	565.5	644.0	654.7	471.0	
25-29 YRS	317	346	377	456	337	346.9	378.6	412.5	498.9	368.7	
30-34 YRS	341	406	366	343	330	383.0	456.1	411.1	385.3	370.7	
35-39 YRS	353	398	475	415	315	488.1	550.3	656.7	573.8	435.5	
40-44 YRS	232	279	316	363	304	377.4	453.8	514.0	590.5	494.5	
45-54 YRS	173	239	275	285	257	160.6	221.9	255.3	264.6	238.6	
55-64 YRS	25	46	52	45	62	38.3	70.5	79.7	68.9	95.0	
65+ YRS	11	5	10	9	5	10.4	4.7	9.4	8.5	4.7	
FEMALE	(ALL)	250	233	349	333	271	65.5	61.0	91.4	87.2	71.0
15-19 YRS	71	67	108	110	69	435.3	410.8	662.2	674.4	423.1	
20-24 YRS	63	75	98	96	69	222.2	264.6	345.7	338.7	243.4	
25-29 YRS	52	32	53	58	50	117.1	72.0	119.3	130.6	112.6	
30-34 YRS	24	18	26	18	33	59.2	44.4	64.2	44.4	81.5	
35-39 YRS	11	10	21	15	19	34.4	31.3	65.7	46.9	59.5	
40-44 YRS	9	13	18	6	9	32.2	46.5	64.4	21.5	32.2	
45-54 YRS	9	11	14	12	9	17.4	21.3	27.1	23.3	17.4	
55-64 YRS	2	2	5	2	4	6.0	6.0	15.1	6.0	12.1	
65+ YRS	2	0	0	1	0	3.2	0.0	0.0	1.6	0.0	


## Cases of GONORRHEA

		Reported cases					Incidence rate				
		2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
MALE	(ALL)	1,539	1,914	2,054	2,123	1,724	389.8	484.8	520.2	537.7	436.6
	15-19 YRS	34	37	61	54	43	199.7	217.3	358.3	317.2	252.6
	20-24 YRS	165	242	261	271	194	595.5	873.4	942.0	978.1	700.2
	25-29 YRS	265	314	324	396	286	564.2	668.5	689.8	843.1	608.9
	30-34 YRS	312	388	338	324	295	643.2	799.8	696.8	667.9	608.1
	35-39 YRS	342	387	453	396	296	847.0	958.5	1122.0	980.8	733.1
	40-44 YRS	223	265	297	357	295	665.2	790.5	885.9	1064.9	879.9
	45-54 YRS	164	228	260	271	247	292.2	406.2	463.2	482.8	440.0
	55-64 YRS	23	44	47	43	57	71.7	137.1	146.5	134.0	177.6
	65+ YRS	9	5	10	8	5	20.3	11.3	22.5	18.0	11.3

## Cases of EARLY SYPHILIS

		Reported cases					Incidence rate				
		2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
(BOTH GENDERS)	(ALL)	527	552	428	420	352	67.8	71.1	55.1	54.1	45.3
	15-19 YRS	3	0	4	3	3	9.0	0.0	12.0	9.0	9.0
	20-24 YRS	25	27	15	20	17	44.6	48.2	26.8	35.7	30.3
	25-29 YRS	49	62	39	40	34	53.6	67.8	42.7	43.8	37.2
	30-34 YRS	123	97	66	54	37	138.2	109.0	74.1	60.7	41.6
	35-39 YRS	136	126	102	99	69	188.0	174.2	141.0	136.9	95.4
	40-44 YRS	96	136	94	88	74	156.2	221.2	152.9	143.1	120.4
	45-54 YRS	76	82	92	87	95	70.6	76.1	85.4	80.8	88.2
	55-64 YRS	15	21	14	24	22	23.0	32.2	21.4	36.8	33.7
	65+ YRS	1	1	2	5	1	0.9	0.9	1.9	4.7	0.9
FEMALE	(ALL)	10	3	8	4	5	2.6	0.8	2.1	1.0	1.3
	15-19 YRS	1	0	1	0	0	6.1	0.0	6.1	0.0	0.0
	20-24 YRS	3	1	1	2	1	10.6	3.5	3.5	7.1	3.5
	25-29 YRS	2	0	0	0	1	4.5	0.0	0.0	0.0	2.3
	30-34 YRS	1	1	0	0	0	2.5	2.5	0.0	0.0	0.0
	35-39 YRS	1	0	3	0	1	3.1	0.0	9.4	0.0	3.1
	40-44 YRS	1	1	0	1	1	3.6	3.6	0.0	3.6	3.6
	45-54 YRS	0	0	3	1	1	0.0	0.0	5.8	1.9	1.9
	55-64 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0
	65+ YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0
MALE	(ALL)	517	549	420	416	347	130.9	139.0	106.4	105.4	87.9
	15-19 YRS	2	0	3	3	3	11.7	0.0	17.6	17.6	17.6
	20-24 YRS	22	26	14	18	16	79.4	93.8	50.5	65.0	57.7
	25-29 YRS	47	62	39	40	33	100.1	132.0	83.0	85.2	70.3
	30-34 YRS	122	96	66	54	37	251.5	197.9	136.1	111.3	76.3
	35-39 YRS	135	126	99	99	68	334.4	312.1	245.2	245.2	168.4
	40-44 YRS	95	135	94	87	73	283.4	402.7	280.4	259.5	217.7
	45-54 YRS	76	82	89	86	94	135.4	146.1	158.5	153.2	167.5
	55-64 YRS	15	21	14	24	22	46.7	65.4	43.6	74.8	68.6
	65+ YRS	1	1	2	5	1	2.3	2.3	4.5	11.3	2.3

## Geography



geographic analyses, but are counted as San Francisco morbidity and included in all other citywide analyses. Cases among homeless patients were also excluded. Note that the rate obtained from combining the rates for each neighborhood will therefore be lower than the actual overall rate for the city.

- San Francisco is divided into 39 districts defined by the Department of Public Health Planning Office (see Figure 46)
- High rates of chlamydia continue to be concentrated in the southeastern sector of the city. Sunnydale and West Hunter's Point had the highest chlamydia rates in 2007. Additionally, chlamydia rates in the Castro, Duboce Triangle, and Western Addition neighborhoods were also disproportionately high. These geographic data likely reflect the two overlapping chlamydia epidemics occurring among adolescents (southeastern sector of the city) and men who have sex with men (Castro area).
- Gonorrhea rates in San Francisco in 2007 were highest in the Castro and Duboce Triangle neighborhoods, with both having a rate in excess of 1,300 cases per 100,000 residents. Rates of gonorrhea in the southeastern sector of the city declined in 2007. Female gonorrhea continues to concentrate in the Sunnydale, West Hunter's Point, and Potrero Hill neighborhoods.
- Syphilis continues to concentrate in the Castro and Duboce Triangle neighborhoods.
- The proportion of reported cases of chlamydia and early syphilis that were homeless declined in 2007, while the proportion increased slightly for gonorrhea. Homeless residents account for less than 3% of total reported STD morbidity.

To examine the geographic distribution of STD cases and to compare disease trends in different neighborhoods, addresses of cases have been grouped into the thirty-nine districts defined by the planner's office of the Department of Public Health (see Figure 45).

The census tract for each address was determined using a computer program. Some addresses were vague or not recognized as valid San Francisco addresses, and therefore could not be assigned to a neighborhood.

Although the law requires addresses to be included in STD reports, they are often missing. In 2007, approximately 14 percent of all gonorrhea and chlamydia reports were missing addresses. Cases that were missing addresses or were otherwise unable to be assigned to a neighborhood are not included in these




Figure 46. Chlamydia rates by neighborhood for San Francisco, 2007.




Figure 47. Chlamydia trends for selected neighborhoods, San Francisco, 2003-2007 (1 of 2).




Figure 48. Chlamydia trends for selected neighborhoods, San Francisco, 2003-2007 (2 of 2).




Figure 49. Gonorrhea rates by neighborhood for San Francisco, 2007.




Figure 50. Gonorrhea trends for selected neighborhoods, San Francisco, 2003-2007 (1 of 2).




Figure 51. Gonorrhea trends for selected neighborhoods, San Francisco, 2003-2007 (2 of 2).




Figure 52. Male early syphilis rates by neighborhood for San Francisco, 2007.




Figure 53. Early syphilis trends for selected neighborhoods, San Francisco, 2003-2007.




Figure 54. Chlamydia rates for San Francisco by neighborhood, 2007.




Figure 55. Male chlamydia rates by neighborhood for San Francisco, 2007.




Figure 56. Female chlamydia rates by neighborhood for San Francisco, 2007.




Figure 57. Male and female chlamydia rates compared by neighborhood, 2007.




Figure 58. Gonorrhea rates for San Francisco by neighborhood, 2007.




Figure 59. Male gonorrhea rates by neighborhood for San Francisco, 2007.




Figure 60. Female gonorrhea rates by neighborhood for San Francisco, 2007.




Figure 61. Male and female gonorrhea rates compared by neighborhood, 2007.




Figure 62. Male and female early syphilis rates for San Francisco by neighborhood, 2007.




Figure 63. Trends in homeless cases reported among persons with an STD by disease for San Francisco, 2003-2007.

Table 15. STD cases among homeless patients and percent of STD cases that are homeless, San Francisco, 2003-2007.

	Homeless cases					Percent homeless				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
CHLAMYDIA	51	46	86	76	51	1.5%	1.3%	2.3%	1.9%	1.3%
GONORRHEA	26	42	70	55	53	1.4%	2.0%	2.9%	2.2%	2.6%
EARLY SYPHILIS	1	9	3	10	6	0.2%	1.6%	0.7%	2.4%	1.7%

Table 16. STD cases and rates by neighborhood, San Francisco, 2003-2007.

#### Cases of CHLAMYDIA

	Reported cases					Incidence rate				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
(WHOLE CITY)	3,350	3,661	3,710	4,052	3,937	431.3	471.3	477.6	521.7	506.9
SEACLIFF	2	2	1	3	1	81.6	81.6	40.8	122.4	40.8
PACIFIC HTS	5	8	8	10	11	58.3	93.2	93.2	116.5	128.2
W TWIN PEAKS	35	39	23	32	39	161.6	180.0	106.2	147.7	180.0
SUNSET/PARKSIDE	109	122	164	186	171	127.9	143.2	192.5	218.3	200.7
WESTWOOD PK	11	13	27	18	28	110.9	131.0	272.1	181.4	282.2
DIAMOND HTS	21	25	19	39	35	261.9	311.8	236.9	486.3	436.5
LAKE MERCED	2	2	3	4	3	131.8	131.8	197.6	263.5	197.6
MARINA	41	50	70	69	64	103.7	126.4	177.0	174.5	161.8
NORTH BEACH	24	32	40	51	46	129.4	172.5	215.6	274.9	247.9
USF/LAUREL HTS	54	57	50	68	67	252.0	266.0	233.3	317.3	312.7
NOB HILL	23	37	48	53	59	158.6	255.1	331.0	365.5	406.8
DUBOCE TRIANGLE	36	29	27	31	45	1202.0	968.3	901.5	1035.1	1502.5
HAIGHT DISTRICT	58	81	85	93	102	360.6	503.6	528.5	578.3	634.2
CASTRO	121	183	175	199	190	700.5	1059.4	1013.1	1152.0	1099.9
GUERRERO	39	40	38	42	52	390.5	400.6	380.5	420.6	520.7
NOE VALLEY	50	33	38	49	40	403.2	266.1	306.4	395.1	322.5

## Cases of CHLAMYDIA

	Reported cases					Incidence rate				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
INNER SUNSET	26	36	30	39	43	146.8	203.3	169.4	220.3	242.9
PARK MERCED	26	40	36	49	41	251.6	387.1	348.4	474.2	396.8
GLEN PK	28	28	52	51	38	182.0	182.0	337.9	331.4	247.0
BERNAL HTS	129	112	124	110	102	517.0	448.9	497.0	440.8	408.8
ALEMANY	168	158	173	209	205	291.8	274.5	300.5	363.0	356.1
PORTOLA	44	35	60	51	52	286.3	227.7	390.4	331.8	338.3
VISITACION VLY	53	76	83	104	86	318.5	456.7	498.8	625.0	516.8
BAYVIEW	79	67	86	109	93	563.0	477.5	612.9	776.8	662.8
NORTH MISSION	84	77	94	110	102	587.2	538.3	657.1	769.0	713.0
MISSION	173	162	196	179	178	481.7	451.1	545.8	498.5	495.7
POTRERO PT	77	91	117	114	79	662.1	782.5	1006.1	980.3	679.3
OMI	112	101	130	135	118	491.2	443.0	570.2	592.1	517.5
PRESIDO	5	4	6	7	5	223.8	179.1	268.6	313.3	223.8
RICHMOND	96	114	103	129	134	145.3	172.5	155.9	195.2	202.8
CHINATOWN	51	57	43	56	86	177.1	197.9	149.3	194.5	298.6
WESTERN ADDITION	268	299	314	358	302	689.6	769.3	807.9	921.1	777.0
W HUNTER'S PT	229	257	211	307	268	1272.8	1428.4	1172.7	1706.3	1489.6
SUNNYDALE	48	48	56	60	62	1253.9	1253.9	1462.9	1567.4	1619.6
DOWNTOWN/TENDERLOIN	186	216	239	254	240	485.2	563.4	623.4	662.6	626.0
CATHEDRAL HILL	48	48	52	58	64	393.4	393.4	426.2	475.3	524.5
SOUTH-OF-MARKET	92	96	110	111	132	464.4	484.6	555.2	560.3	666.3
LINCOLN PARK/VA HOSPITAL	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0
GOLDEN GATE PK	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0

## Cases of GONORRHEA

	Reported cases					Incidence rate				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
(WHOLE CITY)	1,795	2,151	2,411	2,465	2,009	231.1	276.9	310.4	317.4	258.6
SEACLIFF	2	0	0	3	0	81.6	0.0	0.0	122.4	0.0
PACIFIC HTS	8	7	3	4	6	93.2	81.6	35.0	46.6	69.9
W TWIN PEAKS	15	25	22	24	19	69.2	115.4	101.6	110.8	87.7
SUNSET/PARKSIDE	32	40	52	75	27	37.6	46.9	61.0	88.0	31.7
WESTWOOD PK	9	11	14	10	7	90.7	110.9	141.1	100.8	70.5
DIAMOND HTS	20	25	19	28	27	249.4	311.8	236.9	349.2	336.7
LAKE MERCED	0	0	0	1	0	0.0	0.0	0.0	65.9	0.0
MARINA	25	20	31	28	23	63.2	50.6	78.4	70.8	58.2
NORTH BEACH	17	14	20	23	26	91.6	75.5	107.8	124.0	140.1
USF/LAUREL HTS	31	26	30	24	33	144.7	121.3	140.0	112.0	154.0
NOB HILL	21	27	28	35	33	144.8	186.2	193.1	241.3	227.6
DUBOCE TRIANGLE	47	39	70	63	56	1569.3	1302.2	2337.2	2103.5	1869.8
HAIGHT DISTRICT	56	85	93	100	83	348.2	528.5	578.3	621.8	516.1
CASTRO	201	261	250	281	236	1163.6	1510.9	1447.3	1626.7	1366.2
GUERRERO	37	49	44	50	34	370.5	490.7	440.6	500.7	340.5
NOE VALLEY	53	48	53	50	34	427.4	387.0	427.4	403.2	274.1
INNER SUNSET	12	19	25	26	17	67.8	107.3	141.2	146.8	96.0
PARK MERCED	10	14	11	23	17	96.8	135.5	106.5	222.6	164.5
GLEN PK	28	26	45	49	27	182.0	169.0	292.5	318.5	175.5
BERNAL HTS	38	53	69	37	44	152.3	212.4	276.5	148.3	176.3
ALEMANY	38	49	63	61	45	66.0	85.1	109.4	106.0	78.2
PORTOLA	11	9	14	10	17	71.6	58.6	91.1	65.1	110.6
VISITACION VLY	17	14	29	25	19	102.2	84.1	174.3	150.2	114.2
BAYVIEW	16	17	30	34	31	114.0	121.2	213.8	242.3	220.9
NORTH MISSION	73	75	70	102	69	510.3	524.3	489.3	713.0	482.3
MISSION	71	83	114	115	64	197.7	231.1	317.5	320.2	178.2
POTRERO PT	39	38	71	76	51	335.4	326.8	610.5	653.5	438.6
OMI	29	28	49	48	44	127.2	122.8	214.9	210.5	193.0
PRESIDO	5	2	2	1	3	223.8	89.5	89.5	44.8	134.3
RICHMOND	35	36	47	42	37	53.0	54.5	71.1	63.6	56.0

*San Francisco Department of Public Health*

Cases of GONORRHEA

	Reported cases					Incidence rate				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
CHINATOWN	15	25	19	17	24	52.1	86.8	66.0	59.0	83.3
WESTERN ADDITION	223	239	268	273	217	573.8	614.9	689.6	702.4	558.3
W HUNTER'S PT	61	72	85	106	88	339.0	400.2	472.4	589.2	489.1
SUNNYDALE	9	12	6	19	18	235.1	313.5	156.7	496.3	470.2
DOWNTOWN/TENDERLOIN	155	196	225	216	182	404.3	511.3	586.9	563.4	474.7
CATHEDRAL HILL	23	22	47	49	32	188.5	180.3	385.2	401.6	262.3
SOUTH-OF-MARKET	70	99	108	112	93	353.3	499.7	545.1	565.3	469.4
LINCOLN PARK/VA HOSPITAL	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0
GOLDEN GATE PK	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0

Cases of EARLY SYPHILIS

	Reported cases					Incidence rate				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
(WHOLE CITY)	527	552	428	420	352	67.8	71.1	55.1	54.1	45.3
SEACLIFF	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0
PACIFIC HTS	2	2	1	1	0	23.3	23.3	11.7	11.7	0.0
W TWIN PEAKS	3	3	5	3	1	13.8	13.8	23.1	13.8	4.6
SUNSET/PARKSIDE	4	8	11	4	7	4.7	9.4	12.9	4.7	8.2
WESTWOOD PK	7	8	4	3	0	70.5	80.6	40.3	30.2	0.0
DIAMOND HTS	11	8	5	4	5	137.2	99.8	62.4	49.9	62.4
LAKE MERCED	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0
MARINA	4	5	8	7	4	10.1	12.6	20.2	17.7	10.1
NORTH BEACH	2	4	6	2	0	10.8	21.6	32.3	10.8	0.0
USF/LAUREL HTS	12	6	3	4	2	56.0	28.0	14.0	18.7	9.3
NOB HILL	11	3	8	7	7	75.9	20.7	55.2	48.3	48.3
DUBOCE TRIANGLE	17	21	15	18	7	567.6	701.2	500.8	601.0	233.7
HAIGHT DISTRICT	18	28	18	21	22	111.9	174.1	111.9	130.6	136.8
CASTRO	86	98	82	71	65	497.9	567.3	474.7	411.0	376.3
GUERRERO	16	16	13	6	6	160.2	160.2	130.2	60.1	60.1
NOE VALLEY	18	17	15	11	15	145.1	137.1	120.9	88.7	120.9
INNER SUNSET	6	12	7	8	2	33.9	67.8	39.5	45.2	11.3
PARK MERCED	2	3	0	1	2	19.4	29.0	0.0	9.7	19.4
GLEN PK	4	8	5	18	8	26.0	52.0	32.5	117.0	52.0
BERNAL HTS	17	14	13	15	9	68.1	56.1	52.1	60.1	36.1
ALEMANY	9	8	6	7	14	15.6	13.9	10.4	12.2	24.3
PORTOLA	2	1	2	1	1	13.0	6.5	13.0	6.5	6.5
VISITACION VLY	1	6	1	1	3	6.0	36.1	6.0	6.0	18.0
BAYVIEW	2	3	1	2	1	14.3	21.4	7.1	14.3	7.1
NORTH MISSION	25	34	22	15	26	174.8	237.7	153.8	104.9	181.8
MISSION	23	25	18	10	11	64.0	69.6	50.1	27.8	30.6
POTRERO PT	18	9	9	7	4	154.8	77.4	77.4	60.2	34.4
OMI	4	0	4	4	1	17.5	0.0	17.5	17.5	4.4
PRESIDO	0	1	0	0	1	0.0	44.8	0.0	0.0	44.8
RICHMOND	5	7	7	7	5	7.6	10.6	10.6	10.6	7.6
CHINATOWN	7	3	4	9	3	24.3	10.4	13.9	31.3	10.4
WESTERN ADDITION	67	74	51	55	37	172.4	190.4	131.2	141.5	95.2
W HUNTER'S PT	5	4	2	6	4	27.8	22.2	11.1	33.3	22.2
SUNNYDALE	0	0	0	2	1	0.0	0.0	0.0	52.2	26.1
DOWNTOWN/TENDERLOIN	50	37	42	36	40	130.4	96.5	109.6	93.9	104.3
CATHEDRAL HILL	13	7	10	3	1	106.5	57.4	82.0	24.6	8.2
SOUTH-OF-MARKET	20	31	12	21	18	100.9	156.5	60.6	106.0	90.9
LINCOLN PARK/VA HOSPITAL	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0
GOLDEN GATE PK	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0

## *Adolescents*

Chlamydia rates (per 100,000): Adolescents=1923.9; Adults (21+) = 465.4

Gonorrhea rates (per 100,000): Adolescents=341.4; Adults (21+) = 286.0

Early syphilis rates (per 100,000): Adolescents=8.4; Adults (21+) = 54.3

- Chlamydia rates among adolescents were over 4 times higher than among adults. Gonorrhea rates for adolescents were 1.2 times higher than among adults. Syphilis rates among adolescents were nearly 6.5 times lower than among adults.
- Chlamydia rates among adolescents declined by 13.2% in 2007. Adult chlamydia rates were stable from 2003-2007.
- Adolescent gonorrhea rates declined by 29.1% from 2006 to 2007; adult rates declined by 17.7%.
- For both chlamydia and gonorrhea, rates among adolescent females were significantly higher than rates among adolescent males; the trends in these rates have remained stable from 2003 through 2007.
- Racial/ethnic disparities for chlamydia and gonorrhea are large for adolescents. In 2007, the rates of chlamydia and gonorrhea among African-American adolescents were nearly 7 times that of white adolescents.
- Adolescent chlamydia is concentrated in the southeastern sector of San Francisco. While adolescent chlamydia rates declined in West Hunter's Point and Potrero Hill, a nearly 15% increase was seen in Sunnydale. For gonorrhea, increases were seen in the adolescent rates in Sunnydale, Potrero Hill and Bayview.
- A larger proportion of adolescent chlamydia and gonorrhea cases were reported by private doctors and hospitals. From 2003 through 2007, the proportion of chlamydia cases diagnosed through private doctors and hospitals increased from 55.0% to 62.2%; for gonorrhea the proportion increased from 39.0% to 47.8%.




Figure 64. STD rates for adolescents vs. adults, San Francisco, 2007.




Figure 65. Chlamydia trends for adolescents vs. adults, San Francisco, 2003-2007.




Figure 66. Gonorrhea trends for adolescents vs. adults, San Francisco, 2003-2007.




Figure 67. Early syphilis trends for adolescents vs. adults, San Francisco, 2003-2007.




Figure 68. Male/female rate ratios for adolescents and adults, San Francisco, 2007.




Figure 69. Gender-specific chlamydia rates for adolescents, San Francisco, 2003-2007.




Figure 70. Gender-specific gonorrhea rates for adolescents, San Francisco, 2003-2007.




Figure 71. Gender-specific early syphilis rates for adolescents, San Francisco, 2003-2007. Note: male/female ratio is undefined for 2004 and 2007 as there were no female cases.




Figure 72. Ratio of rates among African-American adolescents to rates among white adolescents by disease, San Francisco, 2003-2007. (Early syphilis cases not shown due to small numbers of cases.)




Figure 73. Race/ethnicity-specific chlamydia rates for adolescents compared with adults, San Francisco, 2007.




Figure 74. Race/ethnicity-specific chlamydia trends for adolescents, San Francisco, 2003-2007.




Figure 75. Race/ethnicity-specific gonorrhea rates for adolescents compared with adults, San Francisco, 2007.




Figure 76. Race/ethnicity-specific gonorrhea trends for adolescents, San Francisco, 2003-2007.




Figure 77. Race/ethnicity-specific early syphilis trends for adolescents, San Francisco, 2003-2007.




Figure 78. Adolescent and adult chlamydia rates compared by neighborhood, San Francisco, 2007.




Figure 79. Chlamydia rates for adolescents by neighborhood, San Francisco, 2007.




Figure 80. Gonorrhea rates for adolescents by neighborhood, San Francisco, 2007.




Figure 81. Adolescent and adult gonorrhea rates compared by neighborhood, San Francisco, 2007.




Figure 82. Chlamydia trends among adolescents for selected neighborhoods, San Francisco, 2003-2007.




Figure 83. Gonorrhea trends among adolescents for selected neighborhoods, San Francisco, 2003-2007.




Figure 84. Male adolescent chlamydia rates by neighborhood, San Francisco, 2007.




Figure 85. Female adolescent chlamydia rates by neighborhood, San Francisco, 2007.




Figure 86. Adolescent male and female chlamydia rates compared by neighborhood, 2007.




Figure 87. Male adolescent gonorrhea rates by neighborhood, San Francisco, 2007.




Figure 88. Female adolescent gonorrhea rates by neighborhood, San Francisco, 2007.






Figure 89. Adolescent male and female gonorrhea rates compared by neighborhood, 2007



*Figure 90. Adolescent chlamydia cases by health care provider, San Francisco, 2007. Not included: 22 cases reported by providers outside San Francisco jurisdiction.*



*Figure 91. Adolescent gonorrhea cases by health care provider, San Francisco, 2007. Not included: 4 cases reported by providers outside San Francisco jurisdiction.*




Figure 92. Trends in proportion of adolescent cases identified through public sources (i.e., all but private clinics, physicians, and hospitals), San Francisco, 2003-2007.

Table 17. Comparison of STD cases and rates for adolescents and adults, San Francisco, 2003-2007.

Cases of CHLAMYDIA

	Reported cases					Incidence rate				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
ADOLESCENT (14-20 YRS)	920	863	918	1,061	921	1926.8	1807.4	1922.6	2222.1	1928.9
14-17 YRS	361	310	314	413	341	1457.5	1251.6	1267.7	1667.4	1376.7
18-20 YRS	559	553	604	648	580	2432.7	2406.5	2628.5	2820.0	2524.0
ADULT (21+ YRS)	2,405	2,774	2,768	2,962	2,983	375.2	432.8	431.9	462.1	465.4

Cases of GONORRHEA

	Reported cases					Incidence rate				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
ADOLESCENT (14-20 YRS)	151	155	228	230	163	316.2	324.6	477.5	481.7	341.4
14-17 YRS	59	47	78	84	60	238.2	189.8	314.9	339.1	242.2
18-20 YRS	92	108	150	146	103	400.4	470.0	652.8	635.4	448.2
ADULT (21+ YRS)	1,643	1,987	2,180	2,227	1,833	256.3	310.0	340.1	347.5	286.0

Cases of EARLY SYPHILIS

	Reported cases					Incidence rate				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
ADOLESCENT (14-20 YRS)	8	1	5	8	4	16.8	2.1	10.5	16.8	8.4
14-17 YRS	1	0	0	1	0	4.0	0.0	0.0	4.0	0.0
18-20 YRS	7	1	5	7	4	30.5	4.4	21.8	30.5	17.4
ADULT (21+ YRS)	517	551	423	412	348	80.7	86.0	66.0	64.3	54.3

Table 18. Adolescent cases by disease and health care provider, San Francisco, 2003-2007.

	(ALL PROVIDERS)	Reported cases					Percent of reports				
		2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
CHLAMYDIA	(ALL PROVIDERS)	920	863	918	1,061	921	100%	100%	100%	100%	100%
	OOJ PROVIDERS	33	16	28	22	27	3.5%	1.8%	3.0%	2.0%	2.9%
	CITY CLINIC	68	62	77	65	53	7.3%	7.1%	8.3%	6.1%	5.7%
	PUBLIC CLINIC (CHN)	37	40	28	48	25	4.0%	4.6%	3.0%	4.5%	2.7%
	JAILS	108	131	113	173	113	11.7%	15.1%	12.3%	16.3%	12.2%
	PRIVATE CLINIC/PMD	295	263	334	376	351	32.0%	30.4%	36.3%	35.4%	38.1%
	PRIVATE HOSPITAL	212	189	187	236	222	23.0%	21.9%	20.3%	22.2%	24.1%
	CHPY	64	74	82	104	101	6.9%	8.5%	8.9%	9.8%	10.9%
	SFGH	92	78	61	31	25	10.0%	9.0%	6.6%	2.9%	2.7%
	OUTREACH	11	7	4	0	1	1.1%	0.8%	0.4%	0.0%	0.1%
GONORRHEA	MAGNET	0	3	4	6	3	0.0%	0.3%	0.4%	0.5%	0.3%
	(ALL PROVIDERS)	151	155	228	230	163	100%	100%	100%	100%	100%
	OOJ PROVIDERS	3	4	8	4	2	1.9%	2.5%	3.5%	1.7%	1.2%
	CITY CLINIC	21	36	32	40	28	13.9%	23.2%	14.0%	17.3%	17.1%
	PUBLIC CLINIC (CHN)	9	8	4	9	5	5.9%	5.1%	1.7%	3.9%	3.0%
	JAILS	25	16	36	38	25	16.5%	10.3%	15.7%	16.5%	15.3%
	PRIVATE CLINIC/PMD	31	24	53	45	38	20.5%	15.4%	23.2%	19.5%	23.3%
	PRIVATE HOSPITAL	28	31	43	47	40	18.5%	20.0%	18.8%	20.4%	24.5%
	CHPY	15	18	22	26	16	9.9%	11.6%	9.6%	11.3%	9.8%
	SFGH	18	13	24	9	2	11.9%	8.3%	10.5%	3.9%	1.2%
EARLY SYPHILIS	OUTREACH	1	0	2	1	0	0.6%	0.0%	0.8%	0.4%	0.0%
	MAGNET	0	5	4	11	7	0.0%	3.2%	1.7%	4.7%	4.2%
	(ALL PROVIDERS)	8	1	5	8	4	100%	100%	100%	100%	100%
	OOJ PROVIDERS	0	0	0	0	0	0.0%	0.0%	0.0%	0.0%	0.0%
	CITY CLINIC	5	1	3	3	1	62.5%	100%	60.0%	37.5%	25.0%
	PUBLIC CLINIC (CHN)	0	0	0	1	0	0.0%	0.0%	0.0%	12.5%	0.0%
	JAILS	1	0	0	0	0	12.5%	0.0%	0.0%	0.0%	0.0%
	PRIVATE CLINIC/PMD	0	0	0	2	0	0.0%	0.0%	0.0%	25.0%	0.0%
	PRIVATE HOSPITAL	0	0	0	1	2	0.0%	0.0%	0.0%	12.5%	50.0%
	CHPY	0	0	0	0	1	0.0%	0.0%	0.0%	0.0%	25.0%
ADOLESCENTS	SFGH	2	0	2	1	0	25.0%	0.0%	40.0%	12.5%	0.0%
	OUTREACH	0	0	0	0	0	0.0%	0.0%	0.0%	0.0%	0.0%
	MAGNET	0	0	0	0	0	0.0%	0.0%	0.0%	0.0%	0.0%

### Congenital syphilis

Number of reported congenital syphilis cases (2007) = 0

Number of reported congenital syphilis cases 2003-2007 = 1




Figure 93. Trends in congenital syphilis rate vs. early syphilis among women of childbearing age, San Francisco, 2003-2007.




Figure 94. Congenital syphilis cases by race/ethnicity of mother, San Francisco, 2003-2007.

*Table 19. Congenital syphilis cases and rates by race/ethnicity of mother, San Francisco, 2003-2007. Rates equal cases per 100,000 live births per year. Birth data from Vital Statistics Office, San Francisco Department of Public Health.*

Diagnosis is CONGENITAL SYPHILIS

	Reported cases					Incidence rate				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
(ALL)	0	1	0	0	1	0.0	12.1	0.0	0	0
ASIAN/PI	0	0	0	0	0	0.0	0.0	0.0	0	0
BLACK	0	0	0	0	0	0.0	0.0	0.0	0	0
HISPANIC	0	0	0	0	0	0.0	0.0	0.0	0	0
NATV AMER	0	0	0	0	0	0.0	0.0	0.0	0	0
WHITE	0	1	0	0	0	0.0	33.8	0.0	0	0

*Table 20. Congenital syphilis cases by health care provider, San Francisco, 2003-2007.*

	Reported cases				
	2003	2004	2005	2006	2007
(OTHER PROVIDER)	0	0	0	0	0
OOJ PROVIDER	0	0	0	0	0
OTHER HOSPITAL	0	0	0	0	1
PMD	0	0	0	0	0
SFGH	0	1	0	0	0

### Rectal and pharyngeal infections in men

Number of reported rectal chlamydia cases (2007) = 595

Change in chlamydia cases from 2006 = 15.1% increase

Number of reported rectal gonorrhea cases (2007) = 494

Change in rectal gonorrhea cases from 2006 = 12.3% decline

Rectal and pharyngeal screening for gonorrhea and chlamydia for men who have sex with men began in 2003 and has expanded beyond the STD clinic to local providers serving MSM populations. As a result, increases in case detection may be a function of increased screening for this often asymptomatic infection. Nearly all rectal and pharyngeal specimens collected were tested for both chlamydia and gonorrhea.

- In 2007, cases of rectal chlamydia increased while cases of rectal gonorrhea declined.
- Nearly 60% of rectal chlamydia and gonorrhea cases were among white males. Over half of the reported cases of rectal infections occurred among men 35 years of age or older.
- Since 2003, the proportion of rectal chlamydia cases diagnosed by a private doctor or hospital increased from 6.8% to 20.6%; for gonorrhea the proportions were 17.9% and 18.3%, respectively.




Figure 95. Male rectal gonorrhea cases, San Francisco, 1984-2007. (Data not available before 1984.)




Figure 96. Male rectal chlamydia cases, San Francisco, 2003-2007..




Figure 97. Rectal gonorrhea cases among males by race/ethnicity, San Francisco 2007.




Figure 98. Rectal chlamydia cases among males by race/ethnicity, San Francisco 2007.




Figure 99. Percentage of total rectal gonorrhea cases for selected reporting sources, San Francisco 2003-2007.




Figure 100. Male rectal gonorrhea for San Francisco by age group, 2003-2007.




Figure 101. Male gonorrhea for San Francisco by site of infection, 2003-2007. For patients with multiple sites of infection, cases are included in lower group in bar.




Figure 102. Male chlamydia for San Francisco by site of infection, 2003-2007. For patients with multiple sites of infection, cases are included in lower group in bar.

Table 21. Rectal STDs among male and transgendered residents, San Francisco, 2003-2007.

## Cases of CHLAMYDIA

	Reported cases					Percent of cases				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
<b>Race/ethnicity</b>										
ASIAN OR OTHER	43	57	53	37	63	13.6%	12.6%	10.8%	7.1%	10.5%
AFRICAN AMERICAN	19	13	25	32	33	6.0%	2.8%	5.1%	6.1%	5.5%
HISPANIC	52	73	100	108	131	16.4%	16.2%	20.4%	20.8%	22.0%
NATIVE AMERICAN	3	0	1	1	4	0.9%	0	0.2%	0.1%	0.6%
UNKNOWN	14	53	29	18	28	4.4%	11.8%	5.9%	3.4%	4.7%
WHITE	185	253	280	321	336	58.5%	56.3%	57.3%	62.0%	56.4%
<b>Age group</b>										
(MISSING AGE)	2	0	1	0	2	0.6%	0	0.2%	0	0.3%
15-19 YEARS	8	3	7	7	2	2.5%	0.6%	1.4%	1.3%	0.3%
20-24 YEARS	30	48	57	53	54	9.4%	10.6%	11.6%	10.2%	9.0%
25-29 YEARS	63	83	87	97	90	19.9%	18.4%	17.8%	18.7%	15.1%
30-34 YEARS	61	95	89	80	107	19.3%	21.1%	18.2%	15.4%	17.9%
35-39 YEARS	70	113	114	124	123	22.1%	25.1%	23.3%	23.9%	20.6%
40+ YEARS	82	107	133	156	217	25.9%	23.8%	27.2%	30.1%	36.4%
<b>Reporting source</b>										
OOJ PROVIDERS	1	0	1	2	2	0.3%	0	0.2%	0.3%	0.3%
CITY CLINIC	243	234	254	254	266	76.8%	52.1%	52.0%	49.1%	44.7%
PUBLIC CLINIC (CHN)	5	7	10	10	17	1.5%	1.5%	2.0%	1.9%	2.8%
JAILS	1	0	0	0	0	0.3%	0	0	0	0
PRIVATE CLINIC/PMD	18	36	39	70	112	5.6%	8.0%	7.9%	13.5%	18.8%
PRIVATE HOSPITAL	4	7	4	3	11	1.2%	1.5%	0.8%	0.5%	1.8%
CHPY	2	3	7	6	5	0.6%	0.6%	1.4%	1.1%	0.8%
SFGH	15	14	18	42	28	4.7%	3.1%	3.6%	8.1%	4.7%
OUTREACH	0	2	15	20	26	0	0.4%	3.0%	3.8%	4.3%
MAGNET	27	146	140	110	128	8.5%	32.5%	28.6%	21.2%	21.5%
(TOTAL)	316	449	488	517	595	100%	100%	100%	100%	100%

## Cases of GONORRHEA

	Reported cases					Percent of cases				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
<b>Race/ethnicity</b>										
ASIAN OR OTHER	22	33	32	40	39	7.0%	7.9%	6.7%	7.1%	7.8%
AFRICAN AMERICAN	15	17	22	36	37	4.8%	4.1%	4.6%	6.3%	7.4%
HISPANIC	46	65	98	110	88	14.7%	15.7%	20.8%	19.5%	17.8%
NATIVE AMERICAN	0	3	4	2	5	0	0.7%	0.8%	0.3%	1.0%
UNKNOWN	11	37	29	16	22	3.5%	8.9%	6.1%	2.8%	4.4%
WHITE	217	259	286	359	303	69.7%	62.5%	60.7%	63.7%	61.3%
<b>Age group</b>										
(MISSING AGE)	0	0	0	1	1	0	0	0	0.1%	0.2%
15-19 YEARS	2	3	4	10	4	0.6%	0.7%	0.8%	1.7%	0.8%
20-24 YEARS	31	53	67	71	46	9.9%	12.8%	14.2%	12.6%	9.3%
25-29 YEARS	60	80	83	110	88	19.2%	19.3%	17.6%	19.5%	17.8%
30-34 YEARS	74	105	86	85	99	23.7%	25.3%	18.2%	15.0%	20.0%
35-39 YEARS	78	85	101	115	99	25.0%	20.5%	21.4%	20.4%	20.0%
40+ YEARS	66	88	130	171	157	21.2%	21.2%	27.6%	30.3%	31.7%
<b>Reporting source</b>										
OOJ PROVIDERS	0	2	1	4	0	0	0.4%	0.2%	0.7%	0
CITY CLINIC	222	256	266	314	253	71.3%	61.8%	56.4%	55.7%	51.2%
PUBLIC CLINIC (CHN)	6	3	7	11	15	1.9%	0.7%	1.4%	1.9%	3.0%
JAILS	0	0	1	0	0	0	0	0.2%	0	0
PRIVATE CLINIC/PMD	46	43	45	73	76	14.7%	10.3%	9.5%	12.9%	15.3%
PRIVATE HOSPITAL	10	10	16	12	15	3.2%	2.4%	3.3%	2.1%	3.0%
CHPY	2	7	3	6	3	0.6%	1.6%	0.6%	1.0%	0.6%
SFGH	8	15	14	32	31	2.5%	3.6%	2.9%	5.6%	6.2%

## Cases of GONORRHEA

	Reported cases					Percent of cases				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
OUTREACH	1	0	6	1	8	0.3%	0	1.2%	0.1%	1.6%
MAGNET	16	78	112	110	93	5.1%	18.8%	23.7%	19.5%	18.8%
(TOTAL)	311	414	471	563	494	100%	100%	100%	100%	100%

Table 22. Male chlamydia and gonorrhea among San Francisco residents by site of infection, 2003-2007. (Order of rows reflects precedence for patients with multiple sites of infection.)

	Cases					Percent				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
CHLAMYDIA URETHRAL/URINE	1,277	1,329	1,230	1,185	1,173	77.4%	71.6%	65.3%	61.3%	58.6%
RECTAL	293	415	444	478	546	17.7%	22.3%	23.6%	24.7%	27.2%
PHARYNGEAL	56	78	78	102	82	3.3%	4.2%	4.1%	5.2%	4.0%
UNSPECIFIED/OTHER	22	33	129	167	200	1.3%	1.7%	6.8%	8.6%	9.9%
(TOTAL)	1,648	1,855	1,881	1,932	2,001	100%	100%	100%	100%	100%
GONORRHEA URETHRAL/URINE	847	1,058	1,181	1,081	846	55.0%	55.2%	57.4%	50.9%	49.0%
RECTAL	256	337	358	431	385	16.6%	17.6%	17.4%	20.3%	22.3%
PHARYNGEAL	417	490	397	492	397	27.0%	25.6%	19.3%	23.1%	23.0%
UNSPECIFIED/OTHER	19	29	118	119	96	1.2%	1.5%	5.7%	5.6%	5.5%
(TOTAL)	1,539	1,914	2,054	2,123	1,724	100%	100%	100%	100%	100%

### Other STDs

Number of reported chancroid cases (2007) = 0


Last year a case of chancroid was reported = 2001

Number of reported NGU cases (2007) = 725

Change in reported NGU from 2006 = 15.6% decline

Number of reported PID cases (2007) = 40

Change in reported PID from 2006 = 7.0% decline






Figure 104. Reported pelvic inflammatory disease cases, San Francisco, 2003-2007. "Probable" cases meet CDC case definition for PID.

## II. Prevalence Data

### *Screening program*

STD Prevention and Control Services established a screening program for chlamydia and gonorrhea in 1988. Clinics in the screening program include family planning clinics, teen clinics, correctional facilities and public primary care clinics that serve the indigent and other high-risk populations.

Number of female chlamydia tests provided (2007) = 9,256

Number of male chlamydia tests provided (2007) = 21,142

Number of female gonorrhea tests provided (2007) = 8,297

Number of male gonorrhea tests provided (2007) = 21,586

Number of female syphilis tests provided (2007) = 1,976

Number of male syphilis tests provided (2007) = 7,512

*Table 23. Screening tests performed and STD cases identified by screening site, 2007 only. Syphilis "positives" are confirmed reactive STS at least 1:16.*

Tests for CHLAMYDIA

	(SUTOTAL)	FEMALE			MALE		
		Tests	Cases	Percent	Tests	Cases	Percent
DPH Clinics	CASTRO-MISSION HEALTH CENTER (HC#1)	325	7	2.1%	485	18	3.7%
	CHINATOWN HEALTH CENTER (HC#4)	296	19	6.4%	27	4	14.8%
	HAWKINS VILLAGE	34	2	6.4%	11	0	0.0%
	HIP HOP TO HEALTH CLINIC	145	12	8.3%	24	4	16.6%
	MAXINE HALL HEALTH CENTER (HC#2)	327	13	3.9%	123	7	5.6%
	OCEAN PARK HEALTH CENTER (HC#5)	162	6	3.7%	6	0	0.0%
	POTRERO HILL HEALTH CENTER	114	2	1.7%	62	3	4.8%
	SILVER AVENUE HEALTH CENTER (HC#3)	400	11	2.8%	47	1	2.1%
	SOUTHEAST HEALTH CENTER	538	23	4.3%	151	11	7.3%
	TOM WADDELL HEALTH CENTER	401	7	1.7%	586	19	3.2%
Teen clinics	(SUTOTAL)	1,596	101	6.3%	500	35	7.0%
	BALBOA TEEN HEALTH CENTER	236	8	3.4%	61	2	3.2%
	COLE STREET YOUTH CLINIC	1,015	62	6.1%	148	13	8.7%
Other clinics	LARKIN STREET YOUTH CLINIC	345	31	9.1%	291	20	6.9%
	(SUTOTAL)	1,345	42	3.1%	314	12	3.8%
Detention facilities	CITY COLLEGE HEALTH SERVICE CENTER	560	17	3.0%	156	5	3.2%
	ST. JAMES INFIRMARY	213	4	1.8%	158	7	4.4%
	THE WOMEN'S COMMUNITY CLINIC	572	21	3.7%	(N/A)	(N/A)	(N/A)
	(SUTOTAL)	1,921	142	7.4%	5,618	240	4.2%
	CBS SCREENING CJ9 [CBS/CJ9]	504	45	9.0%	418	24	5.7%
	JHS SCREENING AT CJ1	24	0	0.0%	582	24	4.1%
	JHS SCREENING AT CJ2	1	0	0.0%	514	16	3.1%
	JHS SCREENING AT CJ6	3	0	0.0%	543	22	4.0%
	JHS SCREENING AT CJ8	1,010	54	5.3%	2,316	120	5.2%
	YOUTH GUIDANCE CENTER	379	43	11.4%	1,245	34	2.7%

San Francisco Department of Public Health

Tests for CHLAMYDIA

Community sites	(SUTOTAL)	FEMALE			MALE		
		Tests	Cases	Percent	Tests	Cases	Percent
		205	10	4.9%	1,452	55	3.8%
	440 CLUB	2	0	0.0%	156	5	3.2%
	AGUILAS	(N/A)	(N/A)	(N/A)	4	0	0.0%
	AIDS HEALTH PROJECT	4	0	0.0%	361	18	5.0%
	AIDS OFFICE-CBS	(N/A)	(N/A)	(N/A)	6	0	0.0%
	ASIAN PACIFIC WELLNESS CENTER [API]	(N/A)	(N/A)	(N/A)	29	0	0.0%
	CBS SCREENING [CBS]	(N/A)	(N/A)	(N/A)	1	0	0.0%
	CBS: GOLD'S GYM	15	0	0.0%	438	13	2.9%
	CBS: MACK FOLSOM PRISON	(N/A)	(N/A)	(N/A)	8	0	0.0%
	CENTRAL YMCA	1	0	0.0%	6	0	0.0%
	CLUB PAPI	(N/A)	(N/A)	(N/A)	1	0	0.0%
	ENERGY 92.7 EVENT	4	0	0.0%	107	5	4.6%
	FOLSOM STREET FAIR	19	0	0.0%	158	6	3.8%
	HERMANOS DE LA LUNA SOL	1	0	0.0%	29	1	3.4%
	LATINO GAY PRIDE/BROWN PRIDE	3	0	0.0%	24	0	0.0%
	LPCH ADOLESCENT HEALTH VAN	140	10	7.1%	66	5	7.5%
	NEW LEAF	6	0	0.0%	22	0	0.0%
	SEX CLUB SCREENING [CBS/SC]	7	0	0.0%	3	0	0.0%
	WORLD AIDS DAY FAIR	3	0	0.0%	33	2	6.0%
Studies and anonymous testing	(SUTOTAL)	832	18	2.1%	1,388	48	3.4%
	AIDS OFFICE RESEARCH	2	0	0.0%	193	3	1.5%
	OPTIONS PROJECT	4	0	0.0%	643	34	5.3%
	RED PLUS RESEARCH STUDY	4	0	0.0%	536	11	2.0%
	SWEAT STUDY	713	17	2.3%	9	0	0.0%
	UCSF-UFO STUDY	6	0	0.0%	7	0	0.0%
	WOMEN AND METH (WAM)	103	1	0.9%	(N/A)	(N/A)	(N/A)
Other	(SUTOTAL)	615	20	3.2%	10,348	424	4.1%
	AHP-STOP AIDS	(N/A)	(N/A)	(N/A)	1	0	0.0%
	HEALING ARTS CNTR (THIRD ST CLINIC)	150	12	8.1%	78	7	8.9%
	HOUSING & URBAN HEALTH CLINIC	33	0	0.0%	53	0	0.0%
	LARKIN ST CLINIC - OUTREACH	23	0	0.0%	36	0	0.0%
	MAGNET	27	1	3.7%	6,546	294	4.5%
	MISSION NEIGHBORHOOD RESOURCE CTR	(N/A)	(N/A)	(N/A)	6	0	0.0%
	PHILLIP & SALA BURTON HIGH SCHOOL	168	3	1.7%	231	1	0.4%
	PMD: HASSSLER, SHAWN K.	4	0	0.0%	710	27	3.8%
	PMD: OWEN, WILLIAM F. JR., MD	(N/A)	(N/A)	(N/A)	1,158	45	3.8%
	SFGH -- AIDS (86)	145	2	1.4%	1,252	36	2.8%
	TENDERLOIN HEALTH	2	0	0.0%	10	0	0.0%
	THURGOOD MARSHALL HIGH SCHOOL	59	2	3.3%	79	1	1.2%
	TOM WADDELL HEALTH CENTER - TARC	1	0	0.0%	12	0	0.0%
	UCSF POSITIVE HEALTH PRACTICE	3	0	0.0%	176	13	7.4%
	(TOTAL)	9,256	435	4.7%	21,142	881	4.1%

## Tests for GONORRHEA

			FEMALE			MALE		
			Tests	Cases	Percent	Tests	Cases	Percent
DPH Clinics	(SUTOTAL)		2,580	17	0.6%	1,498	64	4.3%
	CASTRO-MISSION HEALTH CENTER							
	(HC#1)		235	3	1.2%	464	9	1.9%
	CHINATOWN HEALTH CENTER (HC#4)		268	2	0.7%	22	0	0.0%
	HAWKINS VILLAGE		32	0	0.0%	11	2	18.1%
	HIP HOP TO HEALTH CLINIC		139	1	0.7%	23	1	4.3%
	MAXINE HALL HEALTH CENTER							
	(HC#2)		318	3	0.9%	123	4	3.2%
	OCEAN PARK HEALTH CENTER (HC#5)		145	0	0.0%	6	0	0.0%
	POTRERO HILL HEALTH CENTER		115	0	0.0%	60	0	0.0%
	SILVER AVENUE HEALTH CENTER							
	(HC#3)		389	0	0.0%	46	1	2.1%
	SOUTHEAST HEALTH CENTER		537	2	0.3%	152	12	8.0%
	TOM WADDELL HEALTH CENTER		402	6	1.5%	591	35	5.9%
Teen clinics	(SUTOTAL)		1,533	10	0.6%	478	18	3.8%
	BALBOA TEEN HEALTH CENTER		173	0	0.0%	40	0	0.0%
	COLE STREET YOUTH CLINIC		1,014	4	0.3%	148	8	5.4%
	LARKIN STREET YOUTH CLINIC		346	6	1.7%	290	10	3.5%
Other clinics	(SUTOTAL)		694	3	0.4%	313	13	4.2%
	CITY COLLEGE HEALTH SERVICE							
	CENTER		157	0	0.0%	101	4	4.0%
	ST. JAMES INFIRMARY		301	0	0.0%	212	9	4.3%
	THE WOMEN'S COMMUNITY CLINIC		236	3	1.2%	(N/A)	(N/A)	(N/A)
Detention facilities	(SUTOTAL)		1,921	65	3.4%	5,616	52	0.9%
	CBS SCREENING CJ9 [CBS/CJ9]		504	22	4.4%	418	5	1.2%
	JHS SCREENING AT CJ1		24	0	0.0%	581	9	1.5%
	JHS SCREENING AT CJ2		1	0	0.0%	513	11	2.1%
	JHS SCREENING AT CJ6		3	0	0.0%	543	2	0.3%
	JHS SCREENING AT CJ8		1,010	25	2.4%	2,316	19	0.8%
	YOUTH GUIDANCE CENTER		379	18	4.8%	1,245	6	0.4%
Community sites	(SUTOTAL)		124	2	1.6%	1,946	58	2.9%
	440 CLUB		2	0	0.0%	156	6	3.8%
	AGUILAS		(N/A)	(N/A)	(N/A)	4	0	0.0%
	AIDS HEALTH PROJECT		10	1	10.0%	866	31	3.6%
	AIDS OFFICE-CBS		(N/A)	(N/A)	(N/A)	6	0	0.0%
	ASIAN PACIFIC WELLNESS CENTER							
	[API]		(N/A)	(N/A)	(N/A)	56	0	0.0%
	CBS SCREENING [CBS]		(N/A)	(N/A)	(N/A)	1	0	0.0%
	CBS: GOLD'S GYM		15	0	0.0%	438	11	2.5%
	CBS: MACK FOLSOM PRISON		(N/A)	(N/A)	(N/A)	8	0	0.0%
	CENTRAL YMCA		1	0	0.0%	6	0	0.0%
	CLUB PAPI		(N/A)	(N/A)	(N/A)	1	0	0.0%
	ENERGY 92.7 EVENT		4	0	0.0%	107	3	2.8%
	FOLSOM STREET FAIR		19	0	0.0%	158	2	1.2%
	HERMANOS DE LA LUNA SOL		1	0	0.0%	29	1	3.4%
	LATINO GAY PRIDE/BROWN PRIDE		3	0	0.0%	24	1	4.1%
	LPCH ADOLESCENT HEALTH VAN		53	1	1.9%	28	0	0.0%
	NEW LEAF		6	0	0.0%	22	0	0.0%
	SEX CLUB SCREENING [CBS/SC]		7	0	0.0%	3	0	0.0%
	WORLD AIDS DAY FAIR		3	0	0.0%	33	3	9.0%
Studies and anonymous testing	(SUTOTAL)		831	9	1.0%	1,389	40	2.9%
	AIDS OFFICE RESEARCH		2	0	0.0%	195	13	6.6%
	OPTIONS PROJECT		4	0	0.0%	642	20	3.1%
	RED PLUS RESEARCH STUDY		4	0	0.0%	536	7	1.3%
	SWEAT STUDY		713	7	0.9%	9	0	0.0%
	UCSF-UFO STUDY		6	0	0.0%	7	0	0.0%
	WOMEN AND METH (WAM)		102	2	1.9%	(N/A)	(N/A)	(N/A)

San Francisco Department of Public Health

Tests for GONORRHEA

		FEMALE			MALE		
		Tests	Cases	Percent	Tests	Cases	Percent
Other	(SUTOTAL)	614	3	0.4%	10,346	583	5.6%
	AHP-STOP AIDS	(N/A)	(N/A)	(N/A)	1	1	100%
	HEALING ARTS CNTR (THIRD ST CLINIC)	150	2	1.3%	78	3	3.8%
	HOUSING & URBAN HEALTH CLINIC	32	1	3.1%	52	1	1.9%
	LARKIN ST CLINIC - OUTREACH	23	0	0.0%	36	0	0.0%
	MAGNET	27	0	0.0%	6,545	419	6.4%
	MISSION NEIGHBORHOOD RESOURCE CTR	(N/A)	(N/A)	(N/A)	6	0	0.0%
	PHILLIP & SALA BURTON HIGH SCHOOL	168	0	0.0%	231	0	0.0%
	PMD: HASSSLER, SHAWN K.	4	0	0.0%	710	39	5.5%
	PMD: OWEN, WILLIAM F. JR., MD	(N/A)	(N/A)	(N/A)	1,158	47	4.0%
	SFGH -- AIDS (86)	145	0	0.0%	1,252	58	4.6%
	TENDERLOIN HEALTH	2	0	0.0%	10	0	0.0%
	THURGOOD MARSHALL HIGH SCHOOL	59	0	0.0%	79	0	0.0%
	TOM WADDELL HEALTH CENTER - TARC	1	0	0.0%	12	0	0.0%
	UCSF POSITIVE HEALTH PRACTICE	3	0	0.0%	176	15	8.5%
(TOTAL)	(SUTOTAL)	8,297	109	1.3%	21,586	828	3.8%

Tests for SYPHILIS

		FEMALE			MALE		
		Tests	Cases	Percent	Tests	Cases	Percent
DPH Clinics	(SUTOTAL)	665	0	0.0%	2,139	7	0.3%
	CASTRO-MISSION HEALTH CENTER (HC#1)	82	0	0.0%	540	1	0.1%
	CHINATOWN HEALTH CENTER (HC#4)	(N/A)	(N/A)	(N/A)	1	0	0.0%
	HAWKINS VILLAGE	(N/A)	(N/A)	(N/A)	2	0	0.0%
	HIP HOP TO HEALTH CLINIC	7	0	0.0%	2	0	0.0%
	MAXINE HALL HEALTH CENTER (HC#2)	3	0	0.0%	1	0	0.0%
	POTRERO HILL HEALTH CENTER	92	0	0.0%	155	0	0.0%
	TOM WADDELL HEALTH CENTER	481	0	0.0%	1,438	6	0.4%
Teen clinics	(SUTOTAL)	233	0	0.0%	149	3	2.0%
	BALBOA TEEN HEALTH CENTER	11	0	0.0%	5	0	0.0%
	COLE STREET YOUTH CLINIC	191	0	0.0%	76	0	0.0%
	LARKIN STREET YOUTH CLINIC	31	0	0.0%	68	3	4.4%
Other clinics	(SUTOTAL)	234	0	0.0%	113	0	0.0%
	CITY COLLEGE HEALTH SERVICE CENTER	11	0	0.0%	20	0	0.0%
	ST. JAMES INFIRMARY	129	0	0.0%	93	0	0.0%
	THE WOMEN'S COMMUNITY CLINIC	94	0	0.0%	(N/A)	(N/A)	(N/A)
Detention facilities	(SUTOTAL)	479	1	0.2%	736	1	0.1%
	CBS SCREENING CJ9 [CBS/CJ9]	2	0	0.0%	9	0	0.0%
	JHS SCREENING AT CJ1	17	0	0.0%	190	1	0.5%
	JHS SCREENING AT CJ2	2	0	0.0%	140	0	0.0%
	JHS SCREENING AT CJ6	(N/A)	(N/A)	(N/A)	191	0	0.0%
	JHS SCREENING AT CJ7	(N/A)	(N/A)	(N/A)	1	0	0.0%
	JHS SCREENING AT CJ8	432	1	0.2%	56	0	0.0%
	YOUTH GUIDANCE CENTER	26	0	0.0%	149	0	0.0%

## Tests for SYPHILIS

		FEMALE			MALE		
		Tests	Cases	Percent	Tests	Cases	Percent
Community sites	(SUTOTAL)	35	0	0.0%	549	0	0.0%
	440 CLUB	2	0	0.0%	78	0	0.0%
	AGUILAS	(N/A)	(N/A)	(N/A)	2	0	0.0%
	ASIAN PACIFIC WELLNESS CENTER						
	[API]	(N/A)	(N/A)	(N/A)	25	0	0.0%
	CBS SCREENING [CBS]	(N/A)	(N/A)	(N/A)	2	0	0.0%
	CBS: GOLD'S GYM	7	0	0.0%	229	0	0.0%
	CBS: MACK FOLSOM PRISON	(N/A)	(N/A)	(N/A)	4	0	0.0%
	CENTRAL YMCA	1	0	0.0%	4	0	0.0%
	CLUB PAPI	(N/A)	(N/A)	(N/A)	1	0	0.0%
	ENERGY 92.7 EVENT	2	0	0.0%	54	0	0.0%
	FOLSOM STREET FAIR	9	0	0.0%	80	0	0.0%
	HERMANOS DE LA LUNA SOL	1	0	0.0%	18	0	0.0%
	LATINO GAY PRIDE/BROWN PRIDE	1	0	0.0%	14	0	0.0%
	LPCH ADOLESCENT HEALTH VAN	7	0	0.0%	6	0	0.0%
	NEW LEAF	4	0	0.0%	13	0	0.0%
	WORLD AIDS DAY FAIR	1	0	0.0%	19	0	0.0%
Studies and anonymous testing	(SUTOTAL)	295	1	0.3%	555	2	0.3%
	AIDS OFFICE RESEARCH	(N/A)	(N/A)	(N/A)	65	1	1.5%
	OPTIONS PROJECT	17	0	0.0%	485	1	0.2%
	SWEAT STUDY	226	1	0.4%	3	0	0.0%
	UCSF-UFO STUDY	2	0	0.0%	2	0	0.0%
	WOMEN AND METH (WAM)	50	0	0.0%	(N/A)	(N/A)	(N/A)
Other	(SUTOTAL)	35	0	0.0%	3,271	26	0.7%
	AIDS HEALTHCARE FOUNDATION	(N/A)	(N/A)	(N/A)	1	0	0.0%
	LARKIN ST CLINIC - OUTREACH	2	0	0.0%	(N/A)	(N/A)	(N/A)
	MAGNET	13	0	0.0%	3,198	26	0.8%
	TENDERLOIN HEALTH	3	0	0.0%	12	0	0.0%
	TOM WADDELL CLINIC - API	1	0	0.0%	7	0	0.0%
	TOM WADDELL HEALTH CENTER - TARC	16	0	0.0%	53	0	0.0%
(TOTAL)	(SUTOTAL)	1,976	2	0.1%	7,512	39	0.5%

### Sentinel surveillance

Five female clinical sites: City Clinic, Cole Street Youth Clinic, Youth Guidance Center, Ocean Park Health Center, and Southeast Health Center.

Data include testing among women 25 years of age and younger

- Overall, chlamydia positivity among sentinel-surveillance sites declined 13.4% in 2007; gonorrhea positivity declined 27.8%.
- The most dramatic declines for both chlamydia and gonorrhea were among females aged 10-14.




Figure 105. Overall gonorrhea positivity rates by sentinel surveillance sites, 2003-2007.




Figure 106. Chlamydia positivity rates by age group for sentinel surveillance sites, 2003-2007.




Figure 107. Gonorrhea positivity rates by age group for sentinel surveillance sites, 2003-2007.




Figure 108. Chlamydia positivity rates by race/ethnicity for sentinel-surveillance sites, 2003-2007.




Figure 109. Gonorrhea positivity rates by race/ethnicity for sentinel-surveillance sites, 2003-2007.




Figure 110. Positivity rates by race/ethnicity for sentinel-surveillance sites, 2007 only.

Table 24. STD cases identified among women 25 years old or younger and positivity rates for sentinel-surveillance sites, 2003-2007.

		CHLAMYDIA					GONORRHEA				
		2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
CITY CLINIC	positive tests	195	185	245	296	251	81	79	96	125	90
	total tests	2,940	3,507	3,820	3,615	3,553	2,910	3,375	3,616	3,460	3,469
	prevalence	6.6%	5.3%	6.4%	8.2%	7.1%	2.8%	2.3%	2.7%	3.6%	2.6%
COLE STREET	positive tests	96	80	132	137	130	53	58	65	82	67
	total tests	1,327	1,643	1,901	1,723	1,955	1,336	1,539	1,714	1,584	1,880
	prevalence	7.2%	4.9%	6.9%	8.0%	6.6%	4.0%	3.8%	3.8%	5.2%	3.6%
OCEAN PARK	positive tests	34	42	54	52	61	6	7	13	14	4
	total tests	865	1,126	1,270	1,147	1,006	865	1,122	1,266	1,146	1,005
	prevalence	3.9%	3.7%	4.3%	4.5%	6.1%	0.7%	0.6%	1.0%	1.2%	0.4%
SOUTHEAST	positive tests	2	1	0	1	0	0	0	0	0	0
	total tests	56	42	29	25	25	34	32	18	16	18
	prevalence	3.6%	2.4%	0.0%	4.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
YOUTH GUIDANCE	positive tests	18	21	21	40	18	6	5	3	7	2
	total tests	203	271	241	279	195	203	270	239	279	194
	prevalence	8.9%	7.7%	8.7%	14.3%	9.2%	3.0%	1.9%	1.3%	2.5%	1.0%

		CHLAMYDIA					GONORRHEA				
		2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
(ALL AGES)	positive tests	195	185	245	296	251	81	79	96	125	90
	total tests	2,940	3,507	3,820	3,615	3,553	2,910	3,375	3,616	3,460	3,469
	prevalence	6.6%	5.3%	6.4%	8.2%	7.1%	2.8%	2.3%	2.7%	3.6%	2.6%

	CHLAMYDIA					GONORRHEA				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
10-14 YEARS	positive tests	10	6	10	21	9	5	3	4	10
	total tests	169	131	126	179	117	158	129	126	178
	prevalence	5.9%	4.6%	7.9%	11.7%	7.7%	3.2%	2.3%	3.2%	5.6%
15-19 YEARS	positive tests	106	100	106	138	116	29	33	43	45
	total tests	1,315	1,416	1,453	1,331	1,252	1,293	1,391	1,426	1,302
	prevalence	8.1%	7.1%	7.3%	10.4%	9.3%	2.2%	2.4%	3.0%	3.5%
20-25 YEARS	positive tests	79	79	129	137	125	47	43	49	69
	total tests	1,447	1,944	2,237	2,100	2,180	1,450	1,841	2,061	1,976
	prevalence	5.5%	4.1%	5.8%	6.5%	5.7%	3.2%	2.3%	2.4%	3.5%
<hr/>										

		CHLAMYDIA					GONORRHEA				
		2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
(ALL RACES)	positive tests	195	185	245	296	251	81	79	96	125	90
	total tests	2,940	3,507	3,820	3,615	3,553	2,910	3,375	3,616	3,460	3,469
	prevalence	6.6%	5.3%	6.4%	8.2%	7.1%	2.8%	2.3%	2.7%	3.6%	2.6%
ASIAN/PI	positive tests	42	36	41	44	48	17	8	12	10	17
	total tests	639	734	838	766	743	628	682	768	713	718
	prevalence	6.6%	4.9%	4.9%	5.7%	6.5%	2.7%	1.2%	1.6%	1.4%	2.4%
BLACK	positive tests	94	83	94	145	123	37	36	46	74	44
	total tests	1,042	1,156	1,054	1,067	1,003	1,044	1,143	1,041	1,064	1,012
	prevalence	9.0%	7.2%	8.9%	13.6%	12.3%	3.5%	3.1%	4.4%	7.0%	4.3%
HISPANIC	positive tests	30	25	42	47	33	10	9	6	21	9
	total tests	458	558	619	573	544	455	546	590	556	527
	prevalence	6.6%	4.5%	6.8%	8.2%	6.1%	2.2%	1.6%	1.0%	3.8%	1.7%
NATIVE AMERICAN	positive tests	2	4	2	0	1	0	3	2	0	2
	total tests	27	26	22	15	17	27	26	23	13	19
	prevalence	7.4%	15.4%	9.1%	0.0%	5.9%	0.0%	11.5%	8.7%	0.0%	10.5%
UNK/OTHER	positive tests	0	4	2	6	2	0	1	2	2	1
	total tests	5	34	56	62	65	5	35	55	60	65
	prevalence	0.0%	11.8%	3.6%	9.7%	3.1%	0.0%	2.9%	3.6%	3.3%	1.5%
WHITE	positive tests	27	31	49	45	40	17	21	24	16	17
	total tests	755	935	1,080	947	964	737	879	990	872	912
	prevalence	3.6%	3.3%	4.5%	4.8%	4.1%	2.3%	2.4%	2.4%	1.8%	1.9%
<hr/>											

## Detention facilities

Urine-based screening for chlamydia was implemented in the San Francisco County jails in September of 1996, and gonorrhea screening began in March of 1997. We began using the urine-based technology to test for both infections at the youth-detention facility, Youth Guidance Center (YGC), in summer of 1997.

Women in detention facilities were screened for chlamydia and gonorrhea up to age 35. Men were screened for chlamydia and gonorrhea up to age 30. Since gonorrhea was rarely seen in males younger than 18 years of age, we discontinued routine gonorrhea screening of males in youth detention in 2001.

Number of female chlamydia tests provided (2007) = 1,798

Female detention chlamydia positivity (2007) = 8.2%

Number of male chlamydia tests provided (2007) = 4,738

Male detention chlamydia positivity (2007)=4.7%

Number of female gonorrhea tests provided (2007) = 1,554

Female detention gonorrhea positivity (2007) = 3.8%

Number of male gonorrhea tests provided (2007) = 3,498

Male detention gonorrhea positivity (2007) = 0.8%




Figure 111. Chlamydia prevalence by gender and facility, 2003-2007.




Figure 112. Gonorrhea prevalence by gender and facility, 2003-2007. (Males not screened for gonorrhea at YGC since 2000).




Figure 113. Age-specific prevalence by gender for detention screening, 2007.




Figure 114. Prevalence by race/ethnicity for detention screening, 2007.

Table 25. STD cases identified and positivity rates for detention facilities by race/ethnicity of patient, 2003-2007.

		CHLAMYDIA					GONORRHEA				
		2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
(ALL RACES)	positive tests	349	399	385	424	351	73	76	104	110	91
	total tests	6,224	6,421	6,597	6,547	6,292	4,114	4,884	5,023	5,187	5,052
	prevalence	5.6%	6.2%	5.8%	6.4%	5.5%	1.7%	1.5%	2.0%	2.1%	1.8%
ASIAN/PI	positive tests	24	30	22	22	12	2	0	6	4	3
	total tests	528	522	433	407	302	291	348	308	330	255
	prevalence	4.5%	5.7%	5.0%	5.4%	3.9%	0.6%	0.0%	1.9%	1.2%	1.1%
BLACK	positive tests	233	260	224	259	217	53	52	67	75	65
	total tests	3,082	3,142	3,115	2,907	2,741	2,030	2,315	2,375	2,397	2,349
	prevalence	7.5%	8.2%	7.1%	8.9%	7.9%	2.6%	2.2%	2.8%	3.1%	2.7%
HISPANIC	positive tests	45	70	57	82	77	7	6	8	14	6
	total tests	1,455	1,603	1,376	1,446	1,358	911	1,261	1,100	1,216	1,200
	prevalence	3.0%	4.3%	4.1%	5.6%	5.6%	0.7%	0.4%	0.7%	1.1%	0.5%
NATIVE AMERIC-	positive tests	3	4	3	4	2	0	1	0	0	0
CAN	total tests	58	35	22	24	11	49	28	22	24	11
	prevalence	5.1%	11.4%	13.6%	16.6%	18.1%	0.0%	3.5%	0.0%	0.0%	0.0%
UNK/OTHER	positive tests	0	4	1	5	0	0	3	1	1	0
	total tests	22	57	43	52	49	12	40	22	36	37
	prevalence	0.0%	7.0%	2.3%	9.6%	0.0%	0.0%	7.5%	4.5%	2.7%	0.0%
WHITE	positive tests	40	28	41	38	33	11	11	14	14	16
	total tests	994	938	980	917	953	761	810	882	892	938
	prevalence	4.0%	2.9%	4.1%	4.1%	3.4%	1.4%	1.3%	1.5%	1.5%	1.7%

Table 26. STD cases identified and positivity rates for detention facilities by gender of patient, 2003-2007.

Gender is (BOTH GENDERS)

		CHLAMYDIA					GONORRHEA				
		2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
(ALL SITES)	positive tests	349	399	385	424	351	73	76	104	110	91
	total tests	6,224	6,421	6,597	6,547	6,292	4,114	4,884	5,023	5,187	5,052
	prevalence	5.6%	6.2%	5.8%	6.4%	5.5%	1.7%	1.5%	2.0%	2.1%	1.8%
ADULT JAIL	positive tests	266	318	307	303	275	57	67	89	88	74
	total tests	4,429	4,736	4,955	4,750	4,677	3,642	4,472	4,643	4,750	4,679
	prevalence	6.0%	6.7%	6.1%	6.3%	5.8%	1.5%	1.4%	1.9%	1.8%	1.5%
YOUTH GUID- ANCE	positive tests	83	81	78	121	76	16	9	15	22	17
	total tests	1,795	1,685	1,642	1,797	1,615	472	412	380	437	373
	prevalence	4.6%	4.8%	4.7%	6.7%	4.7%	3.3%	2.1%	3.9%	5.0%	4.5%

Gender is FEMALE

		CHLAMYDIA					GONORRHEA				
		2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
(ALL SITES)	positive tests	127	145	158	163	128	46	39	54	55	60
	total tests	1,823	1,532	1,656	1,607	1,554	1,798	1,518	1,652	1,600	1,554
	prevalence	6.9%	9.4%	9.5%	10.1%	8.2%	2.5%	2.5%	3.2%	3.4%	3.8%
ADULT JAIL	positive tests	82	104	119	97	86	30	30	39	33	43
	total tests	1,333	1,107	1,276	1,164	1,181	1,326	1,106	1,272	1,163	1,181
	prevalence	6.1%	9.3%	9.3%	8.3%	7.2%	2.2%	2.7%	3.0%	2.8%	3.6%
YOUTH GUID- ANCE	positive tests	45	41	39	66	42	16	9	15	22	17
	total tests	490	425	380	443	373	472	412	380	437	373
	prevalence	9.1%	9.6%	10.2%	14.8%	11.2%	3.3%	2.1%	3.9%	5.0%	4.5%

Gender is MALE

		CHLAMYDIA					GONORRHEA				
		2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
(ALL SITES)	positive tests	222	254	227	261	223	27	37	50	55	31
	total tests	4,401	4,889	4,941	4,940	4,738	2,316	3,366	3,371	3,587	3,498
	prevalence	5.0%	5.1%	4.5%	5.2%	4.7%	1.1%	1.0%	1.4%	1.5%	0.8%
ADULT JAIL	positive tests	184	214	188	206	189	27	37	50	55	31
	total tests	3,096	3,629	3,679	3,586	3,496	2,316	3,366	3,371	3,587	3,498
	prevalence	5.9%	5.8%	5.1%	5.7%	5.4%	1.1%	1.0%	1.4%	1.5%	0.8%
YOUTH GUID- ANCE	positive tests	38	40	39	55	34	0	0	0	0	0
	total tests	1,305	1,260	1,262	1,354	1,242	0	0	0	0	0
	prevalence	2.9%	3.1%	3.0%	4.0%	2.7%	0	0	0	0	0

Table 27. STD cases identified and positivity rates for detention facilities by age and gender of patient, 2003-2007.

Gender is FEMALE

	CHLAMYDIA					GONORRHEA				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
(ALL AGES)	positive tests	127	145	158	163	128	46	39	54	55
	total tests	1,823	1,532	1,656	1,607	1,554	1,798	1,518	1,652	1,600
	prevalence	6.9%	9.4%	9.5%	10.1%	8.2%	2.5%	2.5%	3.2%	3.4%
10-14 YEARS	positive tests	8	5	7	15	6	4	3	3	9
	total tests	113	79	81	118	74	104	78	81	117
	prevalence	7.0%	6.3%	8.6%	12.7%	8.1%	3.8%	3.8%	3.7%	5.4%
15-19 YEARS	positive tests	59	68	58	79	52	20	10	18	25
	total tests	579	520	493	523	453	571	508	492	518
	prevalence	10.1%	13.0%	11.7%	15.1%	11.4%	3.5%	1.9%	3.6%	4.8%
20-24 YEARS	positive tests	35	47	48	31	40	8	16	23	10
	total tests	572	486	520	435	471	566	485	520	435
	prevalence	6.1%	9.6%	9.2%	7.1%	8.4%	1.4%	3.2%	4.4%	2.3%
25-29 YEARS	positive tests	16	17	28	34	24	12	5	6	9
	total tests	290	254	333	393	384	288	253	331	392
	prevalence	5.5%	6.6%	8.4%	8.6%	6.2%	4.1%	1.9%	1.8%	4.4%
30-34 YEARS	positive tests	9	8	16	4	6	2	5	4	2
	total tests	268	193	219	136	168	268	194	218	136
	prevalence	3.3%	4.1%	7.3%	2.9%	3.5%	0.7%	2.5%	1.8%	2.9%

Gender is MALE

	CHLAMYDIA					GONORRHEA				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
(ALL AGES)	positive tests	222	254	227	261	223	27	37	50	55
	total tests	4,401	4,889	4,941	4,940	4,738	2,316	3,366	3,371	3,587
	prevalence	5.0%	5.1%	4.5%	5.2%	4.7%	1.1%	1.0%	1.4%	0.8%
10-14 YEARS	positive tests	1	3	6	2	2	0	0	0	0
	total tests	303	285	274	303	219	0	0	0	1
	prevalence	0.3%	1.0%	2.1%	0.6%	0.9%	0	0	0	0.0%
15-19 YEARS	positive tests	71	80	60	107	91	5	4	10	6
	total tests	1,429	1,538	1,538	1,597	1,620	347	551	538	554
	prevalence	4.9%	5.2%	3.9%	6.7%	5.6%	1.4%	0.7%	1.8%	0.4%
20-24 YEARS	positive tests	100	110	97	87	82	14	20	25	22
	total tests	1,575	1,723	1,686	1,579	1,482	1,162	1,590	1,532	1,574
	prevalence	6.3%	6.3%	5.7%	5.5%	5.5%	1.2%	1.2%	1.6%	1.0%
25-29 YEARS	positive tests	50	61	64	65	46	8	13	15	27
	total tests	1,090	1,332	1,429	1,454	1,396	805	1,222	1,290	1,453
	prevalence	4.5%	4.5%	4.4%	4.4%	3.2%	0.9%	1.0%	1.1%	0.9%

### III. City Clinic

The San Francisco City Clinic is the only municipal STD clinic in San Francisco, and provides confidential, quality STD services to all residents twelve years of age or older. The clinic is open five days a week for three to eight hours per day (see City Clinic website [www.sfcityclinic.org](http://www.sfcityclinic.org) for exact hours). Appointments are not necessary, though appointments are available to patients returning for follow-up tests or treatments.

The clinic offers evaluation, testing and treatment for gonorrhea, syphilis, chlamydia, and all other STDs. It houses a microbiology lab for STAT testing. In addition, the clinic offers STD patients confidential HIV testing, early care for HIV-infected patients, and family planning services for women, including pregnancy testing and Pap smears.

#### *Patient demographics*

Total patient visits (2007) = 21,842

Total patients (2007) = 13,073

- 3,384 (25.9%) female patient visits, 4,362 (33.4%) men who have sex with men patient visits, 5,255 (40.2%) heterosexual male visits, 65 (0.4%) transgender patient visits
- Average age = 33.5 years old
- 52.6% of patient visits were among whites, 17.2% Hispanics, 15.0% African American, 13.4% Asian/Pacific Islander, and 0.5% Native American
- 79.5% of patient visits among San Francisco residents




Figure 115. Patients and visits at City Clinic, 2003-2007.




Figure 116. City Clinic visits by gender and sexual orientation, 2003-2007.




Figure 117. City Clinic visits by age group for 2007.




Figure 118. Mean age of City Clinic visits and percent under 30 years old, 2003-2007.




Figure 119. City Clinic visits by race/ethnicity for 2007.




Figure 120. City Clinic visits by gender and selected races/ethnicities, 2003-2007.

San Francisco Department of Public Health

Table 28. Demographics of patients and clinic visits. Note: patients and visits missing demographics are not listed, but are included in denominators and totals.

	Orientation	Patients					Percentage of patients				
		2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
(TOTAL)		12,684	13,303	13,690	12,888	13,073	100%	100%	100%	100%	100%
Gender											
FEMALE		2,609	3,179	3,669	3,358	3,384	20.5%	23.8%	26.8%	26.0%	25.8%
TRANSGENDER		41	55	63	70	65	0.3%	0.4%	0.4%	0.5%	0.4%
MALE		10,033	10,069	9,957	9,454	9,617	79.0%	75.6%	72.7%	73.3%	73.5%
Gender	Orientation										
FEMALE	BISEXUAL	235	318	375	329	307	1.8%	2.4%	2.7%	2.5%	2.3%
	LESBIAN	57	85	105	91	84	0.4%	0.6%	0.7%	0.7%	0.6%
	STRAIGHT	2,283	2,746	3,154	2,902	2,944	18.2%	20.8%	23.3%	22.7%	22.8%
	(REFUSED)	5	6	11	10	7					
	(MISSING)	29	24	24	26	42					
TRANSGENDER	BISEXUAL	9	18	17	16	7	0.0%	0.1%	0.1%	0.1%	0.0%
	GAY	17	18	24	21	28	0.1%	0.1%	0.1%	0.1%	0.2%
	STRAIGHT	14	16	20	28	25	0.1%	0.1%	0.1%	0.2%	0.1%
	(REFUSED)	0	1	1	3	0					
	(MISSING)	1	2	1	2	5					
MALE	BISEXUAL	746	704	795	776	811	5.9%	5.3%	5.8%	6.0%	6.2%
	GAY	4,084	3,857	3,626	3,551	3,535	32.6%	29.3%	26.7%	27.8%	27.4%
	STRAIGHT	5,068	5,388	5,417	5,018	5,135	40.5%	40.9%	40.0%	39.4%	39.8%
	(REFUSED)	27	36	51	40	36					
	(MISSING)	108	84	68	69	100					
Ethnicity											
ASIAN/PI		1,564	1,736	1,857	1,729	1,754	12.3%	13.0%	13.5%	13.4%	13.4%
BLACK		1,954	2,014	2,090	2,053	1,971	15.4%	15.1%	15.2%	15.9%	15.0%
HISPANIC		2,283	2,460	2,474	2,392	2,257	17.9%	18.4%	18.0%	18.5%	17.2%
NATV AMER		92	93	82	74	70	0.7%	0.6%	0.5%	0.5%	0.5%
WHITE		6,755	6,957	7,140	6,559	6,884	53.2%	52.2%	52.1%	50.8%	52.6%
Age group											
UNDER 20 YEARS		411	495	565	548	472	3.2%	3.7%	4.1%	4.2%	3.6%
20-24 YEARS		2,008	2,218	2,486	2,282	2,377	15.8%	16.6%	18.1%	17.7%	18.1%
25-29 YEARS		2,713	2,929	2,944	2,828	3,028	21.3%	22.0%	21.5%	21.9%	23.1%
30-34 YEARS		2,487	2,525	2,318	2,204	2,174	19.6%	18.9%	16.9%	17.1%	16.6%
35-39 YEARS		2,001	1,981	2,026	1,819	1,788	15.7%	14.8%	14.7%	14.1%	13.6%
40-44 YEARS		1,354	1,381	1,420	1,360	1,348	10.6%	10.3%	10.3%	10.5%	10.3%
45-54 YEARS		1,296	1,296	1,424	1,325	1,376	10.2%	9.7%	10.4%	10.2%	10.5%
55+ YEARS		411	478	506	521	508	3.2%	3.5%	3.6%	4.0%	3.8%

	Orientation	Visits					Percentage of visits				
		2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
(TOTAL)		21,581	23,040	23,225	21,687	21,842	100%	100%	100%	100%	100%
Gender											
FEMALE		4,122	5,260	6,083	5,447	5,449	19.1%	22.8%	26.1%	25.1%	24.9%
TRANSGENDER		78	112	143	161	124	0.3%	0.4%	0.6%	0.7%	0.5%
MALE		17,380	17,668	16,998	16,071	16,262	80.5%	76.6%	73.1%	74.1%	74.4%
FEMALE	BISEXUAL	360	535	593	532	475	1.6%	2.3%	2.5%	2.4%	2.2%
	LESBIAN	72	104	130	111	115	0.3%	0.4%	0.5%	0.5%	0.5%
	STRAIGHT	3,642	4,573	5,307	4,760	4,789	17.0%	20.0%	23.0%	22.1%	22.2%
	(REFUSED)	12	11	24	13	10					
	(MISSING)	36	37	29	31	60					

		Visits					Percentage of visits				
		2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
TRANSGENDER	BISEXUAL	20	35	37	35	14	0.0%	0.1%	0.1%	0.1%	0.0%
	GAY	34	42	68	48	49	0.1%	0.1%	0.2%	0.2%	0.2%
	STRAIGHT	22	29	35	68	53	0.1%	0.1%	0.1%	0.3%	0.2%
	(REFUSED)	0	3	2	7	0					
	(MISSING)	2	3	1	3	8					
MALE	BISEXUAL	1,489	1,418	1,500	1,489	1,519	6.9%	6.2%	6.5%	6.9%	7.0%
	GAY	8,507	8,309	7,689	7,447	7,252	39.8%	36.4%	33.4%	34.7%	33.6%
	STRAIGHT	7,180	7,758	7,629	6,966	7,282	33.6%	34.0%	33.1%	32.4%	33.7%
	(REFUSED)	45	57	98	67	74					
	(MISSING)	159	126	82	102	135					
Ethnicity											
ASIAN/PI		2,667	2,985	3,125	2,888	2,980	12.3%	12.9%	13.4%	13.3%	13.6%
BLACK		3,250	3,395	3,581	3,435	3,326	15.0%	14.7%	15.4%	15.8%	15.2%
HISPANIC		4,239	4,662	4,683	4,513	4,230	19.6%	20.2%	20.1%	20.8%	19.3%
NATV AMER		174	176	147	126	112	0.8%	0.7%	0.6%	0.5%	0.5%
WHITE		11,207	11,763	11,618	10,612	11,014	51.9%	51.0%	50.0%	48.9%	50.4%
Age group		3	0	1	1	3	0.0%	0	0.0%	0.0%	0.0%
UNDER 20 YEARS		600	753	876	799	684	2.7%	3.2%	3.7%	3.6%	3.1%
20-24 YEARS		3,249	3,773	4,121	3,809	3,749	15.0%	16.3%	17.7%	17.5%	17.1%
25-29 YEARS		4,586	4,963	5,088	4,745	4,993	21.2%	21.5%	21.9%	21.8%	22.8%
30-34 YEARS		4,326	4,375	3,851	3,633	3,714	20.0%	18.9%	16.5%	16.7%	17.0%
35-39 YEARS		3,608	3,561	3,478	3,175	3,101	16.7%	15.4%	14.9%	14.6%	14.1%
40-44 YEARS		2,362	2,572	2,578	2,470	2,426	10.9%	11.1%	11.1%	11.3%	11.1%
45-54 YEARS		2,177	2,211	2,404	2,201	2,302	10.0%	9.5%	10.3%	10.1%	10.5%
55+ YEARS		670	832	828	854	870	3.1%	3.6%	3.5%	3.9%	3.9%

Table 29. Clinic visits by city of residence. Cities with more than 100 visits listed first by frequency of visits; other cities listed alphabetically below. Cities with fewer than two visits in 2007 included in "other." "Homeless" visits only include patients who do not live primarily in one city.

		Visits					Percent				
		2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
SAN FRANCISCO, CA		17,650	18,797	18,615	17,280	17,366	81.7%	81.5%	80.1%	79.6%	79.5%
OAKLAND, CA		794	751	708	662	740	3.6%	3.2%	3.0%	3.0%	3.3%
DALY CITY, CA		370	482	510	462	484	1.7%	2.0%	2.1%	2.1%	2.2%
SOUTH SAN FRANCISCO, CA		138	161	202	204	200	0.6%	0.6%	0.8%	0.9%	0.9%
BERKELEY, CA		157	181	196	187	202	0.7%	0.7%	0.8%	0.8%	0.9%
SAN MATEO, CA		139	121	127	114	120	0.6%	0.5%	0.5%	0.5%	0.5%
PACIFICA, CA		69	78	124	103	98	0.3%	0.3%	0.5%	0.4%	0.4%
SAN JOSE, CA		75	96	107	98	141	0.3%	0.4%	0.4%	0.4%	0.6%
ALAMEDA, CA		81	71	87	75	68	0.3%	0.3%	0.3%	0.3%	0.3%
ALAMO, CA		1	4	2	7	13	0.0%	0.0%	0.0%	0.0%	0.0%
ALBANY, CA		16	12	19	23	18	0.0%	0.0%	0.0%	0.1%	0.0%
ANTIOCH, CA		17	21	34	16	22	0.0%	0.0%	0.1%	0.0%	0.1%
ATHERTON, CA		0	2	2	7	2	0.0%	0.0%	0.0%	0.0%	0.0%
AUSTIN, TX		2	2	2	1	5	0.0%	0.0%	0.0%	0.0%	0.0%
BAKERSFIELD, CA		1	0	2	0	9	0.0%	0.0%	0.0%	0.0%	0.0%
BELMONT, CA		26	23	25	23	25	0.1%	0.0%	0.1%	0.1%	0.1%
BELVEDERE TIBURON, CA		3	11	9	7	2	0.0%	0.0%	0.0%	0.0%	0.0%
BOULDER, CO		2	1	0	0	2	0.0%	0.0%	0.0%	0.0%	0.0%
BRENTWOOD, CA		0	3	3	2	6	0.0%	0.0%	0.0%	0.0%	0.0%
BRISBANE, CA		13	38	39	47	22	0.0%	0.1%	0.1%	0.2%	0.1%
BRODERICK, CA		0	0	1	0	2	0.0%	0.0%	0.0%	0.0%	0.0%
BRONXVILLE, NY		0	0	0	2	2	0.0%	0.0%	0.0%	0.0%	0.0%
BROOKLYN, NY		1	4	5	6	3	0.0%	0.0%	0.0%	0.0%	0.0%

San Francisco Department of Public Health

	Visits					Percent				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
BURLINGAME, CA	42	36	55	55	43	0.1%	0.1%	0.2%	0.2%	0.1%
CALISTOGA, CA	0	1	0	2	2	0.0%	0.0%	0.0%	0.0%	0.0%
CAMBRIDGE, MA	0	0	1	1	4	0.0%	0.0%	0.0%	0.0%	0.0%
CAMPBELL, CA	3	4	6	2	7	0.0%	0.0%	0.0%	0.0%	0.0%
CANOGA PARK, CA	0	0	0	0	2	0.0%	0.0%	0.0%	0.0%	0.0%
CANYON, CA	0	0	0	0	2	0.0%	0.0%	0.0%	0.0%	0.0%
CASTRO VALLEY, CA	9	30	24	13	11	0.0%	0.1%	0.1%	0.0%	0.0%
CLEVELAND, OH	0	1	0	0	2	0.0%	0.0%	0.0%	0.0%	0.0%
CONCORD, CA	68	54	73	64	55	0.3%	0.2%	0.3%	0.2%	0.2%
COOL, CA	0	0	0	0	2	0.0%	0.0%	0.0%	0.0%	0.0%
CORTE MADERA, CA	3	8	15	12	11	0.0%	0.0%	0.0%	0.0%	0.0%
CUPERTINO, CA	11	4	8	3	6	0.0%	0.0%	0.0%	0.0%	0.0%
DANVILLE, CA	9	7	11	11	11	0.0%	0.0%	0.0%	0.0%	0.0%
DAVIS, CA	7	7	5	9	4	0.0%	0.0%	0.0%	0.0%	0.0%
DENVER, CO	0	1	1	6	5	0.0%	0.0%	0.0%	0.0%	0.0%
DORCHESTER, MA	1	0	0	0	2	0.0%	0.0%	0.0%	0.0%	0.0%
DUBLIN, CA	12	8	9	10	11	0.0%	0.0%	0.0%	0.0%	0.0%
DURANGO, CO	0	0	0	0	2	0.0%	0.0%	0.0%	0.0%	0.0%
EL CERRITO, CA	32	21	25	32	31	0.1%	0.0%	0.1%	0.1%	0.1%
EL GRANADA, CA	1	2	1	1	2	0.0%	0.0%	0.0%	0.0%	0.0%
EL SOBRANTE, CA	19	7	10	22	13	0.0%	0.0%	0.0%	0.1%	0.0%
ELK GROVE, CA	3	3	0	3	2	0.0%	0.0%	0.0%	0.0%	0.0%
EMERYVILLE, CA	57	80	66	54	92	0.2%	0.3%	0.2%	0.2%	0.4%
EUGENE, OR	0	1	0	0	3	0.0%	0.0%	0.0%	0.0%	0.0%
EUREKA, CA	0	0	0	0	2	0.0%	0.0%	0.0%	0.0%	0.0%
FAIRFAX, CA	7	8	7	5	6	0.0%	0.0%	0.0%	0.0%	0.0%
FAIRFIELD, CA	4	11	20	20	12	0.0%	0.0%	0.0%	0.0%	0.0%
FOREST KNOLLS, CA	0	0	0	1	6	0.0%	0.0%	0.0%	0.0%	0.0%
FORT LAUDERDALE, FL	1	1	0	0	3	0.0%	0.0%	0.0%	0.0%	0.0%
FOUNTAINVILLE, PA	0	0	2	2	2	0.0%	0.0%	0.0%	0.0%	0.0%
FREMONT, CA	35	62	55	27	49	0.1%	0.2%	0.2%	0.1%	0.2%
FRESNO, CA	2	2	2	6	4	0.0%	0.0%	0.0%	0.0%	0.0%
GOLETA, CA	1	1	0	1	3	0.0%	0.0%	0.0%	0.0%	0.0%
GREENBRAE, CA	6	5	9	1	15	0.0%	0.0%	0.0%	0.0%	0.0%
GROTON, CT	0	0	0	0	2	0.0%	0.0%	0.0%	0.0%	0.0%
GUALALA, CA	0	0	0	1	2	0.0%	0.0%	0.0%	0.0%	0.0%
GUERNEVILLE, CA	1	0	4	4	2	0.0%	0.0%	0.0%	0.0%	0.0%
HALF MOON BAY, CA	1	4	9	5	9	0.0%	0.0%	0.0%	0.0%	0.0%
HAYWARD, CA	74	68	83	101	107	0.3%	0.2%	0.3%	0.4%	0.4%
HERCULES, CA	8	8	9	11	10	0.0%	0.0%	0.0%	0.0%	0.0%
HONOLULU, HI	1	0	2	0	2	0.0%	0.0%	0.0%	0.0%	0.0%
HUNTINGTON BEACH, CA	0	0	4	1	2	0.0%	0.0%	0.0%	0.0%	0.0%
LAFAYETTE, CA	8	9	11	9	11	0.0%	0.0%	0.0%	0.0%	0.0%
LARKSPUR, CA	3	11	3	5	4	0.0%	0.0%	0.0%	0.0%	0.0%
LAS VEGAS, NV	8	6	3	4	3	0.0%	0.0%	0.0%	0.0%	0.0%
LIVERMORE, CA	9	9	13	11	5	0.0%	0.0%	0.0%	0.0%	0.0%
LONG BEACH, CA	1	0	1	0	2	0.0%	0.0%	0.0%	0.0%	0.0%
LONG ISLAND CITY, NY	0	0	1	0	2	0.0%	0.0%	0.0%	0.0%	0.0%
LOS ALTOS, CA	7	9	8	12	10	0.0%	0.0%	0.0%	0.0%	0.0%
LOS ANGELES, CA	12	19	19	15	25	0.0%	0.0%	0.0%	0.0%	0.1%
MANTECA, CA	0	0	3	2	2	0.0%	0.0%	0.0%	0.0%	0.0%
MARIN CITY, CA	1	2	5	10	4	0.0%	0.0%	0.0%	0.0%	0.0%
MARTINEZ, CA	8	4	12	11	8	0.0%	0.0%	0.0%	0.0%	0.0%
MENLO PARK, CA	16	18	6	20	15	0.0%	0.0%	0.0%	0.0%	0.0%
MIAMI, FL	0	3	4	5	3	0.0%	0.0%	0.0%	0.0%	0.0%
MIDDLETOWN, CA	1	1	0	1	2	0.0%	0.0%	0.0%	0.0%	0.0%
MILL VALLEY, CA	38	25	29	35	28	0.1%	0.1%	0.1%	0.1%	0.1%
MILLBRAE, CA	16	36	41	34	30	0.0%	0.1%	0.1%	0.1%	0.1%
MILPITAS, CA	10	8	14	6	3	0.0%	0.0%	0.0%	0.0%	0.0%

	Visits					Percent				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
MINNEAPOLIS, MN	2	2	2	0	2	0.0%	0.0%	0.0%	0.0%	0.0%
MODESTO, CA	5	10	11	10	5	0.0%	0.0%	0.0%	0.0%	0.0%
MONTE RIO, CA	1	1	0	0	2	0.0%	0.0%	0.0%	0.0%	0.0%
MORAGA, CA	8	9	8	5	4	0.0%	0.0%	0.0%	0.0%	0.0%
MOUNTAIN VIEW, CA	47	33	22	22	18	0.2%	0.1%	0.0%	0.1%	0.0%
NAPA, CA	5	12	8	15	6	0.0%	0.0%	0.0%	0.0%	0.0%
NEW HYDE PARK, NY	0	0	0	0	2	0.0%	0.0%	0.0%	0.0%	0.0%
NEW YORK, NY	12	15	11	14	7	0.0%	0.0%	0.0%	0.0%	0.0%
NEWARK, CA	5	7	11	8	11	0.0%	0.0%	0.0%	0.0%	0.0%
NOVATO, CA	14	21	18	32	17	0.0%	0.0%	0.0%	0.1%	0.0%
OMAHA, NE	0	0	0	0	2	0.0%	0.0%	0.0%	0.0%	0.0%
ORINDA, CA	4	0	3	2	6	0.0%	0.0%	0.0%	0.0%	0.0%
PALO ALTO, CA	36	24	33	21	37	0.1%	0.1%	0.1%	0.0%	0.1%
PETALUMA, CA	9	8	7	4	14	0.0%	0.0%	0.0%	0.0%	0.0%
PINOLE, CA	13	9	5	6	9	0.0%	0.0%	0.0%	0.0%	0.0%
PITTSBURG, CA	40	25	25	28	26	0.1%	0.1%	0.1%	0.1%	0.1%
PITTSBURGH, PA	0	0	1	1	2	0.0%	0.0%	0.0%	0.0%	0.0%
PLACERVILLE, CA	0	0	0	2	2	0.0%	0.0%	0.0%	0.0%	0.0%
PLEASANT HILL, CA	9	8	10	2	6	0.0%	0.0%	0.0%	0.0%	0.0%
PLEASANTON, CA	18	5	9	7	16	0.0%	0.0%	0.0%	0.0%	0.0%
PORT COSTA, CA	12	10	8	8	6	0.0%	0.0%	0.0%	0.0%	0.0%
PORTLAND, OR	1	1	2	4	4	0.0%	0.0%	0.0%	0.0%	0.0%
REDDING, CA	3	0	0	0	2	0.0%	0.0%	0.0%	0.0%	0.0%
REDWOOD CITY, CA	53	49	62	49	58	0.2%	0.2%	0.2%	0.2%	0.2%
RENO, NV	0	0	0	1	2	0.0%	0.0%	0.0%	0.0%	0.0%
RICHMOND, CA	78	78	84	81	88	0.3%	0.3%	0.3%	0.3%	0.4%
ROHNERT PARK, CA	6	10	2	1	5	0.0%	0.0%	0.0%	0.0%	0.0%
ROUND ROCK, TX	0	0	0	0	2	0.0%	0.0%	0.0%	0.0%	0.0%
SACRAMENTO, CA	21	18	29	23	15	0.0%	0.0%	0.1%	0.1%	0.0%
SALIDA, CA	1	0	0	0	2	0.0%	0.0%	0.0%	0.0%	0.0%
SALINAS, CA	2	1	3	2	6	0.0%	0.0%	0.0%	0.0%	0.0%
SALT LAKE CITY, UT	2	0	1	0	3	0.0%	0.0%	0.0%	0.0%	0.0%
SAN ANSELMO, CA	4	13	2	5	2	0.0%	0.0%	0.0%	0.0%	0.0%
SAN ANTONIO, TX	1	0	1	0	2	0.0%	0.0%	0.0%	0.0%	0.0%
SAN BRUNO, CA	60	98	94	114	89	0.2%	0.4%	0.4%	0.5%	0.4%
SAN CARLOS, CA	8	18	12	8	17	0.0%	0.0%	0.0%	0.0%	0.0%
SAN DIEGO, CA	9	18	4	7	9	0.0%	0.0%	0.0%	0.0%	0.0%
SAN FERNANDO, CA	0	0	0	0	2	0.0%	0.0%	0.0%	0.0%	0.0%
SAN LEANDRO, CA	67	53	66	71	65	0.3%	0.2%	0.2%	0.3%	0.2%
SAN LORENZO, CA	6	6	5	8	10	0.0%	0.0%	0.0%	0.0%	0.0%
SAN PABLO, CA	54	42	58	26	52	0.2%	0.1%	0.2%	0.1%	0.2%
SAN RAFAEL, CA	24	49	45	49	45	0.1%	0.2%	0.1%	0.2%	0.2%
SAN RAMON, CA	13	11	7	7	8	0.0%	0.0%	0.0%	0.0%	0.0%
SANTA CLARA, CA	11	12	9	12	2	0.0%	0.0%	0.0%	0.0%	0.0%
SANTA CRUZ, CA	7	20	15	13	18	0.0%	0.0%	0.0%	0.0%	0.0%
SANTA ROSA, CA	3	11	18	7	15	0.0%	0.0%	0.0%	0.0%	0.0%
SAUSALITO, CA	18	25	17	26	15	0.0%	0.1%	0.0%	0.1%	0.0%
SEATTLE, WA	4	3	10	4	7	0.0%	0.0%	0.0%	0.0%	0.0%
SEBASTOPOL, CA	8	3	4	6	5	0.0%	0.0%	0.0%	0.0%	0.0%
SONOMA, CA	3	7	6	10	16	0.0%	0.0%	0.0%	0.0%	0.0%
SOQUEL, CA	0	0	0	0	2	0.0%	0.0%	0.0%	0.0%	0.0%
SOUTH LAKE TAHOE, CA	4	1	1	1	4	0.0%	0.0%	0.0%	0.0%	0.0%
SPARKS, NV	0	0	0	1	6	0.0%	0.0%	0.0%	0.0%	0.0%
ST HELENA, CA	1	4	0	0	3	0.0%	0.0%	0.0%	0.0%	0.0%
STOCKTON, CA	7	5	13	23	22	0.0%	0.0%	0.0%	0.1%	0.1%
SUISUN CITY, CA	10	11	2	5	4	0.0%	0.0%	0.0%	0.0%	0.0%
SUNNYVALE, CA	23	19	21	14	19	0.1%	0.0%	0.0%	0.0%	0.0%
SUNOL, CA	0	0	1	2	2	0.0%	0.0%	0.0%	0.0%	0.0%
TACOMA, WA	0	1	0	0	3	0.0%	0.0%	0.0%	0.0%	0.0%


	Visits					Percent				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
TRACY, CA	2	5	10	2	8	0.0%	0.0%	0.0%	0.0%	0.0%
TURLOCK, CA	0	0	0	0	2	0.0%	0.0%	0.0%	0.0%	0.0%
UNION CITY, CA	12	17	12	14	21	0.0%	0.0%	0.0%	0.0%	0.0%
VACAVILLE, CA	6	3	10	16	17	0.0%	0.0%	0.0%	0.0%	0.0%
VALLEJO, CA	53	58	46	42	43	0.2%	0.2%	0.1%	0.1%	0.1%
VANCOUVER, WA	0	1	0	0	2	0.0%	0.0%	0.0%	0.0%	0.0%
VISALIA, CA	0	0	0	0	3	0.0%	0.0%	0.0%	0.0%	0.0%
WALNUT CREEK, CA	24	22	48	42	57	0.1%	0.0%	0.2%	0.1%	0.2%
WASHINGTON, DC	1	0	4	2	3	0.0%	0.0%	0.0%	0.0%	0.0%
(HOMELESS)	281	269	230	212	161	1.3%	1.1%	0.9%	0.9%	0.7%
(MISSING)	125	183	316	359	275	0.5%	0.7%	1.3%	1.6%	1.2%
(OTHER)	214	232	255	239	147	0.9%	1.0%	1.0%	1.1%	0.6%

Table 30. Clinic visits by neighborhood of residence for San Francisco residents.

	Visits					Percent				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
ALEMANY	596	606	651	620	733	2.7%	2.6%	2.8%	2.8%	3.3%
BAYVIEW	185	185	237	236	244	0.8%	0.8%	1.0%	1.0%	1.1%
BERNAL HTS	567	575	609	547	506	2.6%	2.5%	2.6%	2.5%	2.3%
CASTRO	1,220	1,139	1,094	973	1,034	5.7%	5.0%	4.7%	4.5%	4.7%
CATHEDRAL HILL	270	297	332	269	283	1.2%	1.3%	1.4%	1.2%	1.3%
CHINATOWN	359	403	363	362	368	1.6%	1.7%	1.5%	1.6%	1.6%
DIAMOND HTS	181	174	154	159	205	0.8%	0.7%	0.6%	0.7%	0.9%
DOWNTOWN/TENDERLOIN	1,703	1,826	1,839	1,761	1,823	7.9%	8.0%	7.9%	8.2%	8.4%
DUBOCE TRIANGLE	292	243	273	245	235	1.3%	1.0%	1.1%	1.1%	1.0%
GLEN PK	243	277	322	219	292	1.1%	1.2%	1.4%	1.0%	1.3%
GUERRERO	459	442	415	380	391	2.1%	1.9%	1.8%	1.7%	1.8%
HAIGHT DISTRICT	622	605	558	488	493	2.9%	2.6%	2.4%	2.2%	2.2%
INNER SUNSET	244	269	332	271	251	1.1%	1.1%	1.4%	1.2%	1.1%
LAKE MERCED	24	16	16	14	24	0.1%	0.0%	0.0%	0.0%	0.1%
MARINA	365	390	450	390	389	1.7%	1.7%	1.9%	1.8%	1.7%
MISSION	1,143	1,415	1,277	1,140	1,154	5.3%	6.2%	5.5%	5.3%	5.3%
NOB HILL	373	442	394	402	413	1.7%	1.9%	1.7%	1.8%	1.9%
NOE VALLEY	358	326	290	236	231	1.6%	1.4%	1.2%	1.0%	1.0%
NORTH BEACH	248	279	280	256	310	1.1%	1.2%	1.2%	1.1%	1.4%
NORTH MISSION	847	850	772	712	686	3.9%	3.7%	3.3%	3.3%	3.1%
OMI	304	348	372	317	304	1.4%	1.5%	1.6%	1.4%	1.4%
PACIFIC HTS	56	85	80	39	65	0.2%	0.3%	0.3%	0.1%	0.2%
PARK MERCED	121	128	194	132	104	0.5%	0.5%	0.8%	0.6%	0.4%
PORTOLA	120	174	159	138	119	0.5%	0.7%	0.6%	0.6%	0.5%
POTRERO PT	465	350	424	427	413	2.1%	1.5%	1.8%	1.9%	1.9%
PRESIDO	43	51	43	31	51	0.2%	0.2%	0.1%	0.1%	0.2%
RICHMOND	691	733	681	680	706	3.2%	3.2%	2.9%	3.1%	3.2%
SEACLIFF	14	8	11	7	14	0.0%	0.0%	0.0%	0.0%	0.0%
SOUTH-OF-MARKET	914	1,028	960	997	986	4.2%	4.5%	4.1%	4.6%	4.5%
SUNNYDALE	86	83	71	102	90	0.4%	0.3%	0.3%	0.4%	0.4%
SUNSET/PARKSIDE	720	779	756	779	708	3.3%	3.4%	3.2%	3.6%	3.2%
TREASURE ISLAND	276	287	273	301	288	1.2%	1.2%	1.1%	1.4%	1.3%
UNKNOWN	3,961	4,407	4,954	4,601	4,783	18.5%	19.3%	21.5%	21.4%	22.0%
USF/LAUREL HTS	365	481	389	402	419	1.7%	2.1%	1.6%	1.8%	1.9%
VISITACION VLY	211	183	174	177	190	0.9%	0.8%	0.7%	0.8%	0.8%
W HUNTER'S PT	440	536	490	506	513	2.0%	2.3%	2.1%	2.3%	2.3%
W TWIN PEAKS	182	175	166	167	173	0.8%	0.7%	0.7%	0.7%	0.7%
WESTERN ADDITION	1,917	2,083	2,025	1,891	1,591	9.0%	9.1%	8.8%	8.8%	7.3%
WESTWOOD PK	111	89	112	98	94	0.5%	0.3%	0.4%	0.4%	0.4%
(TOTAL)	21,296	22,767	22,992	21,472	21,676	100%	100%	100%	100%	100%

## STDs

- In 2007, City Clinic diagnosed 20.6% of total chlamydia cases, 37.1% of total gonorrhea cases, and 24.1% of total early syphilis.
- City Clinic chlamydia prevalence = 6.2%
  - 4.2% for females; 9.3% MSM; 4.6% other men; 3.4% transgender
- City Clinic gonorrhea prevalence = 5.5%
  - 1.3% for females; 11.9% MSM; 2.2% other men; 4.5% transgender
- Chlamydia prevalence was stable in 2007 for all populations.
- Gonorrhea prevalence declined 21.4% from 2006 through 2007, with most of this attributed to a decline among men.
- Declines were also seen for trichomoniasis (20.0%) and herpes (11.1%) diagnoses; genital warts diagnoses increased in 2007 by 5.3%.






Figure 122. Selected non-reportable STDs diagnosed at City Clinic, 2003-2007.






Figure 123. Gonorrhea prevalence at City Clinic by gender, 2003-2007.



**Table 31. STD cases diagnosed at City Clinic by gender, 2003-2007. Prevalence equals proportion of visits with diagnosis, excluding follow-up visits.**

## Cases among (ALL PATIENTS)

	Reported cases					Prevalence				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
CHLAMYDIA	1,019	970	1,078	1,011	1,023	6.6%	5.9%	6.4%	6.2%	6.2%
GONORRHEA	1,051	1,115	1,116	1,146	906	6.8%	6.8%	6.7%	7.0%	5.5%
SYPHILIS (TOTAL)	243	248	150	167	135	1.6%	1.5%	0.9%	1.0%	0.8%
--PRIMARY	71	83	38	55	29	0.5%	0.5%	0.2%	0.3%	0.2%
--SECONDARY	74	74	44	52	42	0.5%	0.5%	0.3%	0.3%	0.3%
--(TOTAL P&S)	145	157	82	107	71	0.9%	1.0%	0.5%	0.7%	0.4%
--EARLY LATENT	50	49	40	31	32	0.3%	0.3%	0.2%	0.2%	0.2%
--(TOTAL EARLY)	195	206	122	138	103	1.3%	1.3%	0.7%	0.8%	0.6%
--UNKNOWN LATENT [1]	3	2	3	2	1	0.0%	0.0%	0.0%	0.0%	0.0%
--LATE LATENT	45	38	25	27	31	0.3%	0.2%	0.1%	0.2%	0.2%
--NEUROSYPHILIS	0	2	0	0	0	0.0%	0.0%	0.0%	0.0%	0.0%
LYMPHOGRANULOMA VENEREUM	0	14	5	10	4	0.0%	0.1%	0.0%	0.1%	0.0%
TRICHOMONIASIS	116	96	104	89	63	0.7%	0.6%	0.6%	0.5%	0.4%
HERPES	561	471	444	436	389	3.6%	2.9%	2.6%	2.7%	2.4%
GENITAL WARTS	770	679	654	618	655	5.0%	4.1%	3.9%	3.8%	4.0%

## Cases among WOMEN

	Reported cases					Prevalence				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
CHLAMYDIA	125	127	195	182	173	4.0%	3.3%	4.4%	4.4%	4.2%
GONORRHEA	28	46	55	63	53	0.9%	1.2%	1.2%	1.5%	1.3%
SYPHILIS (TOTAL)	5	5	12	4	7	0.2%	0.1%	0.3%	0.1%	0.2%
--PRIMARY	0	0	0	0	0	0.0%	0.0%	0.0%	0.0%	0.0%
--SECONDARY	0	0	2	0	0	0.0%	0.0%	0.0%	0.0%	0.0%
--(TOTAL P&S)	0	0	2	0	0	0.0%	0.0%	0.0%	0.0%	0.0%
--EARLY LATENT	1	0	2	0	0	0.0%	0.0%	0.0%	0.0%	0.0%
--(TOTAL EARLY)	1	0	4	0	0	0.0%	0.0%	0.1%	0.0%	0.0%
--UNKNOWN LATENT [1]	0	0	0	0	0	0.0%	0.0%	0.0%	0.0%	0.0%
--LATE LATENT	4	5	8	4	7	0.1%	0.1%	0.2%	0.1%	0.2%
--NEUROSYPHILIS	0	0	0	0	0	0.0%	0.0%	0.0%	0.0%	0.0%
PID (ALL)	55	44	53	45	49	1.8%	1.1%	1.2%	1.1%	1.2%
--PROBABLE PID [2]	31	26	25	28	35	1.0%	0.7%	0.6%	0.7%	0.8%
--SUSPECT PID	24	18	28	17	14	0.8%	0.5%	0.6%	0.4%	0.3%
MPC	76	65	79	73	62	2.4%	1.7%	1.8%	1.8%	1.5%
LYMPHOGRANULOMA VENEREUM	0	0	0	0	0	0.0%	0.0%	0.0%	0.0%	0.0%
TRICHOMONIASIS	108	91	98	83	60	3.5%	2.3%	2.2%	2.0%	1.4%
HERPES	112	114	134	136	130	3.6%	2.9%	3.0%	3.3%	3.1%
GENITAL WARTS	77	122	130	106	90	2.5%	3.1%	3.0%	2.5%	2.2%

## Cases among MEN

	Reported cases					Prevalence				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
CHLAMYDIA	894	840	877	822	847	7.3%	6.8%	7.1%	6.8%	6.9%
GONORRHEA	1,020	1,065	1,056	1,071	849	8.3%	8.6%	8.6%	8.8%	6.9%
SYPHILIS (TOTAL)	238	241	137	162	124	1.9%	1.9%	1.1%	1.3%	1.0%
--PRIMARY	71	83	38	54	29	0.6%	0.7%	0.3%	0.4%	0.2%
--SECONDARY	74	73	42	52	41	0.6%	0.6%	0.3%	0.4%	0.3%
--(TOTAL P&S)	145	156	80	106	70	1.2%	1.3%	0.7%	0.9%	0.6%
--EARLY LATENT	49	48	38	31	32	0.4%	0.4%	0.3%	0.3%	0.3%
--(TOTAL EARLY)	194	204	118	137	102	1.6%	1.6%	1.0%	1.1%	0.8%
--UNKNOWN LATENT [1]	3	2	3	2	1	0.0%	0.0%	0.0%	0.0%	0.0%
--LATE LATENT	41	33	16	23	21	0.3%	0.3%	0.1%	0.2%	0.2%

## Cases among MEN

	Reported cases					Prevalence				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
---NEUROSYphilis	0	2	0	0	0	0.0%	0.0%	0.0%	0.0%	0.0%
NON-GONOCOCCAL URETHRITIS	1,241	1,224	1,189	1,116	979	10.1%	9.8%	9.7%	9.2%	8.0%
LYMPHOGRANULOMA VENEREUM	0	14	5	10	4	0.0%	0.1%	0.0%	0.1%	0.0%
TRICHOMONIASIS	7	3	6	6	3	0.1%	0.0%	0.0%	0.0%	0.0%
HERPES	448	354	309	300	259	3.6%	2.8%	2.5%	2.5%	2.1%
GENITAL WARTS	687	552	521	507	560	5.6%	4.4%	4.2%	4.2%	4.5%
PROCTITIS	187	152	162	179	132	1.5%	1.2%	1.3%	1.5%	1.1%

<sup>1</sup> cases not known to be less than one year's duration where the patient is 40 years old or less and the initial titer is 1:32 or higher.

<sup>2</sup> PID cases meeting CDC case definition.

Table 32. STD cases diagnosed at City Clinic by gender and sexual orientation for 2007. Percentages equal proportion of visits with diagnosis, excluding follow-up visits.

	(ALL PATIENTS)		WOMEN		GAY/BI MEN		OTHER MEN		TRANSGENDERS	
	cases	percent	cases	percent	cases	percent	cases	percent	cases	percent
CHLAMYDIA	1,023	6.2%	173	4.2%	555	9.3%	292	4.6%	3	3.4%
GONORRHEA	906	5.5%	53	1.3%	711	11.9%	138	2.2%	4	4.5%
SYPHILIS (TOTAL)	135	0.8%	7	0.2%	118	2.0%	6	0.1%	4	4.5%
---PRIMARY	29	0.2%	0	0.0%	29	0.5%	0	0.0%	0	0.0%
---SECONDARY	42	0.3%	0	0.0%	40	0.7%	1	0.0%	1	1.1%
---(TOTAL P&S)	71	0.4%	0	0.0%	69	1.2%	1	0.0%	1	1.1%
---EARLY LATENT	32	0.2%	0	0.0%	31	0.5%	1	0.0%	0	0.0%
---(TOTAL EARLY)	103	0.6%	0	0.0%	100	1.7%	2	0.0%	1	1.1%
---UNKNOWN LATENT [1]	1	0.0%	0	0.0%	1	0.0%	0	0.0%	0	0.0%
---LATE LATENT	31	0.2%	7	0.2%	17	0.3%	4	0.1%	3	3.4%
---NEUROSYphilis	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%
PID (ALL)	(N/A)	(N/A)	49	1.2% (N/A)	(N/A) (N/A)					
---PROBABLE PID [2]	(N/A)	(N/A)	35	0.8% (N/A)	(N/A) (N/A)					
---SUSPECT PID	(N/A)	(N/A)	14	0.3% (N/A)	(N/A) (N/A)					
NON-GONOCOCCAL URETHRITIS	(N/A)	(N/A)	(N/A)	(N/A)	423	7.1%	556	8.8%	(N/A)	(N/A)
MPC	(N/A)	(N/A)	62	1.5% (N/A)	(N/A) (N/A)					
LYMPHOGRANULOMA VENEREUM	4	0.0%	0	0.0%	4	0.1%	0	0.0%	0	0.0%
TRICHOMONIASIS	63	0.4%	60	1.4%	1	0.0%	2	0.0%	0	0.0%
HERPES	389	2.4%	130	3.1%	130	2.2%	129	2.0%	0	0.0%
GENITAL WARTS	655	4.0%	90	2.2%	231	3.9%	329	5.2%	5	5.7%


<sup>1</sup> cases not known to be less than one year's duration where the patient is 40 years old or less and the initial titer is 1:32 or higher.

<sup>2</sup> PID cases meeting CDC case definition.

### HIV testing

- Voluntary, confidential, same-day HIV testing is available to City Clinic patients at no additional cost.
- Overall HIV seropositivity (2007) = 2.2%
- Change in HIV seropositivity from 2006 = 33.3% decline
- HIV seropositivity was highest among males (2.7%) and transgenders (2.1%).
- HIV seropositivity was highest among those who were both MSM and injection-drug users (IDU) (11.4%), MSM (3.1%), and transgenders (2.5%).

Since these HIV data only represent persons who chose to be tested at City Clinic, and because testing criteria have changed over time, the prevalence measured over time may not represent prevalence and incidence in the greater population of San Francisco residents.






Figure 127. HIV positivity rates from voluntary testing at City Clinic by behavior risk profile for 2007. Number of patients tested in BRP group listed at end of bar. (See the legend under Table 34 for abbreviations.)

Table 33. Voluntary HIV test results for STD Control Program, 2003-2007. See list of abbreviations at end of table.

	Seropositive					Post-test counseled				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
<b>sex</b>										
FEMALE	1.2%	0.3%	0.9%	0.3%	0.2%	47.1%	62.8%	61.6%	63.5%	29.1%
MALE	4.2%	4.4%	4.2%	3.6%	2.7%	67.1%	76.2%	74.5%	77.4%	54.8%
TRANS-GENDER	0.0%	0.0%	2.1%	6.4%	2.1%	75.0%	64.5%	59.6%	75.0%	66.7%
<b>Risk group</b>										
MSM NOT IDU (BRP 1)	4.5%	5.0%	4.3%	3.4%	3.1%	68.3%	79.5%	75.4%	78.2%	67.9%
TG NOT IDU (BRP 2)	0.0%	0.0%	2.3%	4.9%	2.5%	81.0%	70.4%	62.8%	76.2%	65.0%
MSM IDU (BRP 3)	2.6%	10.7%	7.5%	8.5%	11.4%	60.3%	81.8%	66.4%	78.0%	65.8%
FEMALE NOT IDU (BRP 4)	1.3%	0.3%	0.9%	0.3%	0.2%	46.6%	62.6%	61.9%	63.5%	29.0%
MSF IDU (BRP 5)	4.8%	0.0%	5.6%	2.5%	0.0%	47.6%	55.3%	83.3%	62.5%	52.6%
TG IDU (BRP 6)	0.0%	0.0%	0.0%	16.7%	0.0%	33.3%	25.0%	25.0%	66.7%	75.0%
FEMALE IDU (BRP 7)	0.0%	0.0%	0.0%	NONE	0.0%	62.5%	100.0%	0.0%	NONE	31.6%
MSF NOT IDU (BRP 8)	2.3%	0.6%	1.7%	2.3%	0.4%	62.1%	61.5%	67.8%	70.5%	19.4%
<b>Age group</b>										
(MISSING AGE)	0.0%	0.0%	0.0%	0.0%	0.0%	53.8%	71.4%	70.0%	50.0%	25.0%
15-19 YEARS	0.0%	0.0%	0.0%	4.3%	1.3%	52.8%	47.7%	50.6%	69.0%	37.7%
20-24 YEARS	2.2%	2.0%	2.0%	3.1%	1.9%	66.5%	67.6%	67.6%	69.9%	40.9%
25-29 YEARS	3.9%	2.7%	4.2%	3.4%	1.9%	63.4%	72.9%	72.9%	73.9%	48.0%
30-34 YEARS	6.6%	4.4%	2.6%	2.6%	2.5%	67.1%	76.2%	72.8%	77.1%	53.0%
35-39 YEARS	3.7%	6.0%	4.7%	4.9%	3.0%	64.7%	78.5%	78.0%	77.7%	55.1%
40-44 YEARS	3.2%	4.0%	5.8%	3.3%	2.3%	63.3%	77.3%	73.4%	79.8%	54.0%
45-54 YEARS	3.2%	3.6%	4.5%	1.8%	2.1%	73.3%	75.2%	71.8%	79.3%	53.8%
55-64 YEARS	1.6%	5.5%	4.6%	3.4%	0.7%	54.7%	80.2%	80.5%	80.9%	58.6%
65+ YEARS	0.0%	4.3%	0.0%	0.0%	3.2%	76.9%	82.6%	93.3%	92.9%	64.5%
<b>Ethnicity</b>										
ASIAN/PI	2.3%	1.9%	2.5%	2.4%	1.8%	68.4%	80.1%	78.7%	79.6%	45.7%

	Seropositive					Post-test counseled				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
BLACK	4.6%	4.0%	5.1%	5.4%	2.2%	58.2%	66.7%	59.8%	68.5%	38.8%
HISPANIC/LATINO	5.2%	5.5%	4.2%	4.7%	2.9%	66.7%	70.2%	72.7%	72.6%	54.5%
NATIVE AMERICAN	0.0%	3.1%	9.1%	5.9%	3.8%	44.4%	56.3%	68.2%	64.7%	53.8%
OTHER	NONE	0.0%	0.0%	0.0%	0.0%	NONE	75.0%	66.7%	60.0%	48.5%
(MISSING)	0.0%	0.0%	0.0%	0.0%	0.0%	57.1%	77.8%	66.7%	85.7%	25.0%
WHITE	3.8%	3.4%	3.6%	2.5%	2.0%	65.8%	75.5%	73.6%	77.9%	52.0%
(ALL PATIENTS)	3.9%	3.7%	3.8%	3.3%	2.2%	65.4%	73.9%	72.5%	75.9%	49.8%

	Seropositive					Post-test counseled				
	2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
<b>sex</b>										
FEMALE	1.2%	0.3%	0.9%	0.3%	0.2%	47.1%	62.8%	61.6%	63.5%	29.1%
MALE	4.2%	4.4%	4.2%	3.6%	2.7%	67.1%	76.2%	74.5%	77.4%	54.8%
TRANS-GENDER	0.0%	0.0%	2.1%	6.4%	2.1%	75.0%	64.5%	59.6%	75.0%	66.7%
<b>Risk group</b>										
MSM NOT IDU (BRP 1)	4.5%	5.0%	4.3%	3.4%	3.1%	68.3%	79.5%	75.4%	78.2%	67.9%
TG NOT IDU (BRP 2)	0.0%	0.0%	2.3%	4.9%	2.5%	81.0%	70.4%	62.8%	76.2%	65.0%
MSM IDU (BRP 3)	2.6%	10.7%	7.5%	8.5%	11.4%	60.3%	81.8%	66.4%	78.0%	65.8%
FEMALE NOT IDU (BRP 4)	1.3%	0.3%	0.9%	0.3%	0.2%	46.6%	62.6%	61.9%	63.5%	29.0%
MSF IDU (BRP 5)	4.8%	0.0%	5.6%	2.5%	0.0%	47.6%	55.3%	83.3%	62.5%	52.6%
TG IDU (BRP 6)	0.0%	0.0%	0.0%	16.7%	0.0%	33.3%	25.0%	25.0%	66.7%	75.0%
FEMALE IDU (BRP 7)	0.0%	0.0%	0.0%	NONE	0.0%	62.5%	100.0%	0.0%	NONE	31.6%
MSF NOT IDU (BRP 8)	2.3%	0.6%	1.7%	2.3%	0.4%	62.1%	61.5%	67.8%	70.5%	19.4%
<b>Age group</b>										
(MISSING AGE)	0.0%	0.0%	0.0%	0.0%	0.0%	53.8%	71.4%	70.0%	50.0%	25.0%
15-19 YEARS	0.0%	0.0%	0.0%	4.3%	1.3%	52.8%	47.7%	50.6%	69.0%	37.7%
20-24 YEARS	2.2%	2.0%	2.0%	3.1%	1.9%	66.5%	67.6%	67.6%	69.9%	40.9%
25-29 YEARS	3.9%	2.7%	4.2%	3.4%	1.9%	63.4%	72.9%	72.9%	73.9%	48.0%
30-34 YEARS	6.6%	4.4%	2.6%	2.6%	2.5%	67.1%	76.2%	72.8%	77.1%	53.0%
35-39 YEARS	3.7%	6.0%	4.7%	4.9%	3.0%	64.7%	78.5%	78.0%	77.7%	55.1%
40-44 YEARS	3.2%	4.0%	5.8%	3.3%	2.3%	63.3%	77.3%	73.4%	79.8%	54.0%
45-54 YEARS	3.2%	3.6%	4.5%	1.8%	2.1%	73.3%	75.2%	71.8%	79.3%	53.8%
55-64 YEARS	1.6%	5.5%	4.6%	3.4%	0.7%	54.7%	80.2%	80.5%	80.9%	58.6%
65+ YEARS	0.0%	4.3%	0.0%	0.0%	3.2%	76.9%	82.6%	93.3%	92.9%	64.5%
<b>Ethnicity</b>										
ASIAN/PI	2.3%	1.9%	2.5%	2.4%	1.8%	68.4%	80.1%	78.7%	79.6%	45.7%
BLACK	4.6%	4.0%	5.1%	5.4%	2.2%	58.2%	66.7%	59.8%	68.5%	38.8%
HISPANIC/LATINO	5.2%	5.5%	4.2%	4.7%	2.9%	66.7%	70.2%	72.7%	72.6%	54.5%
NATIVE AMERICAN	0.0%	3.1%	9.1%	5.9%	3.8%	44.4%	56.3%	68.2%	64.7%	53.8%
OTHER	NONE	0.0%	0.0%	0.0%	0.0%	NONE	75.0%	66.7%	60.0%	48.5%
(MISSING)	0.0%	0.0%	0.0%	0.0%	0.0%	57.1%	77.8%	66.7%	85.7%	25.0%
WHITE	3.8%	3.4%	3.6%	2.5%	2.0%	65.8%	75.5%	73.6%	77.9%	52.0%
(ALL PATIENTS)	3.9%	3.7%	3.8%	3.3%	2.2%	65.4%	73.9%	72.5%	75.9%	49.8%

Abbreviations:

- MSM: men who have sex with men
- MSW: men who have sex with women (i.e., not with men)
- TG: transgender
- IDU: injection drug user
- BRP: behavior risk profile (defined by HPPC)

**Table 34. Voluntary HIV testing for STD Control Program by type of test, 2003-2007. All patients testing negative on standard or rapid HIV test have been pooled for HIV RNA testing since 2003.**

		2003	2004	2005	2006	2007
STANDARD ANTIBODY TEST	Patients tested	2,756	3,733	2,992	2,513	2,083
	Seropositive number	108	130	101	70	84
	percent	3.9%	3.5%	3.4%	2.8%	4.1%
RAPID TEST	Patients tested	1	61	269	591	3,191
	Seropositive number	0	9	21	32	30
	percent	0.0%	15.0%	7.8%	5.4%	0.9%
RNA POOLING	Patients tested	2	8	8	4	1,619
	Seropositive number	0	0	0	0	0
	percent	0.0%	0.0%	0.0%	0.0%	0.0%

## Appendix I. Demographic Breakdowns of STD Morbidity

Table 35. Major STD cases and rates by all demographic combinations, 2003-2007.

Breakdown by (NONE)

Cases of	Gender	Race	Age	Reported cases					Incidence rate				
				2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
Cases of CHLAMYDIA	(BOTH)	(ALL)	15-19 YRS	3350	3663	3710	4053	3939	431.3	471.6	477.6	521.8	507.1
GONORRHEA	(BOTH)	(ALL)	20-24 YRS	1795	2153	2411	2468	2010	231.1	277.2	310.4	317.7	258.8
EARLY SYPHILIS	(BOTH)	(ALL)	25-29 YRS	527	552	428	420	352	67.8	71.1	55.1	54.1	45.3

Breakdown by AGE

Cases of	Gender	Race	Age	Reported cases					Incidence rate				
				2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
Cases of CHLAMYDIA	(BOTH)	(ALL)	15-19 YRS	714	633	671	825	693	2142.0	1899.0	2013.0	2475.0	2079.0
			20-24 YRS	885	997	983	967	995	1578.8	1778.6	1753.7	1725.1	1775.1
			25-29 YRS	602	665	686	776	710	658.7	727.6	750.6	849.1	776.9
			30-34 YRS	421	487	438	435	454	472.9	547.0	492.0	488.6	510.0
			35-39 YRS	310	381	380	408	370	428.6	526.8	525.4	564.1	511.6
			40-44 YRS	196	247	243	302	286	318.8	401.8	395.3	491.2	465.2
			45-54 YRS	146	157	206	229	307	135.5	145.8	191.2	212.6	285.0
			55-64 YRS	25	39	52	44	65	38.3	59.7	79.7	67.4	99.6
			65+ YRS	6	*	6	8	5	5.7	*	5.7	7.5	4.7
Cases of GONORRHEA	(BOTH)	(ALL)	15-19 YRS	106	106	170	167	120	318.0	318.0	510.0	501.0	360.0
			20-24 YRS	228	317	361	367	264	406.8	565.5	644.0	654.7	471.0
			25-29 YRS	317	346	377	456	337	346.9	378.6	412.5	498.9	368.7
			30-34 YRS	341	406	366	343	330	383.0	456.1	411.1	385.3	370.7
			35-39 YRS	353	398	475	415	315	488.1	550.3	656.7	573.8	435.5
			40-44 YRS	232	279	316	363	304	377.4	453.8	514.0	590.5	494.5
			45-54 YRS	173	239	275	285	257	160.6	221.9	255.3	264.6	238.6
			55-64 YRS	25	46	52	45	62	38.3	70.5	79.7	68.9	95.0
			65+ YRS	11	5	10	9	5	10.4	4.7	9.4	8.5	4.7
EARLY SYPHILIS	(BOTH)	(ALL)	15-19 YRS	*	0	*	*	*	0.0	*	*	*	*
			20-24 YRS	25	27	15	20	17	44.6	48.2	26.8	35.7	30.3
			25-29 YRS	49	62	39	40	34	53.6	67.8	42.7	43.8	37.2
			30-34 YRS	123	97	66	54	37	138.2	109.0	74.1	60.7	41.6
			35-39 YRS	136	126	102	99	69	188.0	174.2	141.0	136.9	95.4
			40-44 YRS	96	136	94	88	74	156.2	221.2	152.9	143.1	120.4
			45-54 YRS	76	82	92	87	95	70.6	76.1	85.4	80.8	88.2
			55-64 YRS	15	21	14	24	22	23.0	32.2	21.4	36.8	33.7
			65+ YRS	*	*	*	5	*	*	*	*	4.7	*

Breakdown by RACE

Cases of	Gender	Race	Age	Reported cases					Incidence rate				
				2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
Cases of CHLAMYDIA	(BOTH)	ASIAN/PI	(ALL)	490	470	492	486	540	194.2	186.2	194.9	192.6	214.0
		BLACK	(ALL)	829	885	854	1007	894	1293.9	1381.3	1332.9	1571.7	1395.3
		HISPANIC	(ALL)	534	582	608	646	588	487.7	531.5	555.2	589.9	537.0
		NATV AMER	(ALL)	28	16	26	15	24	614.2	351.0	570.3	329.0	526.4
		WHITE	(ALL)	787	911	970	1044	1038	232.2	268.8	286.2	308.0	306.3
Cases of GONORRHEA	(BOTH)	ASIAN/PI	(ALL)	129	144	161	172	134	51.1	57.1	63.8	68.2	53.1
		BLACK	(ALL)	328	363	515	544	422	511.9	566.6	803.8	849.1	658.7
		HISPANIC	(ALL)	222	267	335	340	281	202.7	243.8	305.9	310.5	256.6
		NATV AMER	(ALL)	9	19	24	10	11	197.4	416.8	526.4	219.3	241.3
		WHITE	(ALL)	901	1047	1075	1170	951	265.9	308.9	317.2	345.2	280.6

## Breakdown by RACE

			Reported cases					Incidence rate				
			2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
EARLY SYPHILIS (BOTH)	ASIAN/PI	(ALL)	40	48	35	27	24	15.8	19.0	13.9	10.7	9.5
	BLACK	(ALL)	34	50	28	51	35	53.1	78.0	43.7	79.6	54.6
	HISPANIC	(ALL)	114	113	89	74	56	104.1	103.2	81.3	67.6	51.1
	NATV AMER	(ALL)	*	6	7	6	*	*	131.6	153.5	131.6	*
	WHITE	(ALL)	330	318	257	245	223	97.4	93.8	75.8	72.3	65.8

## Breakdown by RACE AND AGE

Cases of	Gender	Race	Age	Reported cases					Incidence rate				
				2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
CHLAMYDIA (BOTH)	ASIAN/PI	15-19 YRS	92	59	67	74	69	656.3	420.9	477.9	527.9	492.2	
		20-24 YRS	124	136	151	146	140	671.3	736.2	817.4	790.4	757.9	
		25-29 YRS	103	95	91	80	105	434.8	401.0	384.2	337.7	443.3	
		30-34 YRS	69	75	62	62	75	325.5	353.8	292.5	292.5	353.8	
		35-39 YRS	39	45	57	40	62	191.6	221.1	280.1	196.5	304.6	
		40-44 YRS	31	27	30	41	36	162.0	141.1	156.7	214.2	188.1	
		45-54 YRS	25	24	24	30	34	70.9	68.1	68.1	85.1	96.5	
		55-64 YRS	5	*	8	7	16	21.8	*	34.9	30.5	69.8	
		65+ YRS	*	0	0	*	*	*	0.0	0.0	*	*	
	BLACK	15-19 YRS	313	296	290	395	316	7736.9	7316.7	7168.4	9763.8	7811.1	
		20-24 YRS	259	285	257	254	264	6624.0	7289.0	6572.9	6496.1	6751.9	
		25-29 YRS	112	134	132	165	124	2538.4	3037.0	2991.7	3739.6	2810.4	
		30-34 YRS	50	64	70	64	65	1019.0	1304.3	1426.6	1304.3	1324.7	
		35-39 YRS	35	44	39	51	42	666.8	838.3	743.0	971.7	800.2	
		40-44 YRS	26	32	25	27	27	481.0	592.0	462.5	499.5	499.5	
		45-54 YRS	17	9	21	27	23	185.2	98.1	228.8	294.2	250.6	
		55-64 YRS	*	*	*	*	12	*	*	*	*	204.2	
		65+ YRS	*	0	0	0	0	*	0.0	0.0	0.0	0.0	
HISPANIC	ASIAN/PI	15-19 YRS	112	97	98	143	103	1502.8	1301.5	1314.9	1918.7	1382.0	
		20-24 YRS	172	184	188	169	167	1576.7	1686.7	1723.3	1549.2	1530.8	
		25-29 YRS	107	129	140	155	112	803.1	968.2	1050.7	1163.3	840.6	
		30-34 YRS	52	54	77	69	72	426.4	442.8	631.4	565.8	590.4	
		35-39 YRS	42	64	51	55	56	437.6	666.8	531.4	573.0	583.5	
		40-44 YRS	24	29	28	33	39	299.8	362.3	349.8	412.2	487.2	
		45-54 YRS	11	16	19	16	32	97.8	142.2	168.9	142.2	284.4	
		55-64 YRS	*	*	*	*	*	*	*	*	*	*	
		65+ YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	
NATV AMER	ASIAN/PI	15-19 YRS	*	*	7	*	*	*	*	3703.8	*	*	
		20-24 YRS	10	*	*	5	7	2512.0	*	*	1256.0	1758.4	
		25-29 YRS	7	6	6	*	*	1135.7	973.5	973.5	*	*	
		30-34 YRS	*	*	*	*	*	*	*	*	*	*	
		35-39 YRS	5	*	*	*	5	1002.6	*	*	*	1002.6	
		40-44 YRS	0	*	*	0	*	0.0	*	*	0.0	*	
		45-54 YRS	0	0	*	*	*	0.0	0.0	*	*	*	
		55-64 YRS	0	0	*	0	0	0.0	0.0	*	0.0	0.0	
		65+ YRS	0	0	0	*	0	0.0	0.0	0.0	*	0.0	
WHITE	ASIAN/PI	15-19 YRS	49	62	50	54	53	676.6	856.1	690.4	745.7	731.8	
		20-24 YRS	120	169	185	190	183	554.9	781.5	855.5	878.6	846.2	
		25-29 YRS	149	150	179	204	185	308.4	310.4	370.4	422.2	382.9	
		30-34 YRS	153	175	136	124	153	310.8	355.5	276.3	251.9	310.8	
		35-39 YRS	141	161	163	184	136	392.7	448.4	453.9	512.4	378.7	
		40-44 YRS	87	105	120	149	149	311.7	376.2	430.0	533.9	533.9	
		45-54 YRS	70	66	104	109	150	138.4	130.5	205.6	215.5	296.6	
		55-64 YRS	13	20	27	20	23	44.8	69.0	93.1	69.0	79.3	
		65+ YRS	*	*	*	*	*	*	*	*	*	*	

San Francisco Department of Public Health

Breakdown by RACE AND AGE

GONORRHEA	(BOTH) ASIAN/PI	Reported cases							Incidence rate					
		2003		2004		2005		2006		2003	2004	2005	2006	2007
		15-19 YRS	20-24 YRS	25-29 YRS	30-34 YRS	35-39 YRS	40-44 YRS	45-54 YRS	55-64 YRS	*	*	*	*	*
BLACK		10	7	9	9	6	71.3	49.9	64.2	64.2	42.8			
		20-24 YRS	10	26	33	26	24	54.1	140.7	178.6	140.7	129.9		
		25-29 YRS	32	33	36	34	24	135.1	139.3	152.0	143.5	101.3		
		30-34 YRS	21	28	35	44	35	99.1	132.1	165.1	207.6	165.1		
		35-39 YRS	30	25	23	21	22	147.4	122.8	113.0	103.2	108.1		
		40-44 YRS	16	14	12	21	13	83.6	73.1	62.7	109.7	67.9		
		45-54 YRS	7	10	9	11	6	19.9	28.4	25.5	31.2	17.0		
		55-64 YRS	*	*	*	*	*	*	*	*	*	*	*	*
HISPANIC		65+ YRS	0	0	0	0	*	0.0	0.0	0.0	0.0	0.0	*	
		15-19 YRS	64	47	92	103	65	1582.0	1161.8	2274.1	2546.0	1606.7		
		20-24 YRS	59	88	109	119	85	1508.9	2250.6	2787.7	3043.5	2173.9		
		25-29 YRS	53	57	71	91	80	1201.2	1291.9	1609.2	2062.4	1813.1		
		30-34 YRS	53	38	53	50	52	1080.1	774.4	1080.1	1019.0	1059.7		
		35-39 YRS	28	39	66	54	39	533.5	743.0	1257.4	1028.8	743.0		
		40-44 YRS	28	43	49	61	37	518.0	795.6	906.6	1128.6	684.6		
		45-54 YRS	34	40	56	51	43	370.4	435.8	610.1	555.7	468.5		
NATV AMER		55-64 YRS	*	10	10	6	12	*	170.2	170.2	102.1	204.2		
		65+ YRS	*	*	*	*	0	*	*	*	*	0.0		
		15-19 YRS	9	16	21	25	18	120.8	214.7	281.8	335.4	241.5		
		20-24 YRS	44	37	51	62	50	403.3	339.2	467.5	568.3	458.3		
		25-29 YRS	41	66	83	86	57	307.7	495.3	622.9	645.5	427.8		
		30-34 YRS	51	55	49	51	47	418.2	451.0	401.8	418.2	385.4		
		35-39 YRS	34	51	68	58	48	354.2	531.4	708.5	604.3	500.1		
		40-44 YRS	25	21	33	36	30	312.3	262.3	412.2	449.7	374.8		
WHITE		45-54 YRS	16	17	26	20	25	142.2	151.1	231.1	177.8	222.2		
		55-64 YRS	0	*	*	*	0	0.0	*	*	*	*		
		65+ YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0		
		15-19 YRS	*	*	*	0	0	*	*	*	0.0	0.0		
		20-24 YRS	*	*	*	*	*	*	*	*	*	*		
		25-29 YRS	0	*	*	*	*	0.0	*	*	*	*		
		30-34 YRS	*	9	7	0	*	*	1547.0	1203.2	0.0	*		
		35-39 YRS	*	*	5	*	*	*	*	1002.6	*	*		
		40-44 YRS	*	*	*	*	*	*	*	*	*	*		
		45-54 YRS	0	*	0	0	0	0.0	*	0.0	0.0	0.0		
		55-64 YRS	0	0	*	0	0	0.0	0.0	*	0.0	0.0		
		65+ YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0		
		15-19 YRS	10	15	22	11	13	138.1	207.1	303.8	151.9	179.5		
		20-24 YRS	91	121	120	134	76	420.8	559.5	554.9	619.7	351.4		
		25-29 YRS	161	150	142	195	147	333.2	310.4	293.9	403.6	304.2		
		30-34 YRS	173	211	175	165	161	351.4	428.6	355.5	335.2	327.0		
		35-39 YRS	222	226	251	237	172	618.2	629.4	699.0	660.0	479.0		
		40-44 YRS	132	151	178	216	187	473.0	541.1	637.8	774.0	670.1		
		45-54 YRS	91	137	155	175	158	179.9	270.9	306.5	346.0	312.4		
		55-64 YRS	17	31	27	30	31	58.6	106.9	93.1	103.5	106.9		
		65+ YRS	*	*	5	5	*	*	*	10.6	10.6	*		

## Breakdown by RACE AND AGE

			Reported cases					Incidence rate				
			2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
			15-19 YRS	0	0	*	0	0.0	0.0	*	0.0	0.0
EARLY SYPHILIS (BOTH)	ASIAN/PI	20-24 YRS	*	*	*	0	*	*	*	*	0.0	*
		25-29 YRS	9	10	*	6	*	38.0	42.2	*	25.3	*
		30-34 YRS	16	10	11	*	*	75.5	47.2	51.9	*	*
		35-39 YRS	5	16	6	8	*	24.6	78.6	29.5	39.3	*
		40-44 YRS	7	7	6	*	5	36.6	36.6	31.3	*	26.1
		45-54 YRS	*	*	5	6	6	*	*	14.2	17.0	17.0
		55-64 YRS	*	*	*	0	*	*	*	*	0.0	*
		65+ YRS	0	0	*	0	0	0.0	0.0	*	0.0	0.0
		15-19 YRS	0	0	*	*	*	0.0	0.0	*	*	*
		20-24 YRS	*	*	*	5	*	*	*	*	127.9	*
BLACK		25-29 YRS	*	*	*	*	*	*	*	*	*	*
		30-34 YRS	*	12	*	8	*	*	244.6	*	163.0	*
		35-39 YRS	11	10	5	10	6	209.6	190.5	95.3	190.5	114.3
		40-44 YRS	7	14	9	10	7	129.5	259.0	166.5	185.0	129.5
		45-54 YRS	8	6	7	12	8	87.2	65.4	76.3	130.7	87.2
		55-64 YRS	*	*	0	*	*	*	*	0.0	*	*
		65+ YRS	0	0	0	0	*	0.0	0.0	0.0	0.0	*
		15-19 YRS	*	0	0	0	0	*	0.0	0.0	0.0	0.0
		20-24 YRS	13	9	9	*	*	119.2	82.5	82.5	*	*
		25-29 YRS	16	22	12	14	9	120.1	165.1	90.1	105.1	67.5
HISPANIC		30-34 YRS	25	22	15	9	5	205.0	180.4	123.0	73.8	41.0
		35-39 YRS	27	24	16	16	12	281.3	250.1	166.7	166.7	125.0
		40-44 YRS	18	20	21	14	11	224.9	249.8	262.3	174.9	137.4
		45-54 YRS	9	15	15	15	14	80.0	133.3	133.3	133.3	124.4
		55-64 YRS	*	*	*	*	*	*	*	*	*	*
		65+ YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0
		15-19 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0
		20-24 YRS	0	*	0	0	*	0.0	*	0.0	0.0	*
		25-29 YRS	0	*	*	*	0	0.0	*	*	*	0.0
		30-34 YRS	*	0	*	0	0	*	0.0	*	0.0	0.0
NATV AMER		35-39 YRS	0	*	*	0	0	0.0	*	*	*	0.0
		40-44 YRS	0	*	0	*	0	0.0	*	0.0	*	0.0
		45-54 YRS	*	*	0	*	*	*	*	0.0	*	*
		55-64 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0
		65+ YRS	0	0	0	*	0	0.0	0.0	0.0	*	0.0
		15-19 YRS	0	0	*	0	*	0.0	0.0	*	0.0	*
		20-24 YRS	9	9	*	12	7	41.6	41.6	*	55.5	32.4
		25-29 YRS	23	26	20	17	16	47.6	53.8	41.4	35.2	33.1
		30-34 YRS	75	49	32	32	24	152.3	99.5	65.0	65.0	48.8
		35-39 YRS	91	73	68	59	45	253.4	203.3	189.4	164.3	125.3
WHITE		40-44 YRS	62	88	57	57	50	222.2	315.3	204.2	204.2	179.2
		45-54 YRS	57	55	62	47	61	112.7	108.8	122.6	92.9	120.6
		55-64 YRS	10	17	12	17	19	34.5	58.6	41.4	58.6	65.5
		65+ YRS	*	*	*	*	0	*	*	*	*	0.0

## Breakdown by SEX

Cases of	Gender	Race	Age	Reported cases					Incidence rate				
				2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
CHLAMYDIA	FEMALE	(ALL)	(ALL)	1680	1781	1768	2058	1875	439.9	466.3	462.9	538.9	491.0
				MALE	(ALL)	(ALL)	1648	1855	1881	1932	2001	417.4	469.8
GONORRHEA	FEMALE	(ALL)	(ALL)	250	233	349	333	271	65.5	61.0	91.4	87.2	71.0
				MALE	(ALL)	(ALL)	1539	1914	2054	2123	1724	389.8	484.8

*San Francisco Department of Public Health*

Breakdown by SEX

EARLY SYPHILIS		Reported cases						Incidence rate				
		2003	2004	2005	2006	2007	2003	2004	2005	2006	2007	
		FEMALE (ALL) (ALL)	10	*	8	*	5	2.6	*	2.1	*	1.3
	MALE (ALL) (ALL)	517	549	420	416	347	130.9	139.0	106.4	105.4	87.9	

Breakdown by SEX AND AGE

Cases of	Gender	Race	Age	Reported cases					Incidence rate				
				2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
CHLAMYDIA	FEMALE	(ALL)	15-19 YRS	539	461	494	609	505	3304.7	2826.5	3028.8	3733.9	3096.3
			20-24 YRS	564	630	587	619	645	1989.6	2222.5	2070.8	2183.7	2275.4
			25-29 YRS	276	328	325	401	348	621.3	738.4	731.6	902.7	783.4
			30-34 YRS	136	155	155	169	153	335.7	382.6	382.6	417.1	377.6
			35-39 YRS	55	72	83	96	78	172.1	225.3	259.8	300.5	244.1
			40-44 YRS	39	47	44	62	39	139.5	168.2	157.4	221.8	139.5
			45-54 YRS	30	40	33	44	58	58.2	77.5	64.0	85.3	112.4
			55-64 YRS	7	9	14	10	13	21.1	27.1	42.2	30.1	39.2
			65+ YRS	0	0	*	*	*	0.0	0.0	*	*	*
	MALE	(ALL)	15-19 YRS	167	163	149	185	148	981.0	957.5	875.2	1086.7	869.4
			20-24 YRS	317	360	377	337	339	1144.1	1299.3	1360.7	1216.3	1223.5
			25-29 YRS	322	333	357	363	356	685.5	709.0	760.1	772.8	757.9
			30-34 YRS	281	331	279	262	301	579.3	682.3	575.1	540.1	620.5
			35-39 YRS	255	308	295	311	290	631.6	762.8	730.6	770.3	718.2
			40-44 YRS	157	200	199	240	247	468.3	596.6	593.6	715.9	736.8
			45-54 YRS	116	117	172	184	249	206.6	208.4	306.4	327.8	443.6
			55-64 YRS	18	29	38	34	52	56.1	90.4	118.4	106.0	162.0
			65+ YRS	6	*	*	7	*	13.5	*	*	15.8	*
GONORRHEA	FEMALE	(ALL)	15-19 YRS	71	67	108	110	69	435.3	410.8	662.2	674.4	423.1
			20-24 YRS	63	75	98	96	69	222.2	264.6	345.7	338.7	243.4
			25-29 YRS	52	32	53	58	50	117.1	72.0	119.3	130.6	112.6
			30-34 YRS	24	18	26	18	33	59.2	44.4	64.2	44.4	81.5
			35-39 YRS	11	10	21	15	19	34.4	31.3	65.7	46.9	59.5
			40-44 YRS	9	13	18	6	9	32.2	46.5	64.4	21.5	32.2
			45-54 YRS	9	11	14	12	9	17.4	21.3	27.1	23.3	17.4
			55-64 YRS	*	*	5	*	*	*	*	15.1	*	*
			65+ YRS	*	0	0	*	0	*	0.0	0.0	*	0.0
	MALE	(ALL)	15-19 YRS	34	37	61	54	43	199.7	217.3	358.3	317.2	252.6
			20-24 YRS	165	242	261	271	194	595.5	873.4	942.0	978.1	700.2
			25-29 YRS	265	314	324	396	286	564.2	668.5	689.8	843.1	608.9
			30-34 YRS	312	388	338	324	295	643.2	799.8	696.8	667.9	608.1
			35-39 YRS	342	387	453	396	296	847.0	958.5	1122.0	980.8	733.1
			40-44 YRS	223	265	297	357	295	665.2	790.5	885.9	1064.9	879.9
			45-54 YRS	164	228	260	271	247	292.2	406.2	463.2	482.8	440.0
			55-64 YRS	23	44	47	43	57	71.7	137.1	146.5	134.0	177.6
			65+ YRS	9	5	10	8	5	20.3	11.3	22.5	18.0	11.3

### Breakdown by SEX AND AGE

		Reported cases						Incidence rate				
		2003	2004	2005	2006	2007	2003	2004	2005	2006	2007	
EARLY SYPHILIS FEMALE (ALL)	15-19 YRS	*	0	*	0	0	*	0.0	*	0.0	0.0	
	20-24 YRS	*	*	*	*	*	*	*	*	*	*	
	25-29 YRS	*	0	0	0	*	*	0.0	0.0	0.0	*	
	30-34 YRS	*	*	0	0	0	*	*	0.0	0.0	0.0	
	35-39 YRS	*	0	*	0	*	*	0.0	*	0.0	*	
	40-44 YRS	*	*	0	*	*	*	*	0.0	*	*	
	45-54 YRS	0	0	*	*	*	0.0	0.0	*	*	*	
	55-64 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	
	65+ YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	
	MALE (ALL)	15-19 YRS	*	0	*	*	*	0.0	*	*	*	
	20-24 YRS	22	26	14	18	16	79.4	93.8	50.5	65.0	57.7	
	25-29 YRS	47	62	39	40	33	100.1	132.0	83.0	85.2	70.3	
	30-34 YRS	122	96	66	54	37	251.5	197.9	136.1	111.3	76.3	
	35-39 YRS	135	126	99	99	68	334.4	312.1	245.2	245.2	168.4	
	40-44 YRS	95	135	94	87	73	283.4	402.7	280.4	259.5	217.7	
	45-54 YRS	76	82	89	86	94	135.4	146.1	158.5	153.2	167.5	
	55-64 YRS	15	21	14	24	22	46.7	65.4	43.6	74.8	68.6	
	65+ YRS	*	*	*	5	*	*	*	*	11.3	*	

#### Breakdown by RACE AND SEX

	Cases of	Gender	Race	Reported cases					Incidence rate				
				2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
CHLAMYDIA	FEMALE	ASIAN/PI	(ALL)	308	284	312	331	318	232.0	213.9	235.0	249.3	239.5
		BLACK	(ALL)	467	465	462	586	521	1443.6	1437.4	1428.2	1811.5	1610.5
		HISPANIC	(ALL)	271	304	305	333	268	524.7	588.6	590.5	644.7	518.9
		NATV AMER	(ALL)	16	9	17	10	13	766.4	431.1	814.3	479.0	622.7
		WHITE	(ALL)	170	235	242	256	247	106.5	147.3	151.6	160.4	154.8
	MALE	ASIAN/PI	(ALL)	178	185	176	147	220	148.8	154.7	147.1	122.9	183.9
		BLACK	(ALL)	361	418	388	419	370	1138.1	1317.7	1223.2	1320.9	1166.4
		HISPANIC	(ALL)	261	278	302	308	318	451.1	480.5	522.0	532.4	549.6
		NATV AMER	(ALL)	12	7	9	5	11	485.6	283.2	364.2	202.3	445.1
		WHITE	(ALL)	615	676	723	782	785	343.0	377.0	403.2	436.1	437.8
GONORRHEA	FEMALE	ASIAN/PI	(ALL)	22	17	18	29	23	16.6	12.8	13.6	21.8	17.3
		BLACK	(ALL)	115	96	159	185	125	355.5	296.8	491.5	571.9	386.4
		HISPANIC	(ALL)	21	26	36	28	31	40.7	50.3	69.7	54.2	60.0
		NATV AMER	(ALL)	*	5	9	*	*	*	239.5	431.1	*	*
		WHITE	(ALL)	38	45	62	45	56	23.8	28.2	38.9	28.2	35.1
	MALE	ASIAN/PI	(ALL)	106	127	143	142	111	88.6	106.2	119.6	118.7	92.8
		BLACK	(ALL)	212	267	355	359	295	668.3	841.7	1119.1	1131.8	930.0
		HISPANIC	(ALL)	201	241	299	311	250	347.4	416.6	516.8	537.6	432.1
		NATV AMER	(ALL)	7	14	15	8	7	283.2	566.5	606.9	323.7	283.2
		WHITE	(ALL)	863	1002	1012	1120	894	481.2	558.8	564.3	624.6	498.5
EARLY SYPHILIS	FEMALE	ASIAN/PI	(ALL)	*	0	*	0	0	*	0.0	*	0.0	0.0
		BLACK	(ALL)	*	*	*	*	*	*	*	*	*	*
		HISPANIC	(ALL)	7	*	*	*	*	13.6	*	*	*	*
		NATV AMER	(ALL)	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0
		WHITE	(ALL)	*	*	*	*	0	*	*	*	*	0.0
	MALE	ASIAN/PI	(ALL)	39	48	33	27	24	32.6	40.1	27.6	22.6	20.1
		BLACK	(ALL)	33	49	27	49	33	104.0	154.5	85.1	154.5	104.0
		HISPANIC	(ALL)	107	112	87	73	53	184.9	193.6	150.4	126.2	91.6
		NATV AMER	(ALL)	*	6	7	6	*	*	242.8	283.2	242.8	*
		WHITE	(ALL)	329	317	254	244	223	183.5	176.8	141.6	136.1	124.4

San Francisco Department of Public Health

Breakdown by AGE RACE AND SEX

Cases of CHLAMYDIA	Gender Race	Age	Reported cases					Incidence rate				
			2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
FEMALE ASIAN/PI		15-19 YRS	76	46	53	55	56	1105.9	669.4	771.3	800.4	814.9
		20-24 YRS	90	89	108	116	93	944.8	934.3	1133.8	1217.8	976.3
		25-29 YRS	51	63	58	58	63	414.4	511.9	471.3	471.3	511.9
		30-34 YRS	36	34	30	38	46	328.5	310.3	273.8	346.8	419.8
		35-39 YRS	19	19	26	22	21	180.4	180.4	246.9	208.9	199.4
		40-44 YRS	18	12	15	20	17	181.0	120.7	150.9	201.1	171.0
		45-54 YRS	14	14	14	16	12	74.3	74.3	74.3	84.9	63.7
		55-64 YRS	*	*	6	*	8	*	*	47.3	*	63.0
		65+ YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0
BLACK		15-19 YRS										13596.
			221	205	204	283	232	10618.0	9849.3	9801.2	8	11146.5
		20-24 YRS	145	152	132	151	163	7150.6	7495.8	6509.5	7446.5	8038.3
		25-29 YRS	55	53	54	81	62	2472.7	2382.8	2427.8	3641.6	2787.4
		30-34 YRS	15	23	31	23	21	645.2	989.3	1333.4	989.3	903.3
		35-39 YRS	6	10	12	13	11	250.5	417.4	500.9	542.7	459.2
		40-44 YRS	7	*	6	7	*	278.7	*	238.9	278.7	*
		45-54 YRS	5	*	6	9	8	115.6	*	138.8	208.2	185.0
		55-64 YRS	*	*	*	0	*	*	*	*	0.0	*
		65+ YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0
HISPANIC		15-19 YRS	82	68	69	107	69	2419.6	2006.5	2036.0	3157.3	2036.0
		20-24 YRS	96	117	107	95	101	2028.7	2472.5	2261.2	2007.6	2134.4
		25-29 YRS	51	64	73	78	48	891.1	1118.3	1275.6	1362.9	838.7
		30-34 YRS	15	25	29	24	22	299.4	499.0	578.8	479.0	439.1
		35-39 YRS	10	15	12	12	16	247.9	371.8	297.5	297.5	396.6
		40-44 YRS	*	7	8	8	*	*	191.8	219.2	219.2	*
		45-54 YRS	*	*	*	*	5	*	*	*	*	91.5
		55-64 YRS	*	0	*	0	0	*	0.0	*	0.0	0.0
		65+ YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0
NATV AMER		15-19 YRS	*	*	6	*	*	*	*	6242.6	*	*
		20-24 YRS	8	*	*	*	*	4124.5	*	*	*	*
		25-29 YRS	*	*	5	*	*	*	*	1776.9	*	*
		30-34 YRS	*	0	*	*	0	*	0.0	*	*	0.0
		35-39 YRS	0	0	0	*	*	0.0	0.0	0.0	*	*
		40-44 YRS	0	*	0	0	*	0.0	*	0.0	0.0	*
		45-54 YRS	0	0	0	0	*	0.0	0.0	0.0	0.0	*
		55-64 YRS	0	0	*	0	0	0.0	0.0	*	0.0	0.0
		65+ YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0
WHITE		15-19 YRS	40	47	43	46	42	1086.4	1276.5	1167.8	1249.3	1140.7
		20-24 YRS	67	92	106	100	109	582.2	799.4	921.0	868.9	947.1
		25-29 YRS	36	47	52	72	56	153.9	200.9	222.3	307.7	239.3
		30-34 YRS	15	23	20	17	21	69.5	106.6	92.7	78.8	97.4
		35-39 YRS	*	11	14	9	10	*	75.9	96.6	62.1	69.0
		40-44 YRS	*	7	*	5	*	*	61.3	*	43.8	*
		45-54 YRS	*	5	*	*	5	*	22.4	*	*	22.4
		55-64 YRS	*	*	0	*	0	*	*	0.0	*	0.0
		65+ YRS	0	0	*	0	0	0.0	0.0	0.0	0.0	0.0
MALE	ASIAN/PI	15-19 YRS	15	13	13	15	12	209.9	181.9	181.9	209.9	167.9
		20-24 YRS	32	47	41	29	47	357.7	525.3	458.3	324.1	525.3
		25-29 YRS	51	32	33	21	41	448.1	281.2	289.9	184.5	360.2
		30-34 YRS	33	40	32	23	29	322.3	390.7	312.6	224.7	283.3
		35-39 YRS	20	26	30	18	41	203.7	264.7	305.5	183.3	417.5
		40-44 YRS	13	15	15	21	19	141.3	163.1	163.1	228.3	206.6
		45-54 YRS	11	10	10	14	22	67.1	61.0	61.0	85.3	134.1
		55-64 YRS	*	0	*	*	8	*	0.0	*	*	78.1
		65+ YRS	*	0	0	*	*	*	0.0	0.0	*	*
BLACK		15-19 YRS	92	90	83	110	82	4683.9	4582.1	4225.7	5600.3	4174.8
		20-24 YRS	114	133	125	103	101	6056.7	7066.2	6641.1	5472.3	5366.0
		25-29 YRS	57	81	78	84	62	2605.1	3702.0	3564.9	3839.2	2833.7

## Breakdown by AGE RACE AND SEX

		Reported cases						Incidence rate					
		2003		2004		2005		2006	2007	2003	2004	2005	2006
		30-34 YRS	34	41	39	41	44	1316.8	1587.9	1510.5	1587.9	1704.1	2007
		35-39 YRS	29	34	27	38	31	1016.4	1191.6	946.3	1331.8	1086.5	
		40-44 YRS	19	28	19	20	23	656.7	967.8	656.7	691.3	795.0	
		45-54 YRS	12	8	15	18	15	247.2	164.8	309.0	370.8	309.0	
		55-64 YRS	*	0	0	*	9	*	0.0	0.0	*	312.8	
		65+ YRS	*	0	0	0	0	*	0.0	0.0	0.0	0.0	
HISPANIC	15-19 YRS	29	29	28	33	33	713.6	713.6	689.0	812.0	812.0		
	20-24 YRS	76	67	81	72	66	1230.4	1084.7	1311.3	1165.6	1068.5		
	25-29 YRS	55	65	67	77	64	723.6	855.2	881.5	1013.0	842.0		
	30-34 YRS	37	29	48	45	50	514.9	403.6	668.0	626.2	695.8		
	35-39 YRS	32	49	39	43	40	575.1	880.7	700.9	772.8	718.9		
	40-44 YRS	21	22	20	25	35	482.1	505.1	459.1	573.9	803.5		
	45-54 YRS	7	13	17	12	27	121.0	224.7	293.9	207.4	466.7		
	55-64 YRS	*	*	*	*	*	*	*	*	*	*	*	
	65+ YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	
	NATV AMER	15-19 YRS	0	0	*	0	0	0.0	0.0	*	0.0	0.0	
WHITE	20-24 YRS	*	*	0	*	*	*	*	*	0.0	*	*	
	25-29 YRS	*	*	*	0	0	*	*	*	*	0.0	0.0	
	30-34 YRS	*	*	*	*	*	*	*	*	*	*	*	
	35-39 YRS	5	*	*	*	*	1655.2	*	*	*	*	*	
	40-44 YRS	0	*	*	0	*	0.0	*	*	*	0.0	*	
	45-54 YRS	0	0	*	*	*	0.0	0.0	*	*	*	*	
	55-64 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	
	65+ YRS	0	0	0	*	0	0.0	0.0	0.0	0.0	*	0.0	
	15-19 YRS	9	15	6	8	9	252.8	421.3	168.5	224.7	252.8		
	20-24 YRS	53	77	79	89	73	523.9	761.2	780.9	879.8	721.6		
	25-29 YRS	112	103	126	130	128	449.4	413.3	505.5	521.6	513.6		
	30-34 YRS	137	152	114	105	132	495.3	549.5	412.2	379.6	477.2		
	35-39 YRS	137	150	149	174	124	639.5	700.2	695.5	812.2	578.8		
	40-44 YRS	85	98	119	144	147	515.5	594.4	721.7	873.4	891.6		
	45-54 YRS	67	61	101	108	145	236.9	215.7	357.1	381.9	512.7		
	55-64 YRS	11	19	27	17	23	70.5	121.8	173.1	109.0	147.5		
	65+ YRS	*	*	*	*	*	*	*	*	*	*	*	

San Francisco Department of Public Health

Breakdown by AGE RACE AND SEX

GONORRHEA	FEMALE	ASIAN/PI	Reported cases							Incidence rate						
			2003 2004 2005 2006 2007					2003 2004 2005 2006 2007								
			15-19 YRS	20-24 YRS	25-29 YRS	30-34 YRS	35-39 YRS	40-44 YRS	45-54 YRS	55-64 YRS	65+ YRS	15-19 YRS	20-24 YRS	25-29 YRS		
BLACK			8	*	6	5	*	116.4	*	0.0	0.0	87.3	72.8	*	*	
			*	7	5	10	9	*	40.6	*	0.0	73.5	52.5	105.0	94.5	
			5	*	*	5	*	*	*	*	*	*	*	40.6	*	
			*	*	*	*	*	*	*	*	*	*	*	*	*	
			*	*	*	*	0	*	*	*	*	*	*	*	0.0	
			0	*	0	*	*	*	0.0	*	*	0.0	0.0	*	*	
			*	*	*	*	*	*	*	*	*	*	*	*	*	
			0	0	*	*	*	0.0	0.0	0.0	*	*	*	*	*	
			0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
			45	32	62	72	43	2162.0	1537.4	2978.8	3459.3	2065.9	2219.2	3008.2	1627.4	
HISPANIC			33	34	45	61	33	1627.4	1676.7	674.4	989.1	1079.0	305.1	430.1	387.1	*
			19	12	15	22	24	854.2	539.5	375.7	250.5	250.5	199.1	185.0	*	*
			6	7	10	10	9	258.1	301.1	157.3	126.8	147.9	147.9	126.8	147.9	*
			*	*	9	6	6	*	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			*	*	5	*	*	*	*	*	*	*	*	*	*	
			*	*	8	*	*	*	*	*	*	*	*	*	*	
			0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
			*	0	0	0	0	*	*	0.0	0.0	0.0	0.0	0.0	0.0	
			*	*	12	16	6	*	265.6	354.1	472.1	177.0	157.3	126.8	147.9	
			6	7	7	6	7	126.8	147.9	147.9	126.8	147.9	123.9	126.8	147.9	*
NATV AMER			*	*	*	0	*	*	*	*	*	*	0.0	*	*	
			*	*	*	*	*	*	*	*	*	*	*	*	*	
			*	*	*	0	*	*	*	*	*	*	*	*	*	
			0	*	*	*	5	0.0	*	*	*	*	*	*	*	
			*	*	*	*	*	*	*	*	*	*	*	*	*	
			*	*	*	0	*	*	*	*	*	*	*	*	*	
			0	*	0	0	0	0.0	*	0.0	0.0	0.0	0.0	0.0	0.0	
			0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
			*	*	*	0	0	0.0	*	*	*	*	*	*	*	
			*	*	*	*	*	*	*	*	*	*	*	*	*	
WHITE			15-19 YRS	5	8	9	5	10	135.8	217.3	244.4	135.8	271.6	224.4	178.8	167.7
			20-24 YRS	8	13	20	11	12	69.5	113.0	173.8	95.6	104.3	113.0	173.8	95.6
			25-29 YRS	9	7	14	16	11	38.5	29.9	59.8	68.4	47.0	37.1	*	51.0
			30-34 YRS	6	*	8	*	11	27.8	*	0.0	0.0	0.0	0.0	0.0	0.0
			35-39 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			40-44 YRS	*	0	*	0	*	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			45-54 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			55-64 YRS	0	0	*	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			65+ YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			*	*	*	*	*	*	*	*	*	*	*	*	*	
MALE			15-19 YRS	*	*	*	*	*	*	*	*	*	*	*	*	*
			20-24 YRS	6	19	28	16	15	67.1	212.4	313.0	178.8	167.7	1527.4	1578.3	1069.2
			25-29 YRS	27	30	34	29	21	237.2	263.6	298.7	254.8	184.5	2056.7	2559.4	2559.4
			30-34 YRS	17	27	34	43	32	166.1	263.7	332.1	420.0	312.6	224.0	217.5	130.5
			35-39 YRS	29	24	22	20	22	295.3	244.4	224.0	203.7	224.0	141.3	130.5	120.0
			40-44 YRS	16	13	12	20	12	174.0	141.3	130.5	217.5	130.5	141.3	178.8	167.7
			45-54 YRS	6	9	8	9	5	36.6	54.9	48.8	54.9	30.5	34.5	34.5	34.5
			55-64 YRS	*	*	*	*	*	*	*	*	*	*	*	*	*
			65+ YRS	0	0	0	0	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
			*	*	*	*	*	*	*	*	*	*	*	*	*	
BLACK			15-19 YRS	19	15	30	31	21	967.3	763.7	1527.4	1578.3	1069.2	1200.6	1665.4	1549.2
			20-24 YRS	26	54	64	58	52	1381.4	2869.0	3400.3	3081.5	2762.7	2056.7	2559.4	2559.4
			25-29 YRS	34	45	56	69	56	1553.9	2559.4	3153.6	2559.4	2559.4	1781.6	1665.4	1665.4
			30-34 YRS	46	31	43	40	43	1781.6	1200.6	1665.4	1549.2	1665.4	1665.4	1665.4	1665.4

## Breakdown by AGE RACE AND SEX

		Reported cases						Incidence rate				
		2003 2004 2005 2006 2007					2003	2004	2005	2006	2007	
		35-39 YRS	26	35	57	48	33	911.2	1226.7	1997.7	1682.3	1156.6
HISPANIC	40-44 YRS	26	39	43	58	35	898.7	1348.0	1486.3	2004.7	1209.8	
	45-54 YRS	31	37	48	48	41	638.5	762.1	988.7	988.7	844.5	
	55-64 YRS	*	10	10	6	12	*	347.6	347.6	208.5	417.1	
	65+ YRS	0	*	*	*	0	0.0	*	*	*	0.0	
	15-19 YRS	5	7	9	9	12	123.0	172.2	221.5	221.5	295.3	
	20-24 YRS	38	30	44	56	43	615.2	485.7	712.3	906.6	696.1	
	25-29 YRS	37	63	74	83	51	486.8	828.8	973.6	1092.0	671.0	
	30-34 YRS	48	54	47	50	44	668.0	751.5	654.0	695.8	612.3	
	35-39 YRS	34	49	65	57	43	611.1	880.7	1168.2	1024.4	772.8	
	40-44 YRS	24	20	32	35	29	551.0	459.1	734.6	803.5	665.7	
NATV AMER	45-54 YRS	15	16	24	20	23	259.3	276.6	414.9	345.7	397.6	
	55-64 YRS	0	*	*	*	*	0.0	*	*	*	*	
	65+ YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	
	15-19 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	
	20-24 YRS	*	*	*	*	*	*	*	*	*	*	
	25-29 YRS	0	*	*	*	*	0.0	*	*	*	*	
	30-34 YRS	*	7	6	0	*	*	2037.8	1746.7	0.0	*	
	35-39 YRS	*	*	5	*	*	*	*	1655.2	*	*	
	40-44 YRS	*	*	*	*	*	*	*	*	*	*	
	45-54 YRS	0	*	0	0	0	0.0	*	0.0	0.0	0.0	
WHITE	55-64 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	
	65+ YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	
	15-19 YRS	5	7	13	6	*	140.4	196.6	365.2	168.5	*	
	20-24 YRS	83	108	100	123	64	820.5	1067.6	988.5	1215.9	632.7	
	25-29 YRS	152	143	128	177	136	609.9	573.7	513.6	710.2	545.7	
	30-34 YRS	167	207	167	163	149	603.8	748.4	603.8	589.3	538.7	
	35-39 YRS	218	224	249	229	167	1017.6	1045.6	1162.3	1068.9	779.5	
	40-44 YRS	129	146	172	215	184	782.4	885.5	1043.2	1304.0	1116.0	
	45-54 YRS	89	133	153	171	156	314.7	470.2	541.0	604.6	551.6	
	55-64 YRS	16	30	25	29	30	102.6	192.4	160.3	186.0	192.4	
	65+ YRS	*	*	5	5	*	*	*	25.1	25.1	*	

San Francisco Department of Public Health

Breakdown by AGE RACE AND SEX

			Reported cases					Incidence rate										
			2003 2004 2005 2006 2007					2003 2004 2005 2006 2007										
			15-19 YRS	20-24 YRS	25-29 YRS	30-34 YRS	35-39 YRS	40-44 YRS	45-54 YRS	55-64 YRS	65+ YRS	15-19 YRS	20-24 YRS	25-29 YRS	30-34 YRS	35-39 YRS	40-44 YRS	55-64 YRS
EARLY SYPHILIS FEMALE ASIAN/PI			0	0	*	0	0	0.0	0.0	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			20-24 YRS	*	0	0	0	0	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			25-29 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			30-34 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			35-39 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			40-44 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			45-54 YRS	0	0	*	0	0	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0	0.0	0.0
			55-64 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			65+ YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
BLACK			15-19 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			20-24 YRS	0	*	0	*	0	0.0	*	0.0	0.0	*	0.0	0.0	0.0	0.0	0.0
			25-29 YRS	0	0	0	0	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
			30-34 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			35-39 YRS	*	0	0	0	0	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			40-44 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			45-54 YRS	0	0	*	0	*	0.0	0.0	0.0	0.0	*	0.0	0.0	*	0.0	*
			55-64 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			65+ YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
HISPANIC			15-19 YRS	*	0	0	0	0	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			20-24 YRS	*	0	*	0	*	*	0.0	*	0.0	*	0.0	0.0	*	0.0	*
			25-29 YRS	*	0	0	0	0	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			30-34 YRS	*	0	0	0	0	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			35-39 YRS	0	0	*	0	*	0.0	0.0	0.0	*	0.0	*	0.0	0.0	*	*
			40-44 YRS	*	*	0	0	*	*	*	*	*	0.0	0.0	0.0	0.0	*	*
			45-54 YRS	0	0	0	*	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0
			55-64 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			65+ YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
NATV AMER			15-19 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			20-24 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			25-29 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			30-34 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			35-39 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			40-44 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			45-54 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			55-64 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			65+ YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
WHITE			15-19 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			20-24 YRS	*	0	0	0	0	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			25-29 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			30-34 YRS	0	*	0	0	0	0.0	*	0.0	*	0.0	0.0	0.0	0.0	0.0	0.0
			35-39 YRS	0	0	*	0	0	0.0	0.0	0.0	*	0.0	*	0.0	0.0	0.0	0.0
			40-44 YRS	0	0	0	*	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0
			45-54 YRS	0	0	*	0	0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0
			55-64 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			65+ YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
MALE	ASIAN/PI		15-19 YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			20-24 YRS	0	*	*	0	*	0.0	*	*	*	0.0	*	0.0	*	0.0	*
			25-29 YRS	9	10	*	6	*	79.1	87.9	*	52.7	*	52.7	*	52.7	*	*
			30-34 YRS	16	10	11	*	*	156.3	97.7	107.4	*	*	*	*	*	*	*
			35-39 YRS	5	16	6	8	*	50.9	162.9	61.1	81.5	*	81.5	*	81.5	*	*
			40-44 YRS	7	7	6	*	5	76.1	76.1	65.2	*	*	*	*	54.4	54.4	*
			45-54 YRS	*	*	*	6	6	*	*	*	*	*	*	36.6	36.6	36.6	*
			55-64 YRS	*	*	*	0	*	*	*	*	*	*	*	0.0	0.0	*	*
			65+ YRS	0	0	*	0	0	0.0	0.0	*	0.0	*	0.0	*	0.0	0.0	0.0

## Breakdown by AGE RACE AND SEX

			Reported cases					Incidence rate				
			2003 2004 2005 2006 2007				2003	2004	2005	2006	2007	
			2003	2004	2005	2006						
BLACK	15-19 YRS	0	0	*	*	*	0.0	0.0	*	*	*	
	20-24 YRS	*	*	*	*	*	*	*	*	*	*	
	25-29 YRS	*	*	*	*	*	*	*	*	*	*	
	30-34 YRS	*	12	*	8	*	*	464.8	*	309.8	*	
	35-39 YRS	10	10	5	10	6	350.5	350.5	175.2	350.5	210.3	
	40-44 YRS	7	14	9	10	7	242.0	483.9	311.1	345.6	242.0	
	45-54 YRS	8	6	6	12	7	164.8	123.6	123.6	247.2	144.2	
	55-64 YRS	*	*	0	*	*	*	*	0.0	*	*	
	65+ YRS	0	0	0	0	*	0.0	0.0	0.0	0.0	*	
	NATV AMER	15-19 YRS	*	0	0	0	0	*	0.0	0.0	0.0	
HISPANIC	20-24 YRS	12	9	8	*	*	194.3	145.7	129.5	*	*	
	25-29 YRS	14	22	12	14	9	184.2	289.4	157.9	184.2	118.4	
	30-34 YRS	24	22	15	9	5	334.0	306.2	208.7	125.2	69.6	
	35-39 YRS	27	24	15	16	11	485.3	431.3	269.6	287.6	197.7	
	40-44 YRS	17	19	21	14	10	390.3	436.2	482.1	321.4	229.6	
	45-54 YRS	9	15	15	14	14	155.6	259.3	259.3	242.0	242.0	
	55-64 YRS	*	*	*	*	*	*	*	*	*	*	
	65+ YRS	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0	
	WHITE	15-19 YRS	0	0	*	0	*	0.0	0.0	*	0.0	*
	20-24 YRS	8	9	*	12	7	79.1	89.0	*	118.6	69.2	
	25-29 YRS	23	26	20	17	16	92.3	104.3	80.2	68.2	64.2	
	30-34 YRS	75	48	32	32	24	271.2	173.5	115.7	115.7	86.8	
	35-39 YRS	91	73	66	59	45	424.8	340.8	308.1	275.4	210.1	
	40-44 YRS	62	88	57	56	50	376.0	533.7	345.7	339.6	303.3	
	45-54 YRS	57	55	61	47	61	201.5	194.5	215.7	166.2	215.7	
	55-64 YRS	10	17	12	17	19	64.1	109.0	76.9	109.0	121.8	
	65+ YRS	*	*	*	*	0	*	*	*	*	0.0	

\* less than five cases; exact figure suppressed for confidentiality.

**Table 36. Adolescent vs. adult major STD cases and rates by all demographic combinations, 2003-2007.**

Breakdown by AGE, Age group is ADOLESCENT (14-20 YRS)

Cases of	Gender	Race/ethnicity	Reported cases						Incidence rate			
			2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
CHLAMYDIA	(BOTH)	(ALL)	920	863	918	1,062	954	1926.8	1807.4	1922.6	2224.2	1997.7
GONORRHEA	(BOTH)	(ALL)	151	155	228	231	194	316.2	324.6	477.5	483.8	405.4
EARLY SYPHILIS	(BOTH)	(ALL)	8	*	5	8	*	16.8	*	10.5	16.8	*

Breakdown by AGE, Age group is ADULT (21+ YRS)

Cases of	Gender	Race/ethnicity	Reported cases						Incidence rate			
			2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
CHLAMYDIA	(BOTH)	(ALL)	2,405	2,774	2,766	2,958	2,926	375.2	432.8	431.5	461.5	456.5
GONORRHEA	(BOTH)	(ALL)	1,643	1,987	2,182	2,227	1,900	256.3	310.0	340.4	347.5	296.4
EARLY SYPHILIS	(BOTH)	(ALL)	517	550	422	412	367	80.7	85.8	65.8	64.3	57.3

Breakdown by RACE AND AGE, Age group is ADOLESCENT (14-20 YRS)

Cases of	Gender	Race/ethnicity	Reported cases						Incidence rate			
			2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
CHLAMYDIA	(BOTH)	ASIAN/PI	120	93	104	98	117	602.3	466.8	522.0	491.9	587.1
		BLACK	381	365	367	486	383	6725.9	6443.4	6478.7	8579.5	6763.8
		HISPANIC	149	140	128	174	127	1403.3	1318.5	1205.5	1638.7	1196.5
		NATV AMER	5	*	7	*	6	1847.9	*	2587.1	*	2235.9
		WHITE	62	88	93	84	87	578.7	821.4	868.0	784.0	809.3
GONORRHEA	(BOTH)	ASIAN/PI	10	10	13	16	14	50.2	50.2	65.3	80.3	70.9
		BLACK	79	66	121	132	103	1394.6	1165.1	2136.0	2330.2	1815.5
		HISPANIC	16	22	23	34	28	150.7	207.2	216.6	320.2	265.9
		NATV AMER	*	*	*	0	0	*	*	*	0.0	0.0
		WHITE	26	27	36	27	18	242.7	252.0	336.0	252.0	169.4
EARLY SYPHILIS	(BOTH)	ASIAN/PI	0	0	*	0	0	0.0	0.0	*	0.0	0.0
		BLACK	0	0	*	*	*	0.0	0.0	*	*	*
		HISPANIC	6	0	0	*	*	56.5	0.0	0.0	*	*
		NATV AMER	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0
		WHITE	*	*	*	*	0	*	*	*	*	0.0

Breakdown by RACE AND AGE, Age group is ADULT (21+ YRS)

Cases of	Gender	Race/ethnicity	Reported cases						Incidence rate			
			2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
CHLAMYDIA	(BOTH)	ASIAN/PI	368	371	386	383	432	186.0	187.6	195.1	193.6	218.2
		BLACK	441	507	470	509	472	937.2	5	998.8	7	1002.8
		HISPANIC	378	439	472	458	476	477.2	554.2	595.9	578.2	600.8
		NATV AMER	23	12	19	12	18	598.6	312.3	494.5	312.3	472.4
		WHITE	720	808	862	942	855	234.3	262.9	280.5	306.5	278.2
GONORRHEA	(BOTH)	ASIAN/PI	118	136	148	155	141	59.7	68.8	74.8	78.4	71.4
		BLACK	246	297	395	404	347	522.8	631.2	839.4	858.6	737.1
		HISPANIC	203	239	307	303	254	256.3	301.7	387.6	382.5	320.8
		NATV AMER	8	17	23	10	14	208.2	442.5	598.6	260.3	367.4
		WHITE	870	1,000	1,035	1,134	928	283.1	325.4	336.8	369.0	301.8
EARLY SYPHILIS	(BOTH)	ASIAN/PI	40	48	35	27	22	20.2	24.3	17.7	13.6	11.2
		BLACK	34	49	27	47	40	72.3	104.1	57.4	99.9	85.7
		HISPANIC	108	111	88	73	60	136.3	140.1	111.1	92.2	76.4
		NATV AMER	*	5	6	6	*	*	130.1	156.2	156.2	*
		WHITE	324	319	254	243	230	105.4	103.8	82.6	79.1	74.8

## Breakdown by SEX AND AGE, Age group is ADOLESCENT (14-20 YRS)

Cases of	Gender	Race/ethnicity	Reported cases					Incidence rate				
			2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
CHLAMYDIA	FEMALE	(ALL)	683	622	664	773	704	2934.5	2672.4	2852.8	3321.2	3023.8
	MALE	(ALL)	227	230	218	255	206	927.6	939.8	890.8	1042.0	840.5
GONORRHEA	FEMALE	(ALL)	91	85	133	138	111	391.0	365.2	571.4	592.9	476.5
	MALE	(ALL)	59	68	94	90	73	241.1	277.9	384.1	367.8	296.6
EARLY SYPHILIS	FEMALE	(ALL)	*	0	*	0	0	*	0.0	*	0.0	0.0
	MALE	(ALL)	6	*	*	8	*	24.5	*	*	32.7	*

## Breakdown by SEX AND AGE, Age group is ADULT (21+ YRS)

Cases of	Gender	Race/ethnicity	Reported cases					Incidence rate				
			2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
CHLAMYDIA	FEMALE	(ALL)	980	1,140	1,089	1,262	1,153	310.5	361.2	345.1	399.9	365.5
	MALE	(ALL)	1,414	1,620	1,651	1,667	1,756	434.6	497.9	507.5	512.4	539.9
GONORRHEA	FEMALE	(ALL)	158	145	217	190	165	50.1	45.9	68.8	60.2	52.4
	MALE	(ALL)	1,480	1,840	1,958	2,028	1,726	454.9	565.6	601.8	623.3	530.6
EARLY SYPHILIS	FEMALE	(ALL)	8	*	7	*	8	2.5	*	2.2	*	2.6
	MALE	(ALL)	509	547	415	408	359	156.4	168.1	127.6	125.4	110.3

## Breakdown by AGE, RACE, AND SEX, Age group is ADOLESCENT (14-20 YRS)

Cases of	Gender	Race/ethnicity	Reported cases					Incidence rate				
			2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
CHLAMYDIA	FEMALE	ASIAN/PI	96	71	86	75	93	982.0	726.3	879.7	767.2	948.9
		BLACK	269	252	255	344	288	9346.6	8755.9	8860.2	6	10019.7
GONORRHEA	FEMALE	HISPANIC	104	93	87	121	83	2169.4	1939.9	1814.8	2524.0	1724.6
		NATV AMER	5	*	6	*	6	3688.8	*	4426.6	*	4463.3
EARLY SYPHILIS	MALE	WHITE	47	64	71	65	69	867.5	1181.2	1310.4	1199.7	1265.5
		ASIAN/PI	22	22	17	19	22	216.8	216.8	167.6	187.3	218.6
CHLAMYDIA	MALE	BLACK	112	112	108	140	93	4019.2	4019.2	3875.6	5024.0	3328.8
		HISPANIC	44	47	40	49	42	755.5	807.0	686.8	841.3	727.1
GONORRHEA	MALE	NATV AMER	0	*	*	0	0	0.0	*	*	0.0	0.0
		WHITE	15	24	21	19	16	283.2	453.2	396.5	358.8	304.6
CHLAMYDIA	FEMALE	ASIAN/PI	8	*	8	9	10	81.8	*	81.8	92.1	103.1
		BLACK	54	41	75	91	71	1876.3	1424.6	2605.9	3161.9	2452.4
GONORRHEA	FEMALE	HISPANIC	7	11	11	17	10	146.0	229.5	229.5	354.6	210.3
		NATV AMER	*	*	*	0	0	*	*	*	0.0	0.0
EARLY SYPHILIS	MALE	WHITE	7	7	15	6	8	129.2	129.2	276.9	110.7	148.9
		ASIAN/PI	*	6	5	6	*	*	59.1	49.3	59.1	*
CHLAMYDIA	MALE	BLACK	25	25	46	41	30	897.1	897.1	1650.7	1471.3	1085.5
		HISPANIC	9	11	12	17	18	154.5	188.9	206.0	291.9	311.6
GONORRHEA	MALE	NATV AMER	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0
		WHITE	19	20	21	21	10	358.8	377.6	396.5	396.5	190.4
CHLAMYDIA	FEMALE	ASIAN/PI	0	0	*	0	0	0.0	0.0	*	0.0	0.0
		BLACK	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0
GONORRHEA	FEMALE	HISPANIC	*	0	0	0	0	*	0.0	0.0	0.0	0.0
		NATV AMER	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0
EARLY SYPHILIS	MALE	WHITE	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0
		ASIAN/PI	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0
CHLAMYDIA	MALE	BLACK	0	0	*	*	*	0.0	0.0	*	*	*
		HISPANIC	*	0	0	*	*	*	0.0	0.0	*	*
GONORRHEA	MALE	NATV AMER	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0
		WHITE	*	*	*	*	0	*	*	*	*	0.0

San Francisco Department of Public Health

Breakdown by AGE, RACE, AND SEX, Age group is ADULT (21+ YRS)

Cases of	Gender	Race/ethnicity	Reported cases					Incidence rate				
			2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
CHLAMYDIA	FEMALE	ASIAN/PI	210	208	225	253	232	197.9	196.0	212.1	238.5	218.6
		BLACK	192	203	194	235	220	806.3	852.5	814.7	986.9	923.1
		HISPANIC	161	209	213	200	179	431.8	560.5	571.3	536.4	481.4
		NATV AMER	11	6	11	7	8	636.9	347.4	636.9	405.3	467.1
		WHITE	124	168	164	183	151	86.1	116.7	113.9	127.1	105.0
	MALE	ASIAN/PI	156	162	157	127	200	170.1	176.7	171.2	138.5	217.7
		BLACK	248	302	276	274	252	1067.0	1299.3	1187.5	1178.9	1084.5
		HISPANIC	216	230	259	257	296	515.2	548.6	617.8	613.0	707.0
		NATV AMER	12	6	8	5	10	567.4	283.7	378.3	236.4	476.7
		WHITE	594	640	694	752	702	363.7	391.9	424.9	460.4	429.7
GONORRHEA	FEMALE	ASIAN/PI	14	13	10	19	20	13.2	12.3	9.4	17.9	19.0
		BLACK	60	54	86	90	60	252.0	226.8	361.2	378.0	254.1
		HISPANIC	12	15	24	10	20	32.2	40.2	64.4	26.8	54.1
		NATV AMER	*	*	8	*	8	*	*	463.2	*	467.1
		WHITE	31	36	48	40	38	21.5	25.0	33.3	27.8	26.6
	MALE	ASIAN/PI	103	123	138	136	121	112.3	134.1	150.5	148.3	131.9
		BLACK	185	243	308	314	286	795.9	1045.5	1325.1	1351.0	1232.0
		HISPANIC	191	224	283	292	234	455.6	534.3	675.0	696.5	557.9
		NATV AMER	7	14	15	8	6	331.0	661.9	709.2	378.3	286.0
		WHITE	839	964	986	1,089	889	513.7	590.2	603.7	666.8	544.5
EARLY SYPHILIS	FEMALE	ASIAN/PI	*	0	*	0	0	*	0.0	*	0.0	0.0
		BLACK	*	*	*	*	*	*	*	*	*	*
		HISPANIC	5	*	*	*	*	13.4	*	*	*	*
		NATV AMER	0	0	0	0	0	0.0	0.0	0.0	0.0	0.0
		WHITE	*	*	*	*	0	*	*	*	*	0.0
	MALE	ASIAN/PI	39	48	34	27	22	42.5	52.3	37.1	29.4	24.2
		BLACK	33	48	26	45	36	142.0	206.5	111.9	193.6	156.2
		HISPANIC	103	110	86	72	56	245.7	262.4	205.1	171.7	134.7
		NATV AMER	*	5	6	6	*	*	236.4	283.7	283.7	*
		WHITE	323	318	251	242	230	197.8	194.7	153.7	148.2	140.8

\* Less than five cases: exact figures suppressed for confidentiality.

*Table 37. Adolescent cases and rates by reporting source, 2003-2007, versus adult cases and rates. Unknown and out-of-jurisdiction providers included in percentages but not listed.*

Age is ADOLESCENT (14-20 YRS)

		Reported cases					Percent of reports				
		2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
CHLAMYDIA	All				1,06						
	OOJ PROVIDERS	920	863	918	2	473	100%	100%	100%	100%	100%
	CITY CLINIC	33	16	28	22	14	3.5%	1.8%	3.0%	2.0%	2.9%
	PUBLIC CLINIC (CHN)	68	62	77	65	34	7.3%	7.1%	8.3%	6.1%	7.1%
	JAILS	37	40	28	48	10	4.0%	4.6%	3.0%	4.5%	2.1%
	PRIVATE CLINIC/PMD	108	131	113	173	54	11.7%	15.1%	12.3%	16.2%	11.4%
	PRIVATE HOSPITAL	295	263	334	376	178	32.0%	30.4%	36.3%	35.4%	37.6%
	CHPY	212	189	187	236	111	23.0%	21.9%	20.3%	22.2%	23.4%
	SFGH	64	74	82	104	54	6.9%	8.5%	8.9%	9.7%	11.4%
	OUTREACH	92	78	61	31	10	10.0%	9.0%	6.6%	2.9%	2.1%
GONORRHEA	MAGNET	11	7	4	0	8	1.1%	0.8%	0.4%	0.0%	1.6%
	All	0	3	4	7	0	0.0%	0.3%	0.4%	0.6%	0.0%
	OOJ PROVIDERS	151	155	228	231	96	100%	100%	100%	100%	100%
	CITY CLINIC	3	4	8	4	0	1.9%	2.5%	3.5%	1.7%	0.0%
	PUBLIC CLINIC (CHN)	21	36	32	40	17	13.9%	23.2%	14.0%	17.3%	17.7%
	JAILS	9	8	4	9	1	5.9%	5.1%	1.7%	3.8%	1.0%
	PRIVATE CLINIC/PMD	25	16	36	38	13	16.5%	10.3%	15.7%	16.4%	13.5%
	PRIVATE HOSPITAL	31	24	53	45	27	20.5%	15.4%	23.2%	19.4%	28.1%
	CHPY	28	31	43	47	29	18.5%	20.0%	18.8%	20.3%	30.2%
	SFGH	15	18	22	26	6	9.9%	11.6%	9.6%	11.2%	6.2%
EARLY SYPHILIS	OUTREACH	18	13	24	9	1	11.9%	8.3%	10.5%	3.8%	1.0%
	MAGNET	1	0	2	1	0	0.6%	0.0%	0.8%	0.4%	0.0%
	All	0	5	4	12	2	0.0%	3.2%	1.7%	5.1%	2.0%
	OOJ PROVIDERS	8	1	5	8	0	100%	100%	100%	100%	100%
	CITY CLINIC	0	0	0	0	0	0.0%	0.0%	0.0%	0.0%	0.0%
	PUBLIC CLINIC (CHN)	5	1	3	3	1	62.5%	100%	60.0%	37.5%	50.0%
	JAILS	0	0	0	1	0	0.0%	0.0%	0.0%	12.5%	0.0%
	PRIVATE CLINIC/PMD	1	0	0	0	0	12.5%	0.0%	0.0%	0.0%	0.0%
	PRIVATE HOSPITAL	0	0	0	2	0	0.0%	0.0%	0.0%	25.0%	0.0%
	CHPY	0	0	0	0	0	0.0%	0.0%	0.0%	0.0%	0.0%
MAGNET	SFGH	0	0	2	1	0	25.0%	0.0%	40.0%	12.5%	0.0%
	OUTREACH	0	0	0	0	0	0.0%	0.0%	0.0%	0.0%	0.0%
	All	0	0	0	0	0	0.0%	0.0%	0.0%	0.0%	0.0%

Age is ADULT (21+ YRS)

		Reported cases					Percent of reports				
		2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
CHLAMYDIA	All	2,405	2,776	2,766	2,959	1,451	100%	100%	100%	100%	100%
	OOJ PROVIDERS	78	98	116	84	40	3.2%	3.5%	4.1%	2.8%	2.7%
	CITY CLINIC	777	720	770	726	355	32.3%	25.9%	27.8%	24.5%	24.4%
	PUBLIC CLINIC (CHN)	87	134	114	118	42	3.6%	4.8%	4.1%	3.9%	2.8%
	JAILS	140	183	180	159	63	5.8%	6.5%	6.5%	5.3%	4.3%
	PRIVATE CLINIC/PMD	595	638	703	923	489	24.7%	22.9%	25.4%	31.1%	33.7%
	PRIVATE HOSPITAL	466	534	431	593	282	19.3%	19.2%	15.5%	20.0%	19.4%
	CHPY	14	23	32	19	18	0.5%	0.8%	1.1%	0.6%	1.2%
	SFGH	181	171	167	109	46	7.5%	6.1%	6.0%	3.6%	3.1%
	OUTREACH	10	8	18	23	13	0.4%	0.2%	0.6%	0.7%	0.8%
MAGNET	MAGNET	57	267	235	205	103	2.3%	9.6%	8.4%	6.9%	7.0%

San Francisco Department of Public Health

Age is ADULT (21+ YRS)

		Reported cases					Percent of reports				
		2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
		All	1,643	1,989	2,182	2,230	942	100%	100%	100%	100%
GONORRHEA	OOJ PROVIDERS		28	48	52	30	12	1.7%	2.4%	2.3%	1.3%
	CITY CLINIC		848	894	899	919	369	51.6%	44.9%	41.2%	39.1%
	PUBLIC CLINIC (CHN)		63	73	81	79	37	3.8%	3.6%	3.7%	3.5%
	JAILS		32	57	71	63	32	1.9%	2.8%	3.2%	2.8%
	PRIVATE CLINIC/PMD		308	222	294	373	183	18.7%	11.1%	13.4%	16.7%
	PRIVATE HOSPITAL		207	215	271	272	117	12.5%	10.8%	12.4%	12.1%
	CHPY		12	16	20	25	5	0.7%	0.8%	0.9%	1.1%
	SFGH		69	96	121	76	30	4.1%	4.8%	5.5%	3.4%
	OUTREACH		2	12	13	17	16	0.1%	0.6%	0.5%	0.7%
	MAGNET		74	356	360	376	141	4.5%	17.8%	16.4%	16.8%
EARLY SYPHILIS	All		517	550	422	412	182	100%	100%	100%	100%
	OOJ PROVIDERS		12	13	8	12	6	2.3%	2.3%	1.8%	2.9%
	CITY CLINIC		163	180	97	107	50	31.5%	32.7%	22.9%	25.9%
	PUBLIC CLINIC (CHN)		23	24	14	14	3	4.4%	4.3%	3.3%	3.3%
	JAILS		4	4	3	2	0	0.7%	0.7%	0.7%	0.0%
	PRIVATE CLINIC/PMD		173	136	108	140	56	33.4%	24.7%	25.5%	33.9%
	PRIVATE HOSPITAL		84	104	97	85	33	16.2%	18.9%	22.9%	20.6%
	CHPY		0	1	2	1	0	0.0%	0.1%	0.4%	0.2%
	SFGH		52	40	47	24	14	10.0%	7.2%	11.1%	5.8%
	OUTREACH		1	2	2	0	0	0.1%	0.3%	0.4%	0.0%
	MAGNET		5	46	44	27	20	0.9%	8.3%	10.4%	6.5%
											10.9%

## Appendix II. Demographic Breakdowns for City Clinic

Table 38. City Clinic visits and unduplicated patient counts by all demographic combinations, 2003-2007.

Breakdown by (NONE)

Gender	Ethnicity	Age group	Total patients					Clinic visits				
			2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
(ALL PATIENTS)	(ALL)	(ALL)	12,684	13,305	13,691	12,919	7,471	21,581	23,040	23,225	21,764	11,252

Breakdown by AGE

Gender	Ethnicity	Age group	Total patients					Clinic visits				
			2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
(ALL PATIENTS)	(ALL)	UNDER 20 YRS	411	495	565	551	260	600	753	876	805	352
		20-24 YRS	2,008	2,218	2,486	2,286	1,284	3,249	3,773	4,121	3,825	1,831
		25-29 YRS	2,713	2,929	2,944	2,829	1,708	4,586	4,963	5,088	4,757	2,566
		30-34 YRS	2,487	2,525	2,319	2,209	1,267	4,326	4,375	3,851	3,643	1,940
		35-39 YRS	2,001	1,982	2,026	1,827	1,069	3,608	3,561	3,478	3,184	1,638
		40-44 YRS	1,354	1,382	1,420	1,361	790	2,362	2,572	2,578	2,476	1,272
		45-54 YRS	1,296	1,296	1,424	1,331	809	2,177	2,211	2,404	2,214	1,217
		55-64 YRS	411	478	506	524	281	670	832	828	859	432

Breakdown by RACE

Gender	Ethnicity	Age group	Total patients					Clinic visits				
			2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
(ALL PATIENTS)	ASIAN/PI	(ALL)	1,564	1,737	1,855	1,732	1,016	2,667	2,986	3,120	2,897	1,553
	BLACK	(ALL)	1,954	2,015	2,092	2,059	1,179	3,250	3,398	3,588	3,454	1,731
	HISPANIC	(ALL)	2,284	2,460	2,474	2,399	1,382	4,241	4,663	4,681	4,526	2,277
	NATV AMER	(ALL)	91	93	81	75	47	172	176	144	126	66
	WHITE	(ALL)	6,754	6,956	7,140	6,570	3,796	11,206	11,758	11,620	10,647	5,560

San Francisco Department of Public Health

Breakdown by RACE AND AGE

Gender	Ethnicity	Age group	Total patients					Clinic visits				
			2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
Gender	Ethnicity	Age group										
(ALL PATIENTS)	ASIAN/PI	UNDER 20 YRS	80	96	104	105	52	122	145	177	158	68
		20-24 YRS	297	387	409	372	211	481	585	698	611	309
		25-29 YRS	457	455	496	447	293	790	786	821	736	418
		30-34 YRS	299	338	335	324	191	547	625	550	529	312
		35-39 YRS	212	212	238	229	111	345	390	422	439	196
		40-44 YRS	112	119	132	123	74	221	234	222	207	124
		45-54 YRS	79	105	110	96	70	118	176	182	165	106
		55-64 YRS	27	25	30	35	14	42	45	47	51	20
BLACK		UNDER 20 YRS	117	129	134	139	81	194	183	193	215	109
		20-24 YRS	324	351	384	359	210	575	605	680	661	296
		25-29 YRS	317	351	361	395	215	541	588	639	688	326
		30-34 YRS	343	321	291	293	166	558	515	471	473	255
		35-39 YRS	295	305	282	260	168	477	537	510	437	248
		40-44 YRS	245	236	246	233	131	389	414	432	404	204
		45-54 YRS	249	234	305	277	160	417	409	515	428	230
		55-64 YRS	64	88	89	103	48	99	147	148	148	63
HISPANIC		UNDER 20 YRS	105	128	155	142	61	148	214	250	212	94
		20-24 YRS	445	491	510	496	273	777	865	908	903	433
		25-29 YRS	518	565	551	527	313	964	1,102	1,088	1,010	517
		30-34 YRS	479	487	419	440	257	903	942	829	813	426
		35-39 YRS	334	367	370	354	187	727	742	712	700	325
		40-44 YRS	207	214	231	215	146	396	424	459	467	238
		45-54 YRS	155	160	184	168	116	259	296	350	326	194
		55-64 YRS	41	48	54	57	27	67	78	85	95	48
NATV AMER		UNDER 20 YRS	5	6	*	*	*	6	9	10	*	5
		20-24 YRS	19	11	9	14	10	31	22	20	28	12
		25-29 YRS	15	21	16	10	9	25	33	29	15	13
		30-34 YRS	19	20	19	17	7	29	48	34	33	9
		35-39 YRS	11	13	14	9	7	22	22	20	12	9
		40-44 YRS	12	8	14	11	*	32	20	23	16	*
		45-54 YRS	6	9	*	8	9	9	14	7	15	15
		55-64 YRS	*	5	*	*	0	18	8	*	*	0
WHITE		UNDER 20 YRS	104	134	167	160	60	130	200	245	215	74
		20-24 YRS	917	967	1,162	1,029	567	1,379	1,679	1,797	1,599	764
		25-29 YRS	1,397	1,527	1,508	1,424	868	2,256	2,442	2,492	2,279	1,277
		30-34 YRS	1,340	1,354	1,249	1,118	640	2,277	2,238	1,958	1,766	931
		35-39 YRS	1,144	1,082	1,115	965	586	2,032	1,866	1,804	1,582	849
		40-44 YRS	772	799	792	772	435	1,318	1,473	1,433	1,372	698
		45-54 YRS	803	784	816	779	449	1,368	1,311	1,345	1,276	666
		55-64 YRS	275	309	331	323	191	444	549	546	558	300

Breakdown by SEX

Gender	Ethnicity	Age group	Total patients					Clinic visits				
			2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
Gender	Ethnicity	Age group										
WOMEN	(ALL)	(ALL)	2,610	3,180	3,670	3,363	1,936	4,123	5,261	6,085	5,464	2,772
MEN	(ALL)	(ALL)	10,034	10,073	9,959	9,489	5,501	17,386	17,675	16,999	16,143	8,418
TRANS-GENDER	(ALL)	(ALL)	40	52	62	67	34	72	104	141	157	62

## Breakdown by SEX AND AGE

Gender	Ethnicity	Age group	Total patients					Clinic visits				
			2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
WOMEN	(ALL)	UNDER 20 YRS	228	280	327	322	150	348	436	528	472	201
		20-24 YRS	731	890	1,108	981	552	1,196	1,545	1,921	1,718	794
		25-29 YRS	722	870	984	869	521	1,113	1,462	1,656	1,414	751
		30-34 YRS	371	486	489	494	298	610	783	779	766	423
		35-39 YRS	259	268	336	311	193	398	415	534	486	259
		40-44 YRS	137	176	192	182	112	224	296	314	305	176
		45-54 YRS	141	165	187	152	90	210	251	285	226	134
		55-64 YRS	21	45	46	51	20	24	73	67	76	33
MEN	(ALL)	UNDER 20 YRS	182	213	236	229	110	250	314	345	333	151
		20-24 YRS	1,269	1,322	1,369	1,295	726	2,044	2,215	2,182	2,086	1,030
		25-29 YRS	1,979	2,047	1,942	1,947	1,175	3,451	3,472	3,393	3,312	1,792
		30-34 YRS	2,109	2,023	1,819	1,699	965	3,701	3,560	3,027	2,837	1,505
		35-39 YRS	1,735	1,707	1,684	1,506	868	3,195	3,137	2,936	2,675	1,363
		40-44 YRS	1,215	1,204	1,224	1,168	676	2,136	2,271	2,255	2,144	1,094
		45-54 YRS	1,153	1,127	1,229	1,173	717	1,964	1,951	2,105	1,978	1,081
		55-64 YRS	389	430	456	472	261	642	755	756	778	399
TRANS-GENDER (ALL)	(ALL)	UNDER 20 YRS	*	*	*	0	0	*	*	*	0	0
		20-24 YRS	8	6	9	10	6	9	13	18	21	7
		25-29 YRS	12	12	18	13	12	22	29	39	31	23
		30-34 YRS	7	16	11	16	*	15	32	45	40	12
		35-39 YRS	7	7	6	10	8	15	9	8	23	16
		40-44 YRS	*	*	*	11	*	*	5	9	27	*
		45-54 YRS	*	*	8	6	*	*	9	14	10	*
		55-64 YRS	*	*	*	*	0	*	*	5	5	0

## Breakdown by RACE AND SEX

Gender	Ethnicity	Age group	Total patients					Clinic visits				
			2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
WOMEN	(ALL)	ASIAN/PI (ALL)	482	573	711	656	395	760	961	1,150	1,057	577
		BLACK (ALL)	570	601	661	664	392	1,010	1,072	1,259	1,211	603
		HISPANIC (ALL)	423	507	612	548	302	661	896	1,070	941	428
		NATV AMER (ALL)	29	37	32	26	16	55	62	54	45	19
		WHITE (ALL)	1,094	1,445	1,637	1,443	813	1,623	2,246	2,526	2,168	1,119
		ASIAN/PI (ALL)	1,077	1,159	1,139	1,069	616	1,900	2,015	1,964	1,828	967
		BLACK (ALL)	1,382	1,407	1,422	1,385	783	2,236	2,312	2,312	2,229	1,120
		HISPANIC (ALL)	1,839	1,931	1,837	1,826	1,065	3,541	3,721	3,539	3,512	1,817
MEN	(ALL)	NATV AMER (ALL)	61	56	49	49	31	116	114	90	81	47
		WHITE (ALL)	5,650	5,493	5,483	5,102	2,974	9,562	9,478	9,053	8,421	4,429
		ASIAN/PI (ALL)	5	5	5	7	5	7	10	6	12	9
		BLACK (ALL)	*	7	9	10	*	*	14	17	14	8
		HISPANIC (ALL)	22	22	25	25	15	39	46	72	73	32
		NATV AMER (ALL)	*	0	0	0	0	*	0	0	0	0
		WHITE (ALL)	10	18	20	25	9	21	34	41	58	12

San Francisco Department of Public Health

Breakdown by AGE, RACE, AND SEX

Gender	Ethnicity	Age group	Total patients					Clinic visits				
			2003	2004	2005	2006	2007	2003	2004	2005	2006	2007
WOMEN	ASIAN/PI	UNDER 20 YRS	53	59	74	71	40	86	97	129	107	54
		20-24 YRS	135	180	231	199	115	214	288	365	335	170
		25-29 YRS	155	157	195	173	121	240	273	333	269	165
		30-34 YRS	71	93	101	103	69	133	168	161	157	108
		35-39 YRS	36	45	56	59	28	45	65	79	112	45
		40-44 YRS	16	19	23	27	15	21	41	33	43	23
		45-54 YRS	13	16	25	15	*	18	20	42	19	8
		55-64 YRS	*	*	5	8	*	*	9	7	14	*
	BLACK	UNDER 20 YRS	67	91	83	89	48	118	133	122	138	66
		20-24 YRS	163	170	204	199	116	304	329	427	415	175
		25-29 YRS	108	119	127	132	82	173	215	252	250	134
		30-34 YRS	70	69	71	86	41	116	119	128	142	73
		35-39 YRS	69	61	60	61	39	139	113	119	96	52
		40-44 YRS	46	46	56	44	32	78	89	113	95	54
		45-54 YRS	43	37	51	40	31	78	61	81	61	45
		55-64 YRS	*	8	9	13	*	*	13	17	14	*
HISPANIC	HISPANIC	UNDER 20 YRS	51	55	81	64	23	69	96	136	97	36
		20-24 YRS	110	149	183	169	98	176	256	336	323	147
		25-29 YRS	115	116	152	131	73	188	220	276	224	102
		30-34 YRS	64	94	76	80	45	106	176	131	128	56
		35-39 YRS	44	39	62	48	26	65	61	99	72	35
		40-44 YRS	19	22	22	26	20	32	35	35	46	30
		45-54 YRS	15	22	27	24	14	17	36	44	40	19
		55-64 YRS	5	10	9	6	*	8	16	13	11	*
	NATV AMER	UNDER 20 YRS	*	*	*	*	*	*	7	10	*	*
		20-24 YRS	10	7	*	6	*	17	14	9	11	*
		25-29 YRS	5	7	5	*	*	9	8	9	5	5
		30-34 YRS	*	*	6	5	*	*	8	6	15	*
		35-39 YRS	*	*	*	5	*	11	6	*	6	*
		40-44 YRS	*	5	8	*	0	6	10	13	*	0
WHITE	WHITE	45-54 YRS	*	*	*	*	*	6	6	*	*	*
		55-64 YRS	0	*	0	0	0	0	*	0	0	0
		UNDER 20 YRS	54	69	84	94	35	71	101	130	126	41
		20-24 YRS	309	379	478	398	212	481	649	772	618	288
		25-29 YRS	338	464	502	424	238	502	737	781	659	338
		30-34 YRS	163	224	234	216	140	252	310	351	316	183
		35-39 YRS	104	119	153	137	95	136	170	231	197	121
		40-44 YRS	52	84	81	81	44	85	121	116	114	66
		45-54 YRS	65	87	82	70	38	87	128	115	102	59
		55-64 YRS	9	19	23	23	11	9	30	30	36	22

## Breakdown by AGE, RACE, AND SEX

MEN	ASIAN/PI		Total patients					Clinic visits							
			2003		2004		2005		2006	2007	2003		2004		2005
			UNDER 20 YRS	27	37	30	34	12	36	48	48	51	51	14	
BLACK		20-24 YRS	159	207	175	171	95	263	297	329	272	138			
		25-29 YRS	301	296	300	273	171	548	510	487	466	252			
		30-34 YRS	228	244	233	219	121	414	452	388	369	203			
		35-39 YRS	176	166	182	169	81	300	324	343	324	145			
		40-44 YRS	95	100	109	96	59	199	193	189	164	101			
		45-54 YRS	66	89	85	80	66	100	156	140	145	98			
		55-64 YRS	24	20	25	27	11	39	35	40	37	16			
HISPANIC		UNDER 20 YRS	50	38	51	50	33	76	50	71	77	43			
		20-24 YRS	161	181	180	159	92	271	276	253	245	119			
		25-29 YRS	209	230	231	259	132	368	368	381	433	190			
		30-34 YRS	273	251	220	207	125	442	395	343	331	182			
		35-39 YRS	224	241	220	198	128	334	421	387	337	192			
		40-44 YRS	199	190	188	186	99	311	322	316	306	150			
		45-54 YRS	206	196	252	236	129	339	346	430	366	185			
NATV AMER		55-64 YRS	60	80	80	90	45	95	134	131	134	59			
		UNDER 20 YRS	53	72	72	78	38	77	116	111	115	58			
		20-24 YRS	333	339	321	323	172	599	602	558	573	282			
		25-29 YRS	397	445	394	391	234	767	875	798	770	401			
		30-34 YRS	409	383	337	352	209	785	745	665	657	359			
		35-39 YRS	286	326	305	301	160	652	677	610	618	289			
		40-44 YRS	187	191	208	187	124	363	388	422	410	206			
WHITE		45-54 YRS	138	137	155	143	102	239	256	303	285	175			
		55-64 YRS	36	38	45	51	24	59	62	72	84	45			
		UNDER 20 YRS	*	*	0	0	*	*	*	0	0	*			
		20-24 YRS	8	*	5	8	6	13	8	11	17	8			
		25-29 YRS	10	14	11	7	6	16	25	20	10	8			
		30-34 YRS	17	16	13	12	5	27	40	28	18	7			
		35-39 YRS	7	9	11	*	*	11	16	16	6	5			
		40-44 YRS	10	*	6	10	*	26	10	10	15	*			
		45-54 YRS	*	6	*	5	8	*	8	*	11	14			
		55-64 YRS	*	*	*	*	0	18	5	*	*	0			
		UNDER 20 YRS	50	64	83	66	25	59	98	115	89	33			
		20-24 YRS	606	585	684	628	355	896	1,024	1,025	972	476			
		25-29 YRS	1,054	1,059	999	997	626	1,743	1,691	1,697	1,611	933			
		30-34 YRS	1,176	1,126	1,011	896	500	2,022	1,923	1,596	1,441	748			
		35-39 YRS	1,039	962	961	825	488	1,895	1,695	1,572	1,379	724			
		40-44 YRS	720	714	710	685	391	1,233	1,351	1,313	1,245	632			
		45-54 YRS	738	695	730	706	409	1,281	1,180	1,223	1,167	605			
		55-64 YRS	265	288	305	299	180	431	516	512	517	278			

San Francisco Department of Public Health

Breakdown by AGE, RACE, AND SEX

			Total patients					Clinic visits									
			2003		2004		2005		2006	2007	2003		2004		2005	2006	2007
			UNDER 20 YRS	20-24 YRS	25-29 YRS	30-34 YRS	35-39 YRS	40-44 YRS	45-54 YRS	55-64 YRS	UNDER 20 YRS	20-24 YRS	25-29 YRS	30-34 YRS	35-39 YRS	40-44 YRS	45-54 YRS
TRANS-GENDER ASIAN/PI			0	0	0	0	0	0	*	0	0	0	0	0	0	0	0
		20-24 YRS	*	0	*	*	*	*	*	*	0	*	*	*	*	*	*
		25-29 YRS	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
		30-34 YRS	0	*	*	*	*	*	0	5	*	*	*	*	*	*	*
		35-39 YRS	0	*	0	*	*	0	*	0	*	0	*	0	*	6	
		40-44 YRS	*	0	0	0	0	0	*	0	0	0	0	0	0	0	0
		45-54 YRS	0	0	0	*	0	0	0	0	0	0	0	*	0	*	0
		55-64 YRS	0	*	0	0	0	0	0	*	0	0	0	0	0	0	0
BLACK		UNDER 20 YRS	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		20-24 YRS	0	0	0	*	*	*	0	0	0	0	0	*	*	*	*
		25-29 YRS	0	*	*	*	*	*	0	5	6	5	*	*	*	*	*
		30-34 YRS	0	*	0	0	0	0	0	*	0	0	0	0	0	0	0
		35-39 YRS	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
		40-44 YRS	0	*	*	*	*	0	0	*	*	*	*	*	*	*	0
		45-54 YRS	0	*	*	*	*	0	0	*	*	*	*	*	*	*	0
		55-64 YRS	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
HISPANIC		UNDER 20 YRS	*	*	*	0	0	0	*	*	*	*	*	*	*	0	0
		20-24 YRS	*	*	6	*	*	*	*	7	14	7	*	*	*	*	*
		25-29 YRS	6	*	5	5	6	9	7	14	16	14					
		30-34 YRS	6	10	6	8	*	12	21	33	28	11					
		35-39 YRS	*	*	*	5	*	10	*	*	*	10	*	*	*	*	*
		40-44 YRS	*	*	*	*	*	*	*	*	*	*	*	*	11	*	
		45-54 YRS	*	*	*	*	*	0	*	*	*	*	*	*	*	0	
		55-64 YRS	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
NATV AMER		UNDER 20 YRS	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		20-24 YRS	*	0	0	0	0	0	*	0	0	0	0	0	0	0	0
		25-29 YRS	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		30-34 YRS	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		35-39 YRS	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		40-44 YRS	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		45-54 YRS	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		55-64 YRS	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
WHITE		UNDER 20 YRS	0	*	0	0	0	0	0	*	0	0	0	0	0	0	0
		20-24 YRS	*	*	0	*	0	*	*	6	0	9	0	*	*	*	*
		25-29 YRS	5	*	7	*	*	11	14	14	9	6					
		30-34 YRS	*	*	*	6	0	*	5	11	9	0					
		35-39 YRS	*	*	*	*	*	*	*	*	6	*	*	*	*	*	*
		40-44 YRS	0	*	*	6	0	0	*	*	13	0					
		45-54 YRS	0	*	*	*	*	*	0	*	7	7	*	*	*	*	*
		55-64 YRS	*	*	*	*	0	*	*	*	5	0					

\* Less than five cases: exact figures suppressed for confidentiality.