Table 1. STDs among residents, June, 2020. Female syphilis cases include patients assigned as female at birth.

|                        | 2020  |       | 2019  |       |
|------------------------|-------|-------|-------|-------|
|                        | month | YTD   | month | YTD   |
| Gonorrhea              | 285   | 2,063 | 411   | 2,524 |
| Male rectal gonorrhea  | 82    | 597   | 117   | 680   |
| Chlamydia              | 381   | 3,053 | 747   | 4,775 |
| Male rectal chlamydia  | 111   | 867   | 202   | 1191  |
| Syphilis (adult total) | 138   | 826   | 137   | 914   |
| Primary & secondary    | 47    | 279   | 29    | 260   |
| Early latent           | 56    | 377   | 75    | 481   |
| Unknown latent         | 18    | 77    | 12    | 43    |
| Late latent            | 17    | 93    | 21    | 130   |
| Neurosyphilis          | 1     | 8     | 3     | 14    |
| Congenital syphilis    | 1     | 1     | 1     | 2     |
| Female syphilis        | 22    | 90    | 14    | 73    |




Table 2. Selected STD cases and rates for San Francisco by age and race/ethnicity, 2020 through June only. Rates equal cases per 100,000 residents per year based on 2010 US Census Data.

|                | (All race | (All races) |       | Asian/PI |       | African American |       | Hispanic |       | White    |  |
|----------------|-----------|-------------|-------|----------|-------|------------------|-------|----------|-------|----------|--|
|                | cases     | rate        | cases | rate     | cases | rate             | cases | rate     | cases | rate     |  |
| All ages       |           |             |       |          |       |                  |       |          |       |          |  |
| Chlamydia      | 3,053     | 758.3       | 364   | 274.0    | 326   | 1,393.7          | 460   | 755.5    | 901   | 534.0    |  |
| Gonorrhea      | 2,063     | 512.4       | 190   | 143.0    | 259   | 1,107.3          | 389   | 638.9    | 749   | 443.9    |  |
| Early syphilis | 656       | 162.9       | 79    | 59.5     | 88    | 376.2            | 156   | 256.2    | 254   | 150.5    |  |
| Under 20 yrs   |           |             |       |          |       |                  |       |          |       | <u>.</u> |  |
| Chlamydia      | 246       | 885.0       | 25    | 225.8    | 60    | 2,551.6          | 49    | 774.6    | 21    | 342.5    |  |
| Gonorrhea      | 59        | 212.3       | 5     | 45.2     | 20    | 850.5            | 11    | 173.9    | 4     | 65.2     |  |
| Early syphilis | 1         | 3.6         | 0     | 0.0      | 0     | 0.0              | 0     | 0.0      | 0     | 0.0      |  |

Table 3. HIV testing among City Clinic patients, June, 2020.

|                     | 2020  |       | 2019  |       |
|---------------------|-------|-------|-------|-------|
|                     | month | YTD   | month | YTD   |
| Tests               | 216   | 1,448 | 475   | 2,875 |
| Antibody positive   | 7     | 22    | 2     | 18    |
| Acute HIV infection | 0     | 0     | 0     | 3     |

Note: All statistics are provisional until the annual report is released for the year. Morbidity is based on date of diagnosis. Totals for past months may change due to delays in reporting from labs and providers.




Figure 2. City Clinic visits by gender and orientation.




Figure 3. Average number of recent\* sex partners for City Clinic visits by gender and sexual orientation. \*Recall period is 3 months.

## Syphilis continues to increase among women – please screen and ensure treatment of your patients and their partners.

We encourage all providers to screen their patients according to citywide STD screening guidelines. http://sfcityclinic.org/providers/ScreeningandSurveillance\_CitywideSTDscreeningguidance\_0619.pdf

The San Francisco Department of Public Health (SFDPH) LINCS team (comprised of disease intervention specialists) ensures patients with syphilis are treated and confidentially provides prevention treatment and tests to their sex partners.

- ❖ LINCS partner services are voluntary and offered by skilled and client-centered SFDPH staff.
- ❖ Please encourage patients to work with us at SFDPH to prevent syphilis and help them stay healthy.
- To contact LINCS, call 415-487-5531 and fax to 415-431-4628 a confidential morbidity report (<a href="https://www.sfcityclinic.org/providers/reporting">https://www.sfcityclinic.org/providers/reporting</a>).
- ❖ Visit our website for more information: <a href="https://www.sfcityclinic.org/providers">https://www.sfcityclinic.org/providers</a>

## Novel Coronavirus (COVID-19) Sexual Health Tips

Exposure to COVID-19 can occur during sexual activities through direct contact and potential exchange of saliva, semen, mucus and feces; COVID-19 has not been identified in vaginal fluids. All San Franciscans are required to wear a face covering when outside their residence and within six feet of other people who are not members of their household. To minimize exposure to COVID-19 if you do have sex with others, have as few partners as possible and avoid group sex. If you usually meet your sex partners online or make a living by having sex, consider taking a break from in-person dates. Video dates, sexting or chat rooms may be options for you.

- > Additional SFDPH COVID-19 Sexual Health Tips are available on the SFCC website.
- > SFDPH COVID-19 information for providers and patients are available online: https://www.sfdph.org/dph/alerts/coronavirus.asp
  - For SF Health Care Providers

SFDPH-developed guidance and materials can be found here: https://www.sfcdcp.org/infectious-diseases-a-to-z/coronavirus-2019-novel-coronavirus/coronavirus-2019-information-for-healthcare-providers/

For the SF Public

https://www.sfcdcp.org/infectious-diseases-a-to-z/coronavirus-2019-novel-coronavirus/