

HIV SEMI-ANNUAL SURVEILLANCE REPORT

San Francisco Department of Public Health

HIV Cases Reported Through December 2014

Contents	Page
Surveillance Summary.....	1
Figure 1: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year	2
Figure 2: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year	2
Table 1. Diagnoses of HIV infection by age and year of initial HIV diagnosis.....	3
Table 2: Diagnoses of HIV infection stage 3 (AIDS) by age and year of AIDS diagnosis.....	4
Table 3: Diagnoses of HIV infection by gender and year of initial HIV diagnosis	5
Table 4: Diagnoses of HIV infection stage 3 (AIDS) by gender and year of AIDS diagnosis...	6
Table 5: Diagnoses of HIV infection by race/ethnicity and year of initial HIV diagnosis.....	7
Table 6: Diagnoses of HIV infection stage 3 (AIDS) by race/ethnicity and year of AIDS diagnosis.....	8
Table 7: Diagnoses of HIV infection by transmission category and year of initial HIV diagnosis.....	9
Table 8: Diagnoses of HIV infection stage 3 (AIDS) by transmission category and year of AIDS diagnosis.....	10
Table 9: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year	11
Table 10: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year	12
Table 11: Persons living with HIV infection by age, gender and race/ethnicity.....	13
Table 12: Persons living with HIV infection by transmission category, gender and race/ethnicity.....	14

The HIV Surveillance Report is published semi-annually by the
San Francisco Department of Public Health, Population Health Division
Applied Research, Community Health Epidemiology, and Surveillance Branch (ARCHES)
25 Van Ness Avenue, Suite 500, San Francisco, CA 94102; Phone (415) 437-6336, FAX (415) 431-0353

Director of Health: Barbara A. Garcia, MPA
Director of Population Health Division: Tomas Aragon, MD, DrPH
HIV Surveillance Section Directors: Susan Scheer, PhD, MPH; Ling Hsu, MPH

***The HIV Surveillance Report is accessible via internet:
<http://www.sfdph.org/dph/files/reports/default.asp>***

HIV SURVEILLANCE SUMMARY

San Francisco¹ (as of 12/31/2014)

Persons living with HIV infection ²	15,964
Diagnoses of HIV infection stage 3 (AIDS), cumulative	29,673
Deaths among persons diagnosed with HIV infection stage 3 (AIDS), cumulative	20,112

California³ (as of 06/30/2014)

Persons living with HIV infection ²	122,382
Diagnoses of HIV infection stage 3 (AIDS), cumulative	169,588
Deaths among persons diagnosed with HIV infection stage 3 (AIDS), cumulative	95,529

United States⁴ (as of 12/31/2012)

Persons living with HIV infection ²	904,467
Diagnoses of HIV infection stage 3 (AIDS), cumulative (as of 12/31/2011)	1,191,061
Deaths among persons diagnosed with HIV infection stage 3 (AIDS), cumulative	659,068


¹ Residents of San Francisco at time of HIV or AIDS diagnosis.

² Persons diagnosed with HIV infection regardless of disease stage.

³ CA data source: www.cdph.ca.gov/data/statistics/Pages/OAHIVAIDSStatistics.aspx.


⁴ US data source: CDC HIV Surveillance Report 2012, vol. 24 (Table 2b, 11b, 14b).
www.cdc.gov/hiv/topics/surveillance/resources/reports/.

Figure 1: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year, San Francisco, 2002-2014*



* Reporting for recent years is incomplete. See Table 9 for actual numbers per year.

Figure 2: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year, San Francisco, 1980-2014 †



† Reporting for recent years is incomplete. See Table 10 for actual numbers per year.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2014

Table 1. Diagnoses of HIV infection by age and year of initial HIV diagnosis (1), San Francisco, 2003-2014

Age at initial HIV diagnosis (years)	Year of initial HIV diagnosis					
	2003 No. (%)	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)
0 - 12	1 (0.2)	2 (0.3)	1 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	4 (0.6)	1 (0.1)	1 (0.2)	1 (0.2)	2 (0.4)	4 (0.8)
18 - 24	49 (7.6)	62 (8.8)	50 (9.1)	61 (11.7)	56 (10.6)	51 (9.8)
25 - 29	65 (10.1)	92 (13.0)	70 (12.7)	66 (12.7)	99 (18.8)	85 (16.4)
30 - 39	286 (44.5)	283 (40.0)	209 (38.0)	175 (33.7)	189 (35.8)	185 (35.7)
40 - 49	180 (28.0)	189 (26.7)	156 (28.4)	144 (27.7)	127 (24.1)	147 (28.4)
50 - 59	50 (7.8)	62 (8.8)	51 (9.3)	56 (10.8)	42 (8.0)	39 (7.5)
60 +	8 (1.2)	17 (2.4)	12 (2.2)	17 (3.3)	13 (2.5)	7 (1.4)
Total	643 (100)	708 (100)	550 (100)	520 (100)	528 (100)	518 (100)

Age at initial HIV diagnosis (years)	Year of initial HIV diagnosis					
	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	1 (0.2)	2 (0.5)	1 (0.2)	0 (0.0)	0 (0.0)	1 (0.4)
18 - 24	57 (12.2)	55 (12.5)	44 (10.6)	53 (12.4)	48 (12.9)	32 (11.5)
25 - 29	58 (12.4)	56 (12.7)	64 (15.5)	70 (16.4)	78 (21.0)	48 (17.2)
30 - 39	145 (31.1)	137 (31.1)	107 (25.8)	131 (30.7)	109 (29.4)	83 (29.7)
40 - 49	126 (27.0)	125 (28.3)	129 (31.2)	124 (29.0)	92 (24.8)	67 (24.0)
50 - 59	62 (13.3)	50 (11.3)	54 (13.0)	38 (8.9)	31 (8.4)	34 (12.2)
60 +	17 (3.6)	16 (3.6)	15 (3.6)	11 (2.6)	13 (3.5)	14 (5.0)
Total	466 (100)	441 (100)	414 (100)	427 (100)	371 (100)	279 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of HIV antibody test, viral load or CD4 test, initiation of antiretroviral therapy, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2014

Table 2. Diagnoses of HIV infection stage 3 (AIDS) by age and year of AIDS diagnosis (1), San Francisco, 2003-2014

Age at AIDS diagnosis (years)	Year of AIDS diagnosis					
	2003 No. (%)	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)
0 - 12	1 (0.2)	1 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	1 (0.2)	1 (0.2)	0 (0.0)	0 (0.0)	1 (0.2)	1 (0.2)
18 - 24	20 (3.6)	11 (2.3)	14 (2.9)	13 (2.9)	15 (3.3)	17 (3.9)
25 - 29	34 (6.1)	21 (4.3)	32 (6.7)	25 (5.6)	39 (8.7)	31 (7.2)
30 - 39	211 (37.6)	170 (35.1)	157 (32.7)	125 (27.8)	133 (29.7)	124 (28.6)
40 - 49	196 (34.9)	165 (34.0)	183 (38.1)	183 (40.7)	167 (37.3)	164 (37.9)
50 - 59	83 (14.8)	87 (17.9)	80 (16.7)	83 (18.4)	67 (15.0)	75 (17.3)
60 +	15 (2.7)	29 (6.0)	14 (2.9)	21 (4.7)	26 (5.8)	21 (4.8)
Total	561 (100)	485 (100)	480 (100)	450 (100)	448 (100)	433 (100)

Age at AIDS diagnosis (years)	Year of AIDS diagnosis					
	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	1 (0.3)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
18 - 24	15 (4.6)	15 (5.0)	11 (4.4)	10 (4.1)	9 (4.9)	3 (2.8)
25 - 29	23 (7.1)	27 (9.0)	20 (8.0)	25 (10.4)	17 (9.2)	13 (12.0)
30 - 39	75 (23.1)	73 (24.4)	62 (24.8)	60 (24.9)	42 (22.8)	32 (29.6)
40 - 49	120 (37.0)	99 (33.1)	98 (39.2)	81 (33.6)	58 (31.5)	30 (27.8)
50 - 59	69 (21.3)	62 (20.7)	50 (20.0)	46 (19.1)	39 (21.2)	22 (20.4)
60 +	21 (6.5)	23 (7.7)	9 (3.6)	19 (7.9)	19 (10.3)	8 (7.4)
Total	324 (100)	299 (100)	250 (100)	241 (100)	184 (100)	108 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2014

Table 3. Diagnoses of HIV infection by gender and year of initial HIV diagnosis (1), San Francisco, 2003-2014

Gender	Year of initial HIV diagnosis					
	2003 No. (%)	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)
Male	574 (89.3)	641 (90.5)	491 (89.3)	470 (90.4)	462 (87.5)	460 (88.8)
Female	51 (7.9)	53 (7.5)	45 (8.2)	36 (6.9)	43 (8.1)	41 (7.9)
Transgender	18 (2.8)	14 (2.0)	14 (2.5)	14 (2.7)	23 (4.4)	17 (3.3)
Total	643 (100)	708 (100)	550 (100)	520 (100)	528 (100)	518 (100)

Gender	Year of initial HIV diagnosis					
	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)
Male	424 (91.0)	394 (89.3)	364 (87.9)	401 (93.9)	338 (91.1)	262 (93.9)
Female	25 (5.4)	34 (7.7)	41 (9.9)	22 (5.2)	23 (6.2)	12 (4.3)
Transgender	17 (3.6)	13 (2.9)	9 (2.2)	4 (0.9)	10 (2.7)	5 (1.8)
Total	466 (100)	441 (100)	414 (100)	427 (100)	371 (100)	279 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of HIV antibody test, viral load or CD4 test, initiation of antiretroviral therapy, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2014

Table 4. Diagnoses of HIV infection stage 3 (AIDS) by gender and year of AIDS diagnosis (1), San Francisco, 2003-2014

Gender	Year of AIDS diagnosis					
	2003 No. (%)	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)
Male	496 (88.4)	426 (87.8)	429 (89.4)	403 (89.6)	404 (90.2)	376 (86.8)
Female	40 (7.1)	42 (8.7)	37 (7.7)	31 (6.9)	33 (7.4)	43 (9.9)
Transgender	25 (4.5)	17 (3.5)	14 (2.9)	16 (3.6)	11 (2.5)	14 (3.2)
Total	561 (100)	485 (100)	480 (100)	450 (100)	448 (100)	433 (100)

Gender	Year of AIDS diagnosis					
	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)
Male	289 (89.2)	269 (90.0)	214 (85.6)	210 (87.1)	162 (88.0)	97 (89.8)
Female	24 (7.4)	16 (5.4)	23 (9.2)	25 (10.4)	17 (9.2)	6 (5.6)
Transgender	11 (3.4)	14 (4.7)	13 (5.2)	6 (2.5)	5 (2.7)	5 (4.6)
Total	324 (100)	299 (100)	250 (100)	241 (100)	184 (100)	108 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2014

Table 5. Diagnoses of HIV infection by race/ethnicity and year of initial HIV diagnosis (1), San Francisco, 2003-2014

Race/ethnicity	Year of initial HIV diagnosis					
	2003 No. (%)	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)
White	352 (54.7)	388 (54.8)	293 (53.3)	278 (53.5)	269 (50.9)	256 (49.4)
African American	92 (14.3)	90 (12.7)	80 (14.5)	77 (14.8)	77 (14.6)	81 (15.6)
Latino	131 (20.4)	139 (19.6)	119 (21.6)	110 (21.2)	107 (20.3)	118 (22.8)
Asian/Pacific Islander	43 (6.7)	53 (7.5)	30 (5.5)	33 (6.3)	49 (9.3)	40 (7.7)
Native American	6 (0.9)	5 (0.7)	7 (1.3)	3 (0.6)	1 (0.2)	4 (0.8)
Total (2)	643 (100)	708 (100)	550 (100)	520 (100)	528 (100)	518 (100)

Race/ethnicity	Year of initial HIV diagnosis					
	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)
White	239 (51.3)	209 (47.4)	213 (51.4)	209 (48.9)	164 (44.2)	129 (46.2)
African American	71 (15.2)	62 (14.1)	67 (16.2)	44 (10.3)	49 (13.2)	27 (9.7)
Latino	98 (21.0)	108 (24.5)	83 (20.0)	105 (24.6)	93 (25.1)	77 (27.6)
Asian/Pacific Islander	36 (7.7)	38 (8.6)	32 (7.7)	50 (11.7)	48 (12.9)	33 (11.8)
Native American	1 (0.2)	3 (0.7)	2 (0.5)	6 (1.4)	3 (0.8)	1 (0.4)
Total (2)	466 (100)	441 (100)	414 (100)	427 (100)	371 (100)	279 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of HIV antibody test, viral load or CD4 test, initiation of antiretroviral therapy, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

(2) Total includes persons with multiple or unknown race.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2014

Table 6. Diagnoses of HIV infection stage 3 (AIDS) by race/ethnicity and year of AIDS diagnosis (1), San Francisco, 2003-2014

Race/ethnicity	Year of AIDS diagnosis					
	2003 No. (%)	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)
White	276 (49.2)	257 (53.0)	269 (56.0)	252 (56.0)	257 (57.4)	220 (50.8)
African American	105 (18.7)	77 (15.9)	83 (17.3)	80 (17.8)	76 (17.0)	80 (18.5)
Latino	128 (22.8)	115 (23.7)	91 (19.0)	78 (17.3)	75 (16.7)	74 (17.1)
Asian/Pacific Islander	33 (5.9)	22 (4.5)	25 (5.2)	22 (4.9)	25 (5.6)	37 (8.5)
Native American	4 (0.7)	4 (0.8)	3 (0.6)	4 (0.9)	0 (0.0)	2 (0.5)
Total (2)	561 (100)	485 (100)	480 (100)	450 (100)	448 (100)	433 (100)

Race/ethnicity	Year of AIDS diagnosis					
	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)
White	168 (51.9)	144 (48.2)	130 (52.0)	120 (49.8)	84 (45.7)	40 (37.0)
African American	62 (19.1)	52 (17.4)	44 (17.6)	40 (16.6)	32 (17.4)	24 (22.2)
Latino	65 (20.1)	62 (20.7)	44 (17.6)	50 (20.7)	34 (18.5)	22 (20.4)
Asian/Pacific Islander	13 (4.0)	25 (8.4)	15 (6.0)	17 (7.1)	18 (9.8)	16 (14.8)
Native American	3 (0.9)	2 (0.7)	2 (0.8)	1 (0.4)	1 (0.5)	0 (0.0)
Total (2)	324 (100)	299 (100)	250 (100)	241 (100)	184 (100)	108 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

(2) Total includes persons with multiple or unknown race.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2014

Table 7. Diagnoses of HIV infection by transmission category and year of initial HIV diagnosis (1), San Francisco, 2003-2014

Transmission category (2)	Year of initial HIV diagnosis					
	2003 No. (%)	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)
Gay or bisexual male	439 (68.3)	486 (68.6)	377 (68.5)	365 (70.2)	348 (65.9)	374 (72.2)
Injection drug user (IDU)	63 (9.8)	59 (8.3)	31 (5.6)	37 (7.1)	30 (5.7)	27 (5.2)
Gay or bisexual male IDU	97 (15.1)	118 (16.7)	90 (16.4)	81 (15.6)	89 (16.9)	63 (12.2)
Lesbian or bisexual IDU	1 (0.2)	4 (0.6)	5 (0.9)	2 (0.4)	3 (0.6)	5 (1.0)
Transfusion/Hemophilia	1 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	27 (4.2)	21 (3.0)	31 (5.6)	24 (4.6)	42 (8.0)	33 (6.4)
Perinatal	1 (0.2)	2 (0.3)	1 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)
Risk not reported	14 (2.2)	18 (2.5)	15 (2.7)	11 (2.1)	16 (3.0)	16 (3.1)
Total	643 (100)	708 (100)	550 (100)	520 (100)	528 (100)	518 (100)

Transmission category (2)	Year of initial HIV diagnosis					
	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)
Gay or bisexual male	331 (71.0)	282 (63.9)	297 (71.7)	333 (78.0)	282 (76.0)	208 (74.6)
Injection drug user (IDU)	22 (4.7)	33 (7.5)	25 (6.0)	12 (2.8)	22 (5.9)	16 (5.7)
Gay or bisexual male IDU	73 (15.7)	67 (15.2)	48 (11.6)	42 (9.8)	38 (10.2)	32 (11.5)
Lesbian or bisexual IDU	4 (0.9)	1 (0.2)	3 (0.7)	2 (0.5)	1 (0.3)	2 (0.7)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	23 (4.9)	37 (8.4)	26 (6.3)	23 (5.4)	16 (4.3)	6 (2.2)
Perinatal	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Risk not reported	13 (2.8)	21 (4.8)	15 (3.6)	15 (3.5)	12 (3.2)	15 (5.4)
Total	466 (100)	441 (100)	414 (100)	427 (100)	371 (100)	279 (100)

- (1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of HIV antibody test, viral load or CD4 test, initiation of antiretroviral therapy, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.
- (2) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
- (3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2014

Table 8. Diagnoses of HIV infection stage 3 (AIDS) by transmission category and year of AIDS diagnosis (1), San Francisco, 2003-2014

Transmission category (2)	Year of AIDS diagnosis					
	2003 No. (%)	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)
Gay or bisexual male	355 (63.3)	315 (64.9)	295 (61.5)	288 (64.0)	290 (64.7)	262 (60.5)
Injection drug user (IDU)	70 (12.5)	50 (10.3)	49 (10.2)	37 (8.2)	38 (8.5)	30 (6.9)
Gay or bisexual male IDU	105 (18.7)	93 (19.2)	103 (21.5)	95 (21.1)	85 (19.0)	94 (21.7)
Lesbian or bisexual IDU	3 (0.5)	1 (0.2)	3 (0.6)	2 (0.4)	0 (0.0)	4 (0.9)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	21 (3.7)	15 (3.1)	19 (4.0)	21 (4.7)	29 (6.5)	32 (7.4)
Perinatal	1 (0.2)	1 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Risk not reported	6 (1.1)	10 (2.1)	11 (2.3)	7 (1.6)	6 (1.3)	11 (2.5)
Total	561 (100)	485 (100)	480 (100)	450 (100)	448 (100)	433 (100)

Transmission category (2)	Year of AIDS diagnosis					
	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)
Gay or bisexual male	210 (64.8)	170 (56.9)	150 (60.0)	156 (64.7)	110 (59.8)	71 (65.7)
Injection drug user (IDU)	33 (10.2)	24 (8.0)	32 (12.8)	20 (8.3)	19 (10.3)	13 (12.0)
Gay or bisexual male IDU	53 (16.4)	81 (27.1)	46 (18.4)	36 (14.9)	42 (22.8)	16 (14.8)
Lesbian or bisexual IDU	2 (0.6)	0 (0.0)	1 (0.4)	2 (0.8)	0 (0.0)	0 (0.0)
Transfusion/Hemophilia	1 (0.3)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	17 (5.2)	16 (5.4)	16 (6.4)	18 (7.5)	10 (5.4)	5 (4.6)
Perinatal	1 (0.3)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Risk not reported	7 (2.2)	8 (2.7)	5 (2.0)	9 (3.7)	3 (1.6)	3 (2.8)
Total	324 (100)	299 (100)	250 (100)	241 (100)	184 (100)	108 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

(2) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.

(3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2014

Table 9. Diagnoses of HIV infection, HIV deaths and persons living with HIV by year, San Francisco, 2002-2014

Year	Number of cases diagnosed per year (1)	Number of deaths occurred per year (1)	Number of persons living with HIV infection (1)
2002 (2)	676	350	13386
2003	643	338	13691
2004	708	344	14055
2005	550	348	14257
2006	520	327	14450
2007	528	324	14654
2008	518	264	14908
2009	466	255	15119
2010	441	247	15313
2011	414	235	15492
2012	427	236	15683
2013	371	208	15846
2014	279	161	15964

(1) Data in recent years are incomplete due to delays in case/death reporting.

(2) Year of the mandatory reporting of HIV non-AIDS cases in California began.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2014

Table 10. Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year, San Francisco, 1980-2014

Year	Number of cases diagnosed per year (1)	Number of deaths occurred per year (1)	Number of persons living with AIDS (1)
1980	2	0	2
1981	26	9	19
1982	99	32	86
1983	274	111	249
1984	558	273	534
1985	860	534	860
1986	1237	807	1290
1987	1632	878	2044
1988	1764	1040	2768
1989	2160	1279	3649
1990	2046	1364	4331
1991	2286	1512	5105
1992	2333	1641	5797
1993	2068	1603	6262
1994	1785	1601	6446
1995	1559	1485	6520
1996	1078	993	6605
1997	805	424	6986
1998	693	402	7277
1999	578	353	7502
2000	558	350	7710
2001	515	324	7901
2002	494	323	8072
2003	561	303	8330
2004	485	310	8505
2005	480	312	8673
2006	450	289	8834
2007	448	271	9011
2008	433	229	9215
2009	324	212	9327
2010	299	195	9431
2011	250	189	9492
2012	241	183	9550
2013	184	153	9581
2014	108	128	9561
Total	29673	20112	

(1) Data in recent years are incomplete due to delays in case/death reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2014

Table 11. Persons living with HIV infection by age, gender and race/ethnicity, San Francisco

Current age (years)	White		African American		Latino		Asian/ Pacific Islander/ Native American		Total (1)	
	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
Male										
0 - 12	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	1	(0.0)
13 - 17	0	(0.0)	0	(0.0)	1	(0.0)	0	(0.0)	1	(0.0)
18 - 24	26	(0.3)	20	(1.3)	32	(1.2)	16	(1.8)	97	(0.7)
25 - 29	159	(1.7)	57	(3.7)	130	(5.0)	41	(4.6)	403	(2.7)
30 - 39	752	(8.0)	134	(8.7)	490	(19.0)	191	(21.6)	1630	(11.1)
40 - 49	2361	(25.1)	374	(24.3)	834	(32.3)	315	(35.6)	3995	(27.2)
50 - 59	3688	(39.3)	575	(37.4)	796	(30.8)	230	(26.0)	5383	(36.6)
60 +	2403	(25.6)	376	(24.5)	301	(11.6)	92	(10.4)	3201	(21.8)
Subtotal	9389	(100)	1536	(100)	2584	(100)	885	(100)	14711	(100)
Female										
0 - 12	0	(0.0)	1	(0.3)	1	(0.5)	0	(0.0)	2	(0.2)
13 - 17	0	(0.0)	0	(0.0)	2	(1.1)	0	(0.0)	2	(0.2)
18 - 24	3	(1.1)	4	(1.1)	7	(3.8)	1	(1.4)	17	(1.9)
25 - 29	6	(2.3)	7	(1.9)	10	(5.5)	1	(1.4)	24	(2.7)
30 - 39	33	(12.5)	37	(10.3)	31	(17.0)	16	(22.2)	125	(13.9)
40 - 49	83	(31.4)	85	(23.6)	46	(25.3)	22	(30.6)	243	(26.9)
50 - 59	99	(37.5)	145	(40.3)	48	(26.4)	23	(31.9)	322	(35.7)
60 +	40	(15.2)	81	(22.5)	37	(20.3)	9	(12.5)	167	(18.5)
Subtotal	264	(100)	360	(100)	182	(100)	72	(100)	902	(100)
Transgender										
13 - 39	13	(17.8)	37	(31.6)	36	(32.7)	9	(23.1)	102	(29.1)
40 +	60	(82.2)	80	(68.4)	74	(67.3)	30	(76.9)	249	(70.9)
Subtotal	73	(100)	117	(100)	110	(100)	39	(100)	351	(100)
Total	9726	(100)	2013	(100)	2876	(100)	996	(100)	15964	(100)

(1) Total includes persons with multiple or unknown race.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2014

Table 12. Persons living with HIV infection by transmission category, gender and race/ethnicity, San Francisco

Transmission category (1)	White		African American		Latino		Asian/ Pacific Islander/ Native American		Total (2)	
	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
Male										
Gay or bisexual male	7635	(81.3)	882	(57.4)	2104	(81.4)	742	(83.8)	11581	(78.7)
Injection drug user (IDU)	187	(2.0)	230	(15.0)	62	(2.4)	20	(2.3)	511	(3.5)
Gay or bisexual male IDU	1461	(15.6)	313	(20.4)	312	(12.1)	92	(10.4)	2250	(15.3)
Transfusion/Hemophiliac	6	(0.1)	3	(0.2)	2	(0.1)	3	(0.3)	14	(0.1)
Heterosexual (3)	30	(0.3)	67	(4.4)	52	(2.0)	12	(1.4)	163	(1.1)
Perinatal	1	(0.0)	2	(0.1)	2	(0.1)	2	(0.2)	9	(0.1)
Risk not reported	69	(0.7)	39	(2.5)	50	(1.9)	14	(1.6)	183	(1.2)
Subtotal	9389	(100)	1536	(100)	2584	(100)	885	(100)	14711	(100)
Female										
Injection drug user (IDU)	129	(48.9)	180	(50.0)	51	(28.0)	14	(19.4)	382	(42.4)
Lesbian or bisexual IDU	29	(11.0)	14	(3.9)	7	(3.8)	3	(4.2)	59	(6.5)
Transfusion/Hemophilia	5	(1.9)	2	(0.6)	2	(1.1)	2	(2.8)	11	(1.2)
Heterosexual (3)	80	(30.3)	135	(37.5)	92	(50.5)	43	(59.7)	353	(39.1)
Perinatal	1	(0.4)	6	(1.7)	9	(4.9)	0	(0.0)	18	(2.0)
Risk not reported	20	(7.6)	23	(6.4)	21	(11.5)	10	(13.9)	79	(8.8)
Subtotal	264	(100)	360	(100)	182	(100)	72	(100)	902	(100)
Transgender	73	(100)	117	(100)	110	(100)	39	(100)	351	(100)
Total	9726	(100)	2013	(100)	2876	(100)	996	(100)	15964	(100)

- (1) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
- (2) Total includes persons with multiple or unknown race.
- (3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.