

HIV SEMI-ANNUAL SURVEILLANCE REPORT

San Francisco Department of Public Health HIV Cases Reported Through December 2013

Contents	Page
Surveillance Summary.....	1
Figure 1: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year	2
Figure 2: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year	2
Table 1. Diagnoses of HIV infection by age and year of initial HIV diagnosis.....	3
Table 2: Diagnoses of HIV infection stage 3 (AIDS) by age and year of AIDS diagnosis.....	4
Table 3: Diagnoses of HIV infection by gender and year of initial HIV diagnosis	5
Table 4: Diagnoses of HIV infection stage 3 (AIDS) by gender and year of AIDS diagnosis...	6
Table 5: Diagnoses of HIV infection by race/ethnicity and year of initial HIV diagnosis.....	7
Table 6: Diagnoses of HIV infection stage 3 (AIDS) by race/ethnicity and year of AIDS diagnosis.....	8
Table 7: Diagnoses of HIV infection by transmission category and year of initial HIV diagnosis.....	9
Table 8: Diagnoses of HIV infection stage 3 (AIDS) by transmission category and year of AIDS diagnosis.....	10
Table 9: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year	11
Table 10: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year	12
Table 11: Persons living with HIV infection by age, gender and race/ethnicity.....	13
Table 12: Persons living with HIV infection by transmission category, gender and race/ethnicity.....	14

The HIV Surveillance Report is published semi-annually by the
San Francisco Department of Public Health, Population Health Division
Applied Research, Community Health Epidemiology, and Surveillance Branch (ARCHES)
25 Van Ness Avenue, Suite 500, San Francisco, CA 94102; Phone (415) 437-6336, FAX (415) 431-0353

Director of Health: Barbara A. Garcia, MPA
Director of Population Health Division: Tomas Aragon, MD, DrPH
HIV Surveillance Section Directors: Susan Scheer, PhD, MPH; Ling Hsu, MPH

***The HIV Surveillance Report is accessible via internet:
<http://www.sfdph.org/dph/files/reports/default.asp>***

HIV SURVEILLANCE SUMMARY

San Francisco¹ (as of 12/31/2013)

Persons living with HIV infection ²	15,867
Diagnoses of HIV infection stage 3 (AIDS), cumulative	29,531
Deaths among persons diagnosed with HIV infection stage 3 (AIDS), cumulative	19,902

California³ (as of 12/31/2013)

Persons living with HIV infection ²	120,480
Diagnoses of HIV infection stage 3 (AIDS), cumulative	168,602
Deaths among persons diagnosed with HIV infection stage 3 (AIDS), cumulative	95,032

United States⁴ (as of 12/31/2010)

Persons living with HIV infection ²	898,529
Diagnoses of HIV infection stage 3 (AIDS), cumulative (as of 12/31/2011)	1,172,489
Deaths among persons diagnosed with HIV infection stage 3 (AIDS), cumulative	642,603


¹ Residents of San Francisco at time of HIV or AIDS diagnosis.

² Persons diagnosed with HIV infection regardless of disease stage.

³ CA data source: www.cdph.ca.gov/data/statistics/Pages/OAHIVAIDSStatistics.aspx.


⁴ US data source: CDC HIV Surveillance Report 2011, vol. 23 (Table 2b, 12b, 15b).
www.cdc.gov/hiv/topics/surveillance/resources/reports/.

Figure 1: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year, San Francisco, 2002-2013*



* Reporting for recent years is incomplete. See Table 9 for actual numbers per year.

Figure 2: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year, San Francisco, 1980-2013 †



† Reporting for recent years is incomplete. See Table 10 for actual numbers per year.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2013

Table 1. Diagnoses of HIV infection by age and year of initial HIV diagnosis (1), San Francisco, 2002-2013

Age at initial HIV diagnosis (years)	Year of initial HIV diagnosis					
	2002 No. (%)	2003 No. (%)	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)
0 - 12	1 (0.1)	1 (0.2)	2 (0.3)	1 (0.2)	0 (0.0)	0 (0.0)
13 - 17	3 (0.4)	4 (0.6)	1 (0.1)	1 (0.2)	1 (0.2)	2 (0.4)
18 - 24	57 (8.4)	48 (7.6)	63 (9.1)	48 (8.8)	60 (11.7)	57 (10.7)
25 - 29	89 (13.1)	63 (10.0)	90 (13.0)	71 (13.0)	64 (12.4)	98 (18.4)
30 - 39	302 (44.5)	281 (44.5)	273 (39.3)	205 (37.6)	175 (34.0)	190 (35.7)
40 - 49	158 (23.3)	176 (27.8)	186 (26.8)	158 (29.0)	143 (27.8)	128 (24.1)
50 - 59	60 (8.8)	51 (8.1)	62 (8.9)	49 (9.0)	56 (10.9)	44 (8.3)
60 +	9 (1.3)	8 (1.3)	17 (2.4)	12 (2.2)	16 (3.1)	13 (2.4)
Total	679 (100)	632 (100)	694 (100)	545 (100)	515 (100)	532 (100)

Age at initial HIV diagnosis (years)	Year of initial HIV diagnosis					
	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	4 (0.8)	0 (0.0)	2 (0.5)	1 (0.2)	0 (0.0)	0 (0.0)
18 - 24	48 (9.4)	56 (12.1)	53 (12.2)	42 (10.3)	53 (12.4)	43 (13.0)
25 - 29	82 (16.0)	59 (12.7)	55 (12.7)	61 (14.9)	70 (16.4)	73 (22.0)
30 - 39	189 (36.8)	144 (31.1)	134 (30.9)	109 (26.7)	133 (31.1)	93 (28.0)
40 - 49	143 (27.9)	127 (27.4)	124 (28.6)	127 (31.1)	122 (28.5)	82 (24.7)
50 - 59	40 (7.8)	60 (13.0)	50 (11.5)	54 (13.2)	39 (9.1)	29 (8.7)
60 +	7 (1.4)	17 (3.7)	16 (3.7)	15 (3.7)	11 (2.6)	12 (3.6)
Total	513 (100)	463 (100)	434 (100)	409 (100)	428 (100)	332 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of HIV antibody test, viral load or CD4 test, initiation of antiretroviral therapy, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report

Cases Reported as of 12/31/2013

Table 2. Diagnoses of HIV infection stage 3 (AIDS) by age and year of AIDS diagnosis (1), San Francisco, 2002-2013

Age at AIDS diagnosis (years)	Year of AIDS diagnosis					
	2002 No. (%)	2003 No. (%)	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)
0 - 12	1 (0.2)	1 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	1 (0.2)	1 (0.2)	1 (0.2)	0 (0.0)	0 (0.0)	1 (0.2)
18 - 24	7 (1.4)	20 (3.6)	11 (2.3)	14 (2.9)	13 (2.9)	15 (3.3)
25 - 29	33 (6.7)	34 (6.1)	21 (4.3)	32 (6.7)	25 (5.6)	39 (8.7)
30 - 39	180 (36.4)	210 (37.4)	170 (35.1)	157 (32.8)	123 (27.6)	134 (29.8)
40 - 49	181 (36.6)	197 (35.1)	166 (34.2)	183 (38.2)	183 (41.0)	168 (37.3)
50 - 59	81 (16.4)	83 (14.8)	87 (17.9)	79 (16.5)	81 (18.2)	67 (14.9)
60 +	10 (2.0)	15 (2.7)	29 (6.0)	14 (2.9)	21 (4.7)	26 (5.8)
Total	494 (100)	561 (100)	485 (100)	479 (100)	446 (100)	450 (100)

Age at AIDS diagnosis (years)	Year of AIDS diagnosis					
	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	1 (0.2)	1 (0.3)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
18 - 24	17 (3.9)	15 (4.6)	14 (4.8)	12 (4.8)	9 (3.8)	7 (4.6)
25 - 29	31 (7.2)	23 (7.1)	26 (8.9)	19 (7.6)	24 (10.1)	16 (10.5)
30 - 39	123 (28.4)	74 (22.9)	72 (24.6)	62 (24.8)	61 (25.6)	38 (24.8)
40 - 49	163 (37.6)	120 (37.2)	97 (33.1)	98 (39.2)	79 (33.2)	48 (31.4)
50 - 59	76 (17.6)	69 (21.4)	62 (21.2)	50 (20.0)	46 (19.3)	31 (20.3)
60 +	22 (5.1)	21 (6.5)	22 (7.5)	9 (3.6)	19 (8.0)	13 (8.5)
Total	433 (100)	323 (100)	293 (100)	250 (100)	238 (100)	153 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2013

Table 3. Diagnoses of HIV infection by gender and year of initial HIV diagnosis (1), San Francisco, 2002-2013

Gender	Year of initial HIV diagnosis					
	2002 No. (%)	2003 No. (%)	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)
Male	606 (89.2)	563 (89.1)	626 (90.2)	487 (89.4)	467 (90.7)	466 (87.6)
Female	48 (7.1)	51 (8.1)	53 (7.6)	44 (8.1)	35 (6.8)	43 (8.1)
Transgender	25 (3.7)	18 (2.8)	15 (2.2)	14 (2.6)	13 (2.5)	23 (4.3)
Total	679 (100)	632 (100)	694 (100)	545 (100)	515 (100)	532 (100)

Gender	Year of initial HIV diagnosis					
	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)
Male	458 (89.3)	421 (90.9)	389 (89.6)	357 (87.3)	404 (94.4)	300 (90.4)
Female	38 (7.4)	24 (5.2)	34 (7.8)	43 (10.5)	22 (5.1)	23 (6.9)
Transgender	17 (3.3)	18 (3.9)	11 (2.5)	9 (2.2)	2 (0.5)	9 (2.7)
Total	513 (100)	463 (100)	434 (100)	409 (100)	428 (100)	332 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of HIV antibody test, viral load or CD4 test, initiation of antiretroviral therapy, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2013

Table 4. Diagnoses of HIV infection stage 3 (AIDS) by gender and year of AIDS diagnosis (1), San Francisco, 2002-2013

Gender	Year of AIDS diagnosis					
	2002 No. (%)	2003 No. (%)	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)
Male	439 (88.9)	496 (88.4)	427 (88.0)	428 (89.4)	401 (89.9)	406 (90.2)
Female	35 (7.1)	40 (7.1)	41 (8.5)	37 (7.7)	30 (6.7)	33 (7.3)
Transgender	20 (4.0)	25 (4.5)	17 (3.5)	14 (2.9)	15 (3.4)	11 (2.4)
Total	494 (100)	561 (100)	485 (100)	479 (100)	446 (100)	450 (100)

Gender	Year of AIDS diagnosis					
	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)
Male	376 (86.8)	288 (89.2)	263 (89.8)	214 (85.6)	209 (87.8)	133 (86.9)
Female	42 (9.7)	24 (7.4)	16 (5.5)	24 (9.6)	25 (10.5)	15 (9.8)
Transgender	15 (3.5)	11 (3.4)	14 (4.8)	12 (4.8)	4 (1.7)	5 (3.3)
Total	433 (100)	323 (100)	293 (100)	250 (100)	238 (100)	153 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2013

Table 5. Diagnoses of HIV infection by race/ethnicity and year of initial HIV diagnosis (1), San Francisco, 2002-2013

Race/ethnicity	Year of initial HIV diagnosis					
	2002 No. (%)	2003 No. (%)	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)
White	383 (56.4)	349 (55.2)	389 (56.1)	296 (54.3)	281 (54.6)	274 (51.5)
African American	106 (15.6)	92 (14.6)	91 (13.1)	78 (14.3)	75 (14.6)	78 (14.7)
Latino	123 (18.1)	127 (20.1)	132 (19.0)	116 (21.3)	109 (21.2)	107 (20.1)
Asian/Pacific Islander	38 (5.6)	42 (6.6)	51 (7.3)	30 (5.5)	32 (6.2)	48 (9.0)
Native American	7 (1.0)	6 (0.9)	5 (0.7)	7 (1.3)	3 (0.6)	1 (0.2)
Total (2)	679 (100)	632 (100)	694 (100)	545 (100)	515 (100)	532 (100)

Race/ethnicity	Year of initial HIV diagnosis					
	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)
White	256 (49.9)	239 (51.6)	208 (47.9)	212 (51.8)	207 (48.4)	156 (47.0)
African American	80 (15.6)	69 (14.9)	60 (13.8)	64 (15.6)	45 (10.5)	37 (11.1)
Latino	117 (22.8)	95 (20.5)	107 (24.7)	83 (20.3)	107 (25.0)	86 (25.9)
Asian/Pacific Islander	40 (7.8)	38 (8.2)	40 (9.2)	32 (7.8)	51 (11.9)	41 (12.3)
Native American	4 (0.8)	1 (0.2)	3 (0.7)	2 (0.5)	7 (1.6)	2 (0.6)
Total (2)	513 (100)	463 (100)	434 (100)	409 (100)	428 (100)	332 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of HIV antibody test, viral load or CD4 test, initiation of antiretroviral therapy, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

(2) Total includes persons with multiple or unknown race.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2013

Table 6. Diagnoses of HIV infection stage 3 (AIDS) by race/ethnicity and year of AIDS diagnosis (1), San Francisco, 2002-2013

Race/ethnicity	Year of AIDS diagnosis					
	2002 No. (%)	2003 No. (%)	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)
White	286 (57.9)	279 (49.7)	258 (53.2)	270 (56.4)	250 (56.1)	261 (58.0)
African American	87 (17.6)	105 (18.7)	76 (15.7)	82 (17.1)	79 (17.7)	76 (16.9)
Latino	77 (15.6)	128 (22.8)	115 (23.7)	91 (19.0)	78 (17.5)	75 (16.7)
Asian/Pacific Islander	28 (5.7)	33 (5.9)	22 (4.5)	25 (5.2)	23 (5.2)	27 (6.0)
Native American	2 (0.4)	4 (0.7)	4 (0.8)	3 (0.6)	4 (0.9)	0 (0.0)
Total (2)	494 (100)	561 (100)	485 (100)	479 (100)	446 (100)	450 (100)

Race/ethnicity	Year of AIDS diagnosis					
	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)
White	225 (52.0)	170 (52.6)	144 (49.1)	131 (52.4)	119 (50.0)	73 (47.7)
African American	79 (18.2)	61 (18.9)	52 (17.7)	43 (17.2)	39 (16.4)	25 (16.3)
Latino	73 (16.9)	65 (20.1)	61 (20.8)	45 (18.0)	51 (21.4)	30 (19.6)
Asian/Pacific Islander	39 (9.0)	13 (4.0)	25 (8.5)	15 (6.0)	17 (7.1)	16 (10.5)
Native American	2 (0.5)	3 (0.9)	2 (0.7)	2 (0.8)	1 (0.4)	1 (0.7)
Total (2)	433 (100)	323 (100)	293 (100)	250 (100)	238 (100)	153 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

(2) Total includes persons with multiple or unknown race.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2013

Table 7. Diagnoses of HIV infection by transmission category and year of initial HIV diagnosis (1), San Francisco, 2002-2013

Transmission category (2)	Year of initial HIV diagnosis					
	2002 No. (%)	2003 No. (%)	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)
Gay or bisexual male	459 (67.6)	429 (67.9)	475 (68.4)	376 (69.0)	366 (71.1)	350 (65.8)
Injection drug user (IDU)	63 (9.3)	63 (10.0)	59 (8.5)	31 (5.7)	36 (7.0)	32 (6.0)
Gay or bisexual male IDU	116 (17.1)	96 (15.2)	114 (16.4)	86 (15.8)	75 (14.6)	88 (16.5)
Lesbian or bisexual IDU	3 (0.4)	1 (0.2)	4 (0.6)	5 (0.9)	2 (0.4)	3 (0.6)
Transfusion/Hemophilia	0 (0.0)	1 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	19 (2.8)	28 (4.4)	21 (3.0)	31 (5.7)	24 (4.7)	42 (7.9)
Perinatal	1 (0.1)	1 (0.2)	2 (0.3)	1 (0.2)	0 (0.0)	0 (0.0)
Risk not reported	18 (2.7)	13 (2.1)	19 (2.7)	15 (2.8)	12 (2.3)	17 (3.2)
Total	679 (100)	632 (100)	694 (100)	545 (100)	515 (100)	532 (100)

Transmission category (2)	Year of initial HIV diagnosis					
	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)
Gay or bisexual male	374 (72.9)	333 (71.9)	282 (65.0)	289 (70.7)	335 (78.3)	255 (76.8)
Injection drug user (IDU)	26 (5.1)	20 (4.3)	33 (7.6)	25 (6.1)	12 (2.8)	16 (4.8)
Gay or bisexual male IDU	60 (11.7)	70 (15.1)	61 (14.1)	46 (11.2)	40 (9.3)	31 (9.3)
Lesbian or bisexual IDU	4 (0.8)	4 (0.9)	1 (0.2)	3 (0.7)	2 (0.5)	2 (0.6)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	32 (6.2)	21 (4.5)	36 (8.3)	27 (6.6)	23 (5.4)	13 (3.9)
Perinatal	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Risk not reported	17 (3.3)	15 (3.2)	21 (4.8)	19 (4.6)	16 (3.7)	15 (4.5)
Total	513 (100)	463 (100)	434 (100)	409 (100)	428 (100)	332 (100)

- (1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of HIV antibody test, viral load or CD4 test, initiation of antiretroviral therapy, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.
- (2) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
- (3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2013

Table 8. Diagnoses of HIV infection stage 3 (AIDS) by transmission category and year of AIDS diagnosis (1), San Francisco, 2002-2013

Transmission category (2)	Year of AIDS diagnosis					
	2002 No. (%)	2003 No. (%)	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)
Gay or bisexual male	302 (61.1)	354 (63.1)	315 (64.9)	298 (62.2)	287 (64.3)	290 (64.4)
Injection drug user (IDU)	57 (11.5)	71 (12.7)	50 (10.3)	49 (10.2)	37 (8.3)	38 (8.4)
Gay or bisexual male IDU	109 (22.1)	105 (18.7)	93 (19.2)	99 (20.7)	92 (20.6)	85 (18.9)
Lesbian or bisexual IDU	4 (0.8)	3 (0.5)	1 (0.2)	3 (0.6)	2 (0.4)	0 (0.0)
Transfusion/Hemophilia	1 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	14 (2.8)	21 (3.7)	16 (3.3)	19 (4.0)	20 (4.5)	29 (6.4)
Perinatal	1 (0.2)	1 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Risk not reported	6 (1.2)	6 (1.1)	10 (2.1)	11 (2.3)	8 (1.8)	8 (1.8)
Total	494 (100)	561 (100)	485 (100)	479 (100)	446 (100)	450 (100)

Transmission category (2)	Year of AIDS diagnosis					
	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)
Gay or bisexual male	264 (61.0)	210 (65.0)	168 (57.3)	147 (58.8)	154 (64.7)	89 (58.2)
Injection drug user (IDU)	30 (6.9)	33 (10.2)	24 (8.2)	32 (12.8)	20 (8.4)	15 (9.8)
Gay or bisexual male IDU	93 (21.5)	52 (16.1)	77 (26.3)	46 (18.4)	35 (14.7)	38 (24.8)
Lesbian or bisexual IDU	4 (0.9)	2 (0.6)	0 (0.0)	1 (0.4)	2 (0.8)	0 (0.0)
Transfusion/Hemophilia	0 (0.0)	1 (0.3)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	31 (7.2)	17 (5.3)	16 (5.5)	16 (6.4)	18 (7.6)	7 (4.6)
Perinatal	0 (0.0)	1 (0.3)	0 (0.0)	1 (0.4)	0 (0.0)	0 (0.0)
Risk not reported	11 (2.5)	7 (2.2)	8 (2.7)	7 (2.8)	9 (3.8)	4 (2.6)
Total	433 (100)	323 (100)	293 (100)	250 (100)	238 (100)	153 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

(2) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.

(3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2013

Table 9. Diagnoses of HIV infection, HIV deaths and persons living with HIV by year, San Francisco, 2002-2013

Year	Number of cases diagnosed per year (1)	Number of deaths occurred per year (1)	Number of persons living with HIV infection (1)
2002 (2)	679	347	13411
2003	632	338	13705
2004	694	343	14056
2005	545	348	14253
2006	515	327	14441
2007	532	323	14650
2008	513	262	14901
2009	463	251	15113
2010	434	246	15301
2011	409	231	15479
2012	428	208	15699
2013	332	164	15867

(1) Data in recent years are incomplete due to delays in case/death reporting.

(2) Year of the mandatory reporting of HIV non-AIDS cases in California began.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2013

Table 10. Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year, San Francisco, 1980-2013

Year	Number of cases diagnosed per year (1)	Number of deaths occurred per year (1)	Number of persons living with AIDS (1)
1980	3	0	3
1981	26	9	20
1982	99	32	87
1983	274	111	250
1984	558	273	535
1985	860	534	861
1986	1236	807	1290
1987	1630	878	2042
1988	1763	1039	2766
1989	2162	1278	3650
1990	2046	1365	4331
1991	2288	1510	5109
1992	2331	1641	5799
1993	2071	1600	6270
1994	1787	1594	6463
1995	1561	1484	6540
1996	1079	992	6627
1997	808	424	7011
1998	696	402	7305
1999	577	353	7529
2000	557	350	7736
2001	514	324	7926
2002	494	320	8100
2003	561	303	8358
2004	485	310	8533
2005	479	312	8700
2006	446	289	8857
2007	450	270	9037
2008	433	229	9241
2009	323	210	9354
2010	293	194	9453
2011	250	186	9517
2012	238	163	9592
2013	153	116	9629
Total	29531	19902	

(1) Data in recent years are incomplete due to delays in case/death reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2013

Table 11. Persons living with HIV infection by age, gender and race/ethnicity, San Francisco

Current age (years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander/ Native American No. (%)	Total (1) No. (%)
Male					
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.0)
13 - 17	0 (0.0)	0 (0.0)	1 (0.0)	0 (0.0)	1 (0.0)
18 - 24	28 (0.3)	25 (1.6)	38 (1.5)	16 (1.9)	111 (0.8)
25 - 29	159 (1.7)	48 (3.1)	130 (5.2)	45 (5.2)	394 (2.7)
30 - 39	798 (8.5)	133 (8.6)	472 (18.9)	183 (21.3)	1648 (11.3)
40 - 49	2658 (28.2)	427 (27.6)	877 (35.0)	321 (37.3)	4389 (30.0)
50 - 59	3588 (38.1)	586 (37.9)	728 (29.1)	208 (24.2)	5184 (35.5)
60 +	2187 (23.2)	326 (21.1)	257 (10.3)	87 (10.1)	2879 (19.7)
Subtotal	9418 (100)	1545 (100)	2503 (100)	860 (100)	14607 (100)
Female					
0 - 12	0 (0.0)	1 (0.3)	1 (0.6)	0 (0.0)	2 (0.2)
13 - 17	0 (0.0)	0 (0.0)	2 (1.1)	0 (0.0)	3 (0.3)
18 - 24	2 (0.7)	5 (1.4)	9 (5.0)	1 (1.4)	18 (2.0)
25 - 29	10 (3.7)	6 (1.6)	12 (6.7)	1 (1.4)	29 (3.2)
30 - 39	32 (11.9)	42 (11.5)	32 (17.9)	16 (21.9)	131 (14.4)
40 - 49	89 (33.2)	95 (26.0)	45 (25.1)	25 (34.2)	259 (28.5)
50 - 59	97 (36.2)	149 (40.8)	46 (25.7)	22 (30.1)	321 (35.4)
60 +	38 (14.2)	67 (18.4)	32 (17.9)	8 (11.0)	145 (16.0)
Subtotal	268 (100)	365 (100)	179 (100)	73 (100)	908 (100)
Transgender					
13 - 39	12 (16.9)	39 (31.7)	43 (40.2)	8 (21.1)	110 (31.3)
40 +	59 (83.1)	84 (68.3)	64 (59.8)	30 (78.9)	242 (68.8)
Subtotal	71 (100)	123 (100)	107 (100)	38 (100)	352 (100)
Total	9757 (100)	2033 (100)	2789 (100)	971 (100)	15867 (100)

(1) Total includes persons with multiple or unknown race.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2013

Table 12. Persons living with HIV infection by transmission category, gender and race/ethnicity, San Francisco

Transmission category (1)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander/ Native American No. (%)	Total (2) No. (%)
Male					
Gay or bisexual male	7660 (81.3)	874 (56.6)	2040 (81.5)	716 (83.3)	11488 (78.6)
Injection drug user (IDU)	193 (2.0)	239 (15.5)	61 (2.4)	21 (2.4)	523 (3.6)
Gay or bisexual male IDU	1452 (15.4)	317 (20.5)	298 (11.9)	92 (10.7)	2216 (15.2)
Transfusion/Hemophilia	6 (0.1)	3 (0.2)	2 (0.1)	3 (0.3)	14 (0.1)
Heterosexual (3)	30 (0.3)	69 (4.5)	49 (2.0)	11 (1.3)	163 (1.1)
Perinatal	1 (0.0)	3 (0.2)	2 (0.1)	2 (0.2)	10 (0.1)
Risk not reported	76 (0.8)	40 (2.6)	51 (2.0)	15 (1.7)	193 (1.3)
Subtotal	9418 (100)	1545 (100)	2503 (100)	860 (100)	14607 (100)
Female					
Injection drug user (IDU)	133 (49.6)	190 (52.1)	51 (28.5)	13 (17.8)	394 (43.4)
Lesbian or bisexual IDU	27 (10.1)	15 (4.1)	7 (3.9)	3 (4.1)	58 (6.4)
Transfusion/Hemophilia	5 (1.9)	2 (0.5)	2 (1.1)	2 (2.7)	11 (1.2)
Heterosexual (3)	78 (29.1)	130 (35.6)	91 (50.8)	45 (61.6)	347 (38.2)
Perinatal	1 (0.4)	6 (1.6)	9 (5.0)	0 (0.0)	18 (2.0)
Risk not reported	24 (9.0)	22 (6.0)	19 (10.6)	10 (13.7)	80 (8.8)
Subtotal	268 (100)	365 (100)	179 (100)	73 (100)	908 (100)
Transgender	71 (100)	123 (100)	107 (100)	38 (100)	352 (100)
Total	9757 (100)	2033 (100)	2789 (100)	971 (100)	15867 (100)

(1) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.

(2) Total includes persons with multiple or unknown race.

(3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.