

HIV SEMI-ANNUAL SURVEILLANCE REPORT

San Francisco Department of Public Health

HIV Cases Reported Through June 2014

Contents	Page
Surveillance Summary.....	1
Figure 1: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year	2
Figure 2: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year	2
Table 1. Diagnoses of HIV infection by age and year of initial HIV diagnosis.....	3
Table 2: Diagnoses of HIV infection stage 3 (AIDS) by age and year of AIDS diagnosis.....	4
Table 3: Diagnoses of HIV infection by gender and year of initial HIV diagnosis	5
Table 4: Diagnoses of HIV infection stage 3 (AIDS) by gender and year of AIDS diagnosis...	6
Table 5: Diagnoses of HIV infection by race/ethnicity and year of initial HIV diagnosis.....	7
Table 6: Diagnoses of HIV infection stage 3 (AIDS) by race/ethnicity and year of AIDS diagnosis.....	8
Table 7: Diagnoses of HIV infection by transmission category and year of initial HIV diagnosis.....	9
Table 8: Diagnoses of HIV infection stage 3 (AIDS) by transmission category and year of AIDS diagnosis.....	10
Table 9: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year	11
Table 10: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year	12
Table 11: Persons living with HIV infection by age, gender and race/ethnicity.....	13
Table 12: Persons living with HIV infection by transmission category, gender and race/ethnicity.....	14

The HIV Surveillance Report is published semi-annually by the
San Francisco Department of Public Health, Population Health Division
Applied Research, Community Health Epidemiology, and Surveillance Branch (ARCHES)
25 Van Ness Avenue, Suite 500, San Francisco, CA 94102; Phone (415) 437-6336, FAX (415) 431-0353

Director of Health: Barbara A. Garcia, MPA
Director of Population Health Division: Tomas Aragon, MD, DrPH
HIV Surveillance Section Directors: Susan Scheer, PhD, MPH; Ling Hsu, MPH

***The HIV Surveillance Report is accessible via internet:
<http://www.sfdph.org/dph/files/reports/default.asp>***

HIV SURVEILLANCE SUMMARY

San Francisco¹ (as of 06/30/2014)

Persons living with HIV infection ²	15,971
Diagnoses of HIV infection stage 3 (AIDS), cumulative	29,605
Deaths among persons diagnosed with HIV infection stage 3 (AIDS), cumulative	19,978

California³ (as of 06/30/2014)

Persons living with HIV infection ²	122,382
Diagnoses of HIV infection stage 3 (AIDS), cumulative	169,588
Deaths among persons diagnosed with HIV infection stage 3 (AIDS), cumulative	95,529

United States⁴ (as of 12/31/2010)

Persons living with HIV infection ²	898,529
Diagnoses of HIV infection stage 3 (AIDS), cumulative (as of 12/31/2011)	1,172,489
Deaths among persons diagnosed with HIV infection stage 3 (AIDS), cumulative	642,603


¹ Residents of San Francisco at time of HIV or AIDS diagnosis.

² Persons diagnosed with HIV infection regardless of disease stage.

³ CA data source: www.cdph.ca.gov/data/statistics/Pages/OAHIVAIDSStatistics.aspx.


⁴ US data source: CDC HIV Surveillance Report 2011, vol. 23 (Table 2b, 12b, 15b).
www.cdc.gov/hiv/topics/surveillance/resources/reports/.

Figure 1: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year, San Francisco, 2002-2014*



* Reporting for recent years is incomplete. See Table 9 for actual numbers per year.

Figure 2: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year, San Francisco, 1980-2014 †



† Reporting for recent years is incomplete. See Table 10 for actual numbers per year.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2014

Table 1. Diagnoses of HIV infection by age and year of initial HIV diagnosis (1), San Francisco, 2003-2014

Age at initial HIV diagnosis (years)	Year of initial HIV diagnosis					
	2003 No. (%)	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)
0 - 12	1 (0.2)	2 (0.3)	1 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	4 (0.6)	1 (0.1)	1 (0.2)	1 (0.2)	2 (0.4)	4 (0.8)
18 - 24	48 (7.5)	63 (8.9)	48 (8.8)	61 (11.7)	56 (10.5)	51 (9.8)
25 - 29	64 (10.0)	92 (13.0)	71 (13.0)	65 (12.5)	98 (18.5)	83 (16.0)
30 - 39	285 (44.5)	278 (39.4)	208 (38.0)	177 (33.9)	191 (36.0)	188 (36.3)
40 - 49	179 (28.0)	190 (27.0)	157 (28.7)	145 (27.8)	127 (23.9)	146 (28.2)
50 - 59	51 (8.0)	62 (8.8)	49 (9.0)	57 (10.9)	44 (8.3)	39 (7.5)
60 +	8 (1.3)	17 (2.4)	12 (2.2)	16 (3.1)	13 (2.4)	7 (1.4)
Total	640 (100)	705 (100)	547 (100)	522 (100)	531 (100)	518 (100)

Age at initial HIV diagnosis (years)	Year of initial HIV diagnosis					
	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	1 (0.2)	2 (0.5)	1 (0.2)	0 (0.0)	0 (0.0)	1 (0.7)
18 - 24	56 (12.1)	55 (12.4)	43 (10.5)	54 (12.7)	46 (12.7)	15 (11.0)
25 - 29	58 (12.6)	56 (12.6)	62 (15.2)	68 (16.0)	78 (21.5)	26 (19.1)
30 - 39	143 (31.0)	137 (30.9)	107 (26.2)	131 (30.8)	104 (28.7)	46 (33.8)
40 - 49	126 (27.3)	127 (28.6)	127 (31.1)	123 (28.9)	90 (24.9)	28 (20.6)
50 - 59	61 (13.2)	51 (11.5)	54 (13.2)	38 (8.9)	31 (8.6)	12 (8.8)
60 +	17 (3.7)	16 (3.6)	15 (3.7)	11 (2.6)	13 (3.6)	8 (5.9)
Total	462 (100)	444 (100)	409 (100)	425 (100)	362 (100)	136 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of HIV antibody test, viral load or CD4 test, initiation of antiretroviral therapy, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2014

Table 2. Diagnoses of HIV infection stage 3 (AIDS) by age and year of AIDS diagnosis (1), San Francisco, 2003-2014

Age at AIDS diagnosis (years)	Year of AIDS diagnosis					
	2003 No. (%)	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)
0 - 12	1 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	1 (0.2)	1 (0.2)	0 (0.0)	0 (0.0)	1 (0.2)	1 (0.2)
18 - 24	20 (3.6)	11 (2.3)	14 (2.9)	13 (2.9)	15 (3.3)	17 (3.9)
25 - 29	34 (6.1)	21 (4.3)	32 (6.7)	25 (5.6)	39 (8.7)	31 (7.2)
30 - 39	211 (37.6)	170 (35.1)	157 (32.7)	124 (27.7)	134 (29.8)	123 (28.5)
40 - 49	196 (34.9)	165 (34.1)	183 (38.1)	182 (40.6)	167 (37.2)	164 (38.0)
50 - 59	83 (14.8)	87 (18.0)	80 (16.7)	83 (18.5)	67 (14.9)	75 (17.4)
60 +	15 (2.7)	29 (6.0)	14 (2.9)	21 (4.7)	26 (5.8)	21 (4.9)
Total	561 (100)	484 (100)	480 (100)	448 (100)	449 (100)	432 (100)

Age at AIDS diagnosis (years)	Year of AIDS diagnosis					
	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	1 (0.3)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.8)
18 - 24	15 (4.6)	14 (4.7)	12 (4.8)	10 (4.2)	8 (4.6)	3 (5.4)
25 - 29	23 (7.1)	26 (8.8)	20 (8.0)	24 (10.0)	17 (9.8)	13 (23.2)
30 - 39	74 (22.9)	73 (24.7)	62 (24.7)	60 (25.1)	41 (23.6)	18 (32.1)
40 - 49	120 (37.2)	97 (32.9)	98 (39.0)	80 (33.5)	55 (31.6)	13 (23.2)
50 - 59	69 (21.4)	62 (21.0)	50 (19.9)	46 (19.2)	36 (20.7)	7 (12.5)
60 +	21 (6.5)	23 (7.8)	9 (3.6)	19 (7.9)	17 (9.8)	1 (1.8)
Total	323 (100)	295 (100)	251 (100)	239 (100)	174 (100)	56 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2014

Table 3. Diagnoses of HIV infection by gender and year of initial HIV diagnosis (1), San Francisco, 2003-2014

Gender	Year of initial HIV diagnosis					
	2003 No. (%)	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)
Male	571 (89.2)	637 (90.4)	489 (89.4)	472 (90.4)	465 (87.6)	460 (88.8)
Female	51 (8.0)	53 (7.5)	44 (8.0)	35 (6.7)	43 (8.1)	41 (7.9)
Transgender	18 (2.8)	15 (2.1)	14 (2.6)	15 (2.9)	23 (4.3)	17 (3.3)
Total	640 (100)	705 (100)	547 (100)	522 (100)	531 (100)	518 (100)

Gender	Year of initial HIV diagnosis					
	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)
Male	420 (90.9)	399 (89.9)	359 (87.8)	402 (94.6)	328 (90.6)	129 (94.9)
Female	25 (5.4)	34 (7.7)	41 (10.0)	21 (4.9)	24 (6.6)	6 (4.4)
Transgender	17 (3.7)	11 (2.5)	9 (2.2)	2 (0.5)	10 (2.8)	1 (0.7)
Total	462 (100)	444 (100)	409 (100)	425 (100)	362 (100)	136 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of HIV antibody test, viral load or CD4 test, initiation of antiretroviral therapy, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2014

Table 4. Diagnoses of HIV infection stage 3 (AIDS) by gender and year of AIDS diagnosis (1), San Francisco, 2003-2014

Gender	Year of AIDS diagnosis					
	2003 No. (%)	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)
Male	496 (88.4)	426 (88.0)	429 (89.4)	403 (90.0)	405 (90.2)	375 (86.8)
Female	40 (7.1)	41 (8.5)	37 (7.7)	30 (6.7)	33 (7.3)	43 (10.0)
Transgender	25 (4.5)	17 (3.5)	14 (2.9)	15 (3.3)	11 (2.4)	14 (3.2)
Total	561 (100)	484 (100)	480 (100)	448 (100)	449 (100)	432 (100)

Gender	Year of AIDS diagnosis					
	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)
Male	288 (89.2)	265 (89.8)	214 (85.3)	210 (87.9)	153 (87.9)	52 (92.9)
Female	24 (7.4)	16 (5.4)	24 (9.6)	25 (10.5)	16 (9.2)	3 (5.4)
Transgender	11 (3.4)	14 (4.7)	13 (5.2)	4 (1.7)	5 (2.9)	1 (1.8)
Total	323 (100)	295 (100)	251 (100)	239 (100)	174 (100)	56 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2014

Table 5. Diagnoses of HIV infection by race/ethnicity and year of initial HIV diagnosis (1), San Francisco, 2003-2014

Race/ethnicity	Year of initial HIV diagnosis					
	2003 No. (%)	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)
White	353 (55.2)	391 (55.5)	295 (53.9)	285 (54.6)	273 (51.4)	258 (49.8)
African American	92 (14.4)	92 (13.0)	78 (14.3)	76 (14.6)	77 (14.5)	82 (15.8)
Latino	130 (20.3)	135 (19.1)	118 (21.6)	109 (20.9)	107 (20.2)	118 (22.8)
Asian/Pacific Islander	43 (6.7)	54 (7.7)	30 (5.5)	33 (6.3)	49 (9.2)	40 (7.7)
Native American	6 (0.9)	5 (0.7)	7 (1.3)	3 (0.6)	1 (0.2)	4 (0.8)
Total (2)	640 (100)	705 (100)	547 (100)	522 (100)	531 (100)	518 (100)

Race/ethnicity	Year of initial HIV diagnosis					
	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)
White	241 (52.2)	213 (48.0)	211 (51.6)	209 (49.2)	163 (45.0)	61 (44.9)
African American	68 (14.7)	63 (14.2)	66 (16.1)	45 (10.6)	47 (13.0)	14 (10.3)
Latino	96 (20.8)	109 (24.5)	81 (19.8)	105 (24.7)	90 (24.9)	37 (27.2)
Asian/Pacific Islander	36 (7.8)	40 (9.0)	32 (7.8)	49 (11.5)	49 (13.5)	18 (13.2)
Native American	1 (0.2)	3 (0.7)	2 (0.5)	6 (1.4)	3 (0.8)	0 (0.0)
Total (2)	462 (100)	444 (100)	409 (100)	425 (100)	362 (100)	136 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of HIV antibody test, viral load or CD4 test, initiation of antiretroviral therapy, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

(2) Total includes persons with multiple or unknown race.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2014

Table 6. Diagnoses of HIV infection stage 3 (AIDS) by race/ethnicity and year of AIDS diagnosis (1), San Francisco, 2003-2014

Race/ethnicity	Year of AIDS diagnosis					
	2003 No. (%)	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)
White	279 (49.7)	258 (53.3)	270 (56.3)	252 (56.3)	261 (58.1)	224 (51.9)
African American	105 (18.7)	76 (15.7)	83 (17.3)	79 (17.6)	76 (16.9)	80 (18.5)
Latino	128 (22.8)	115 (23.8)	91 (19.0)	78 (17.4)	74 (16.5)	72 (16.7)
Asian/Pacific Islander	33 (5.9)	22 (4.5)	25 (5.2)	23 (5.1)	26 (5.8)	38 (8.8)
Native American	4 (0.7)	4 (0.8)	3 (0.6)	4 (0.9)	0 (0.0)	2 (0.5)
Total (2)	561 (100)	484 (100)	480 (100)	448 (100)	449 (100)	432 (100)

Race/ethnicity	Year of AIDS diagnosis					
	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)
White	170 (52.6)	145 (49.2)	131 (52.2)	119 (49.8)	81 (46.6)	18 (32.1)
African American	62 (19.2)	52 (17.6)	44 (17.5)	40 (16.7)	30 (17.2)	10 (17.9)
Latino	65 (20.1)	61 (20.7)	45 (17.9)	51 (21.3)	34 (19.5)	17 (30.4)
Asian/Pacific Islander	13 (4.0)	26 (8.8)	15 (6.0)	17 (7.1)	18 (10.3)	8 (14.3)
Native American	3 (0.9)	2 (0.7)	2 (0.8)	1 (0.4)	1 (0.6)	0 (0.0)
Total (2)	323 (100)	295 (100)	251 (100)	239 (100)	174 (100)	56 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

(2) Total includes persons with multiple or unknown race.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2014

Table 7. Diagnoses of HIV infection by transmission category and year of initial HIV diagnosis (1), San Francisco, 2003-2014

Transmission category (2)	Year of initial HIV diagnosis					
	2003 No. (%)	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)
Gay or bisexual male	436 (68.1)	485 (68.8)	377 (68.9)	369 (70.7)	351 (66.1)	374 (72.2)
Injection drug user (IDU)	63 (9.8)	59 (8.4)	30 (5.5)	36 (6.9)	30 (5.6)	27 (5.2)
Gay or bisexual male IDU	96 (15.0)	115 (16.3)	89 (16.3)	80 (15.3)	88 (16.6)	63 (12.2)
Lesbian or bisexual IDU	1 (0.2)	4 (0.6)	5 (0.9)	2 (0.4)	3 (0.6)	5 (1.0)
Transfusion/Hemophilia	1 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	28 (4.4)	21 (3.0)	31 (5.7)	24 (4.6)	42 (7.9)	33 (6.4)
Perinatal	1 (0.2)	2 (0.3)	1 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)
Risk not reported	14 (2.2)	19 (2.7)	14 (2.6)	11 (2.1)	17 (3.2)	16 (3.1)
Total	640 (100)	705 (100)	547 (100)	522 (100)	531 (100)	518 (100)

Transmission category (2)	Year of initial HIV diagnosis					
	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)
Gay or bisexual male	330 (71.4)	289 (65.1)	290 (70.9)	334 (78.6)	277 (76.5)	106 (77.9)
Injection drug user (IDU)	21 (4.5)	33 (7.4)	25 (6.1)	12 (2.8)	20 (5.5)	6 (4.4)
Gay or bisexual male IDU	70 (15.2)	62 (14.0)	47 (11.5)	40 (9.4)	36 (9.9)	13 (9.6)
Lesbian or bisexual IDU	4 (0.9)	1 (0.2)	3 (0.7)	2 (0.5)	1 (0.3)	2 (1.5)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	23 (5.0)	37 (8.3)	26 (6.4)	22 (5.2)	15 (4.1)	2 (1.5)
Perinatal	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Risk not reported	14 (3.0)	22 (5.0)	18 (4.4)	15 (3.5)	13 (3.6)	7 (5.1)
Total	462 (100)	444 (100)	409 (100)	425 (100)	362 (100)	136 (100)

- (1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of HIV antibody test, viral load or CD4 test, initiation of antiretroviral therapy, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.
- (2) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
- (3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2014

Table 8. Diagnoses of HIV infection stage 3 (AIDS) by transmission category and year of AIDS diagnosis (1), San Francisco, 2003-2014

Transmission category (2)	Year of AIDS diagnosis					
	2003 No. (%)	2004 No. (%)	2005 No. (%)	2006 No. (%)	2007 No. (%)	2008 No. (%)
Gay or bisexual male	355 (63.3)	314 (64.9)	297 (61.9)	288 (64.3)	290 (64.6)	262 (60.6)
Injection drug user (IDU)	71 (12.7)	50 (10.3)	49 (10.2)	37 (8.3)	38 (8.5)	30 (6.9)
Gay or bisexual male IDU	104 (18.5)	93 (19.2)	101 (21.0)	93 (20.8)	85 (18.9)	93 (21.5)
Lesbian or bisexual IDU	3 (0.5)	1 (0.2)	3 (0.6)	2 (0.4)	0 (0.0)	4 (0.9)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	21 (3.7)	16 (3.3)	19 (4.0)	20 (4.5)	29 (6.5)	32 (7.4)
Perinatal	1 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Risk not reported	6 (1.1)	10 (2.1)	11 (2.3)	8 (1.8)	7 (1.6)	11 (2.5)
Total	561 (100)	484 (100)	480 (100)	448 (100)	449 (100)	432 (100)

Transmission category (2)	Year of AIDS diagnosis					
	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)
Gay or bisexual male	209 (64.7)	169 (57.3)	149 (59.4)	155 (64.9)	103 (59.2)	38 (67.9)
Injection drug user (IDU)	33 (10.2)	24 (8.1)	32 (12.7)	20 (8.4)	18 (10.3)	1 (1.8)
Gay or bisexual male IDU	53 (16.4)	78 (26.4)	46 (18.3)	35 (14.6)	41 (23.6)	10 (17.9)
Lesbian or bisexual IDU	2 (0.6)	0 (0.0)	1 (0.4)	2 (0.8)	0 (0.0)	1 (1.8)
Transfusion/Hemophilia	1 (0.3)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	17 (5.3)	16 (5.4)	16 (6.4)	18 (7.5)	9 (5.2)	4 (7.1)
Perinatal	1 (0.3)	0 (0.0)	1 (0.4)	0 (0.0)	0 (0.0)	0 (0.0)
Risk not reported	7 (2.2)	8 (2.7)	6 (2.4)	9 (3.8)	3 (1.7)	2 (3.6)
Total	323 (100)	295 (100)	251 (100)	239 (100)	174 (100)	56 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

(2) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.

(3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2014

Table 9. Diagnoses of HIV infection, HIV deaths and persons living with HIV by year, San Francisco, 2002-2014

Year	Number of cases diagnosed per year (1)	Number of deaths occurred per year (1)	Number of persons living with HIV infection (1)
2002 (2)	677	347	13411
2003	640	339	13712
2004	705	343	14074
2005	547	347	14274
2006	522	327	14469
2007	531	323	14677
2008	518	263	14932
2009	462	253	15141
2010	444	245	15340
2011	409	231	15518
2012	425	207	15736
2013	362	189	15909
2014	136	74	15971

(1) Data in recent years are incomplete due to delays in case/death reporting.

(2) Year of the mandatory reporting of HIV non-AIDS cases in California began.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2014

Table 10. Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year, San Francisco, 1980-2014

Year	Number of cases diagnosed per year (1)	Number of deaths occurred per year (1)	Number of persons living with AIDS (1)
1980	3	0	3
1981	26	9	20
1982	99	32	87
1983	274	111	250
1984	558	273	535
1985	860	534	861
1986	1236	807	1290
1987	1630	878	2042
1988	1763	1039	2766
1989	2162	1278	3650
1990	2046	1365	4331
1991	2287	1510	5108
1992	2331	1641	5798
1993	2070	1600	6268
1994	1785	1594	6459
1995	1559	1484	6534
1996	1079	992	6621
1997	806	424	7003
1998	695	402	7296
1999	578	353	7521
2000	558	350	7729
2001	515	323	7921
2002	493	320	8094
2003	561	303	8352
2004	484	310	8526
2005	480	311	8695
2006	448	289	8854
2007	449	270	9033
2008	432	229	9236
2009	323	212	9347
2010	295	193	9449
2011	251	186	9514
2012	239	162	9591
2013	174	135	9630
2014	56	59	9627
Total	29605	19978	

(1) Data in recent years are incomplete due to delays in case/death reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2014

Table 11. Persons living with HIV infection by age, gender and race/ethnicity, San Francisco

Current age (years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander/ Native American No. (%)	Total (1) No. (%)
Male					
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.0)
13 - 17	0 (0.0)	0 (0.0)	2 (0.1)	0 (0.0)	2 (0.0)
18 - 24	20 (0.2)	26 (1.7)	42 (1.6)	17 (1.9)	110 (0.7)
25 - 29	165 (1.7)	55 (3.6)	125 (4.9)	45 (5.1)	402 (2.7)
30 - 39	788 (8.3)	133 (8.6)	477 (18.7)	186 (21.1)	1646 (11.2)
40 - 49	2498 (26.5)	407 (26.3)	857 (33.6)	326 (37.0)	4197 (28.5)
50 - 59	3667 (38.8)	580 (37.5)	770 (30.2)	217 (24.6)	5317 (36.1)
60 +	2305 (24.4)	345 (22.3)	280 (11.0)	90 (10.2)	3043 (20.7)
Subtotal	9443 (100)	1546 (100)	2553 (100)	881 (100)	14718 (100)
Female					
0 - 12	0 (0.0)	1 (0.3)	1 (0.6)	0 (0.0)	2 (0.2)
13 - 17	0 (0.0)	0 (0.0)	2 (1.1)	0 (0.0)	2 (0.2)
18 - 24	3 (1.1)	4 (1.1)	9 (5.0)	1 (1.4)	19 (2.1)
25 - 29	8 (3.0)	5 (1.4)	10 (5.6)	1 (1.4)	24 (2.7)
30 - 39	33 (12.5)	41 (11.3)	33 (18.4)	17 (23.3)	133 (14.7)
40 - 49	84 (31.8)	91 (25.0)	43 (24.0)	23 (31.5)	246 (27.2)
50 - 59	97 (36.7)	146 (40.1)	47 (26.3)	23 (31.5)	320 (35.4)
60 +	39 (14.8)	76 (20.9)	34 (19.0)	8 (11.0)	157 (17.4)
Subtotal	264 (100)	364 (100)	179 (100)	73 (100)	903 (100)
Transgender					
13 - 39	12 (17.1)	38 (31.7)	40 (37.7)	10 (25.0)	109 (31.1)
40 +	58 (82.9)	82 (68.3)	66 (62.3)	30 (75.0)	241 (68.9)
Subtotal	70 (100)	120 (100)	106 (100)	40 (100)	350 (100)
Total	9777 (100)	2030 (100)	2838 (100)	994 (100)	15971 (100)

(1) Total includes persons with multiple or unknown race.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2014

Table 12. Persons living with HIV infection by transmission category, gender and race/ethnicity, San Francisco

Transmission category (1)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander/ Native American No. (%)	Total (2) No. (%)
Male					
Gay or bisexual male	7681 (81.3)	882 (57.1)	2085 (81.7)	738 (83.8)	11591 (78.8)
Injection drug user (IDU)	192 (2.0)	233 (15.1)	60 (2.4)	21 (2.4)	517 (3.5)
Gay or bisexual male IDU	1457 (15.4)	315 (20.4)	306 (12.0)	91 (10.3)	2232 (15.2)
Transfusion/Hemophiliac	6 (0.1)	3 (0.2)	2 (0.1)	3 (0.3)	14 (0.1)
Heterosexual (3)	30 (0.3)	68 (4.4)	50 (2.0)	11 (1.2)	162 (1.1)
Perinatal	1 (0.0)	3 (0.2)	2 (0.1)	2 (0.2)	10 (0.1)
Risk not reported	76 (0.8)	42 (2.7)	48 (1.9)	15 (1.7)	192 (1.3)
Subtotal	9443 (100)	1546 (100)	2553 (100)	881 (100)	14718 (100)
Female					
Injection drug user (IDU)	128 (48.5)	186 (51.1)	51 (28.5)	14 (19.2)	386 (42.7)
Lesbian or bisexual IDU	29 (11.0)	14 (3.8)	7 (3.9)	3 (4.1)	59 (6.5)
Transfusion/Hemophilia	5 (1.9)	2 (0.5)	2 (1.1)	2 (2.7)	11 (1.2)
Heterosexual (3)	79 (29.9)	134 (36.8)	90 (50.3)	44 (60.3)	350 (38.8)
Perinatal	1 (0.4)	6 (1.6)	9 (5.0)	0 (0.0)	18 (2.0)
Risk not reported	22 (8.3)	22 (6.0)	20 (11.2)	10 (13.7)	79 (8.7)
Subtotal	264 (100)	364 (100)	179 (100)	73 (100)	903 (100)
Transgender	70 (100)	120 (100)	106 (100)	40 (100)	350 (100)
Total	9777 (100)	2030 (100)	2838 (100)	994 (100)	15971 (100)

- (1) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
- (2) Total includes persons with multiple or unknown race.
- (3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.