

HIV SEMI-ANNUAL SURVEILLANCE REPORT

San Francisco Department of Public Health

HIV Cases Reported Through June 2017

Contents	Page
Surveillance Summary.....	1
Figure 1: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year	2
Figure 2: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year	2
Table 1. Diagnoses of HIV infection by age and year of initial HIV diagnosis.....	3
Table 2: Diagnoses of HIV infection stage 3 (AIDS) by age and year of AIDS diagnosis.....	4
Table 3: Diagnoses of HIV infection by gender and year of initial HIV diagnosis	5
Table 4: Diagnoses of HIV infection stage 3 (AIDS) by gender and year of AIDS diagnosis...	6
Table 5: Diagnoses of HIV infection by race/ethnicity and year of initial HIV diagnosis.....	7
Table 6: Diagnoses of HIV infection stage 3 (AIDS) by race/ethnicity and year of AIDS diagnosis.....	8
Table 7: Diagnoses of HIV infection by transmission category and year of initial HIV diagnosis.....	9
Table 8: Diagnoses of HIV infection stage 3 (AIDS) by transmission category and year of AIDS diagnosis.....	10
Table 9: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year	11
Table 10: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year	12
Table 11: Persons living with HIV infection by age, gender and race/ethnicity.....	13
Table 12: Persons living with HIV infection by transmission category, gender and race/ethnicity.....	14

The HIV Surveillance Report is published semi-annually by the
San Francisco Department of Public Health, Population Health Division
Applied Research, Community Health Epidemiology, and Surveillance Branch (ARCHES)
25 Van Ness Avenue, Suite 500, San Francisco, CA 94102; Phone (415) 437-6336, FAX (415) 431-0353

Director of Health: Barbara A. Garcia, MPA
Director of Population Health Division: Tomas Aragon, MD, DrPH
Director of ARCHES: Wayne Enanoria, PhD, MPH
Directors of HIV Surveillance Section: Susan Scheer, PhD, MPH; Ling Hsu, MPH

***The HIV Surveillance Report is accessible via internet:
<http://www.sfdph.org/dph/files/reports/default.asp>***

HIV SURVEILLANCE SUMMARY

San Francisco¹ (as of 06/30/2017)

Persons living with HIV infection ²	16,057
Diagnoses of HIV infection stage 3 (AIDS), cumulative	29,956
Deaths among persons diagnosed with HIV infection stage 3 (AIDS), cumulative	20,606

California³ (as of 12/31/2014)

Persons living with HIV infection ²	126,241
Diagnoses of HIV infection stage 3 (AIDS), cumulative	NA
Deaths among persons diagnosed with HIV infection stage 3 (AIDS), cumulative	NA

United States⁴ (as of 12/31/2014)

Persons living with HIV infection ²	972,813
Diagnoses of HIV infection stage 3 (AIDS), cumulative (as of 12/31/2015)	1,252,890
Deaths among persons diagnosed with HIV infection stage 3 (AIDS), cumulative	704,032


¹ Residents of San Francisco at time of HIV or AIDS diagnosis.

² Persons diagnosed with HIV infection regardless of disease stage.

³ CA data source: <http://www.cdph.ca.gov/programs/aids/Pages/TOASurv.aspx>.


⁴ US data source: CDC HIV Surveillance Report 2015, vol. 27 (Table 2b, 13b, 18b).
<https://www.cdc.gov/hiv/library/reports/hiv-surveillance.html>.

Figure 1: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year, San Francisco, 2002-2017*



* Reporting for recent years is incomplete. See Table 9 for actual numbers per year.

Figure 2: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year, San Francisco, 1980-2017 †



† Reporting for recent years is incomplete. See Table 10 for actual numbers per year.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2017

Table 1. Diagnoses of HIV infection by age and year of initial HIV diagnosis (1), San Francisco, 2006-2017

Age at initial HIV diagnosis (years)	Year of initial HIV diagnosis					
	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	0 (0.0)	3 (0.6)	4 (0.8)	1 (0.2)	3 (0.7)	1 (0.2)
18 - 24	65 (12.2)	53 (10.0)	56 (10.8)	60 (12.8)	56 (12.1)	44 (10.5)
25 - 29	64 (12.0)	101 (19.0)	80 (15.4)	65 (13.9)	63 (13.7)	66 (15.7)
30 - 39	180 (33.8)	184 (34.7)	182 (35.1)	141 (30.1)	145 (31.5)	111 (26.4)
40 - 49	150 (28.2)	131 (24.7)	150 (28.9)	122 (26.0)	129 (28.0)	131 (31.2)
50 - 59	54 (10.2)	46 (8.7)	39 (7.5)	63 (13.4)	49 (10.6)	52 (12.4)
60 +	19 (3.6)	13 (2.4)	8 (1.5)	17 (3.6)	16 (3.5)	15 (3.6)
Total	532 (100)	531 (100)	519 (100)	469 (100)	461 (100)	420 (100)

Age at initial HIV diagnosis (years)	Year of initial HIV diagnosis					
	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	0 (0.0)	0 (0.0)	1 (0.3)	3 (1.1)	0 (0.0)	2 (1.9)
18 - 24	59 (13.0)	55 (14.0)	37 (11.9)	35 (13.1)	31 (13.8)	13 (12.5)
25 - 29	73 (16.0)	79 (20.1)	55 (17.6)	60 (22.5)	53 (23.7)	16 (15.4)
30 - 39	139 (30.5)	113 (28.8)	92 (29.5)	82 (30.7)	73 (32.6)	38 (36.5)
40 - 49	130 (28.6)	98 (24.9)	74 (23.7)	57 (21.3)	34 (15.2)	20 (19.2)
50 - 59	42 (9.2)	35 (8.9)	40 (12.8)	17 (6.4)	23 (10.3)	10 (9.6)
60 +	12 (2.6)	13 (3.3)	13 (4.2)	13 (4.9)	10 (4.5)	5 (4.8)
Total	455 (100)	393 (100)	312 (100)	267 (100)	224 (100)	104 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of positive HIV antibody test, positive HIV antigen/antibody combination test, detectable viral load, physician-documented diagnosis in absence of sufficient laboratory evidence, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2017

Table 2. Diagnoses of HIV infection stage 3 (AIDS) by age and year of AIDS diagnosis (1), San Francisco, 2006-2017

Age at AIDS diagnosis (years)	Year of AIDS diagnosis					
	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	0 (0.0)	1 (0.2)	1 (0.2)	1 (0.3)	0 (0.0)	0 (0.0)
18 - 24	13 (2.9)	15 (3.4)	17 (3.9)	15 (4.6)	15 (5.0)	12 (4.8)
25 - 29	25 (5.6)	39 (8.7)	31 (7.1)	23 (7.1)	27 (9.0)	20 (8.0)
30 - 39	125 (27.8)	133 (29.8)	123 (28.3)	75 (23.1)	74 (24.6)	62 (24.7)
40 - 49	185 (41.1)	167 (37.4)	167 (38.4)	120 (37.0)	100 (33.2)	99 (39.4)
50 - 59	81 (18.0)	65 (14.6)	75 (17.2)	69 (21.3)	62 (20.6)	49 (19.5)
60 +	21 (4.7)	26 (5.8)	21 (4.8)	21 (6.5)	23 (7.6)	9 (3.6)
Total	450 (100)	446 (100)	435 (100)	324 (100)	301 (100)	251 (100)

Age at AIDS diagnosis (years)	Year of AIDS diagnosis					
	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.9)
18 - 24	11 (4.5)	10 (5.3)	5 (3.6)	4 (3.3)	2 (2.1)	1 (1.9)
25 - 29	25 (10.3)	17 (9.0)	14 (10.0)	8 (6.5)	8 (8.4)	6 (11.5)
30 - 39	60 (24.8)	43 (22.8)	40 (28.6)	34 (27.6)	27 (28.4)	14 (26.9)
40 - 49	82 (33.9)	59 (31.2)	40 (28.6)	35 (28.5)	22 (23.2)	19 (36.5)
50 - 59	44 (18.2)	39 (20.6)	27 (19.3)	28 (22.8)	17 (17.9)	3 (5.8)
60 +	20 (8.3)	21 (11.1)	14 (10.0)	14 (11.4)	19 (20.0)	8 (15.4)
Total	242 (100)	189 (100)	140 (100)	123 (100)	95 (100)	52 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2017

Table 3. Diagnoses of HIV infection by gender and year of initial HIV diagnosis (1), San Francisco, 2006-2017

Gender	Year of initial HIV diagnosis					
	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)
Male	483 (90.8)	459 (86.4)	462 (89.0)	417 (88.9)	410 (88.9)	365 (86.9)
Female	36 (6.8)	48 (9.0)	40 (7.7)	30 (6.4)	36 (7.8)	41 (9.8)
Transgender	13 (2.4)	24 (4.5)	17 (3.3)	22 (4.7)	15 (3.3)	14 (3.3)
Total	532 (100)	531 (100)	519 (100)	469 (100)	461 (100)	420 (100)

Gender	Year of initial HIV diagnosis					
	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)
Male	424 (93.2)	360 (91.6)	291 (93.3)	234 (87.6)	192 (85.7)	88 (84.6)
Female	23 (5.1)	22 (5.6)	13 (4.2)	26 (9.7)	25 (11.2)	14 (13.5)
Transgender	8 (1.8)	11 (2.8)	8 (2.6)	7 (2.6)	7 (3.1)	2 (1.9)
Total	455 (100)	393 (100)	312 (100)	267 (100)	224 (100)	104 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of positive HIV antibody test, positive HIV antigen/antibody combination test, detectable viral load, physician-documented diagnosis in absence of sufficient laboratory evidence, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2017

Table 4. Diagnoses of HIV infection stage 3 (AIDS) by gender and year of AIDS diagnosis (1), San Francisco, 2006-2017

Gender	Year of AIDS diagnosis					
	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)
Male	402 (89.3)	401 (89.9)	376 (86.4)	288 (88.9)	269 (89.4)	213 (84.9)
Female	32 (7.1)	32 (7.2)	43 (9.9)	24 (7.4)	16 (5.3)	24 (9.6)
Transgender	16 (3.6)	13 (2.9)	16 (3.7)	12 (3.7)	16 (5.3)	14 (5.6)
Total	450 (100)	446 (100)	435 (100)	324 (100)	301 (100)	251 (100)

Gender	Year of AIDS diagnosis					
	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)
Male	210 (86.8)	166 (87.8)	125 (89.3)	99 (80.5)	82 (86.3)	42 (80.8)
Female	24 (9.9)	17 (9.0)	7 (5.0)	18 (14.6)	10 (10.5)	8 (15.4)
Transgender	8 (3.3)	6 (3.2)	8 (5.7)	6 (4.9)	3 (3.2)	2 (3.8)
Total	242 (100)	189 (100)	140 (100)	123 (100)	95 (100)	52 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2017

Table 5. Diagnoses of HIV infection by race/ethnicity and year of initial HIV diagnosis (1), San Francisco, 2006-2017

Race/ethnicity	Year of initial HIV diagnosis					
	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)
White	291 (54.7)	269 (50.7)	255 (49.1)	226 (48.2)	220 (47.7)	219 (52.1)
African American	77 (14.5)	80 (15.1)	80 (15.4)	64 (13.6)	63 (13.7)	64 (15.2)
Latino	113 (21.2)	105 (19.8)	119 (22.9)	112 (23.9)	111 (24.1)	83 (19.8)
Asian/Pacific Islander/ Native American	33 (6.2)	49 (9.2)	43 (8.3)	40 (8.5)	41 (8.9)	36 (8.6)
Multiple races	18 (3.4)	28 (5.3)	22 (4.2)	26 (5.5)	25 (5.4)	17 (4.0)
Total (2)	532 (100)	531 (100)	519 (100)	469 (100)	461 (100)	420 (100)

Race/ethnicity	Year of initial HIV diagnosis					
	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)
White	227 (49.9)	178 (45.3)	136 (43.6)	110 (41.2)	87 (38.8)	36 (34.6)
African American	46 (10.1)	50 (12.7)	32 (10.3)	40 (15.0)	34 (15.2)	15 (14.4)
Latino	110 (24.2)	98 (24.9)	87 (27.9)	69 (25.8)	63 (28.1)	28 (26.9)
Asian/Pacific Islander/ Native American	57 (12.5)	51 (13.0)	43 (13.8)	32 (12.0)	35 (15.6)	18 (17.3)
Multiple races	14 (3.1)	14 (3.6)	14 (4.5)	14 (5.2)	5 (2.2)	7 (6.7)
Total (2)	455 (100)	393 (100)	312 (100)	267 (100)	224 (100)	104 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of positive HIV antibody test, positive HIV antigen/antibody combination test, detectable viral load, physician-documented diagnosis in absence of sufficient laboratory evidence, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

(2) Total includes persons with unknown race.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2017

Table 6. Diagnoses of HIV infection stage 3 (AIDS) by race/ethnicity and year of AIDS diagnosis (1), San Francisco, 2006-2017

Race/ethnicity	Year of AIDS diagnosis					
	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)
White	248 (55.1)	255 (57.2)	219 (50.3)	171 (52.8)	145 (48.2)	129 (51.4)
African American	82 (18.2)	74 (16.6)	80 (18.4)	60 (18.5)	52 (17.3)	45 (17.9)
Latino	79 (17.6)	76 (17.0)	77 (17.7)	66 (20.4)	63 (20.9)	46 (18.3)
Asian/Pacific Islander/ Native American	25 (5.6)	26 (5.8)	39 (9.0)	15 (4.6)	25 (8.3)	15 (6.0)
Multiple races	16 (3.6)	15 (3.4)	20 (4.6)	12 (3.7)	16 (5.3)	16 (6.4)
Total (2)	450 (100)	446 (100)	435 (100)	324 (100)	301 (100)	251 (100)

Race/ethnicity	Year of AIDS diagnosis					
	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)
White	121 (50.0)	93 (49.2)	52 (37.1)	56 (45.5)	46 (48.4)	20 (38.5)
African American	39 (16.1)	32 (16.9)	29 (20.7)	19 (15.4)	15 (15.8)	9 (17.3)
Latino	54 (22.3)	34 (18.0)	31 (22.1)	32 (26.0)	20 (21.1)	15 (28.8)
Asian/Pacific Islander/ Native American	15 (6.2)	18 (9.5)	18 (12.9)	5 (4.1)	9 (9.5)	6 (11.5)
Multiple races	13 (5.4)	12 (6.3)	10 (7.1)	10 (8.1)	5 (5.3)	2 (3.8)
Total (2)	242 (100)	189 (100)	140 (100)	123 (100)	95 (100)	52 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

(2) Total includes persons with unknown race.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2017

Table 7. Diagnoses of HIV infection by transmission category and year of initial HIV diagnosis (1), San Francisco, 2006-2017

Transmission category (2)	Year of initial HIV diagnosis					
	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)
Gay or bisexual male	366 (68.8)	343 (64.6)	373 (71.9)	326 (69.5)	300 (65.1)	300 (71.4)
Injection drug user (IDU)	38 (7.1)	40 (7.5)	34 (6.6)	25 (5.3)	37 (8.0)	28 (6.7)
Gay or bisexual male IDU	93 (17.5)	90 (16.9)	66 (12.7)	80 (17.1)	69 (15.0)	55 (13.1)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	26 (4.9)	43 (8.1)	35 (6.7)	25 (5.3)	37 (8.0)	28 (6.7)
Perinatal	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Other/Unknown	9 (1.7)	15 (2.8)	11 (2.1)	13 (2.8)	18 (3.9)	9 (2.1)
Total	532 (100)	531 (100)	519 (100)	469 (100)	461 (100)	420 (100)

Transmission category (2)	Year of initial HIV diagnosis					
	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)
Gay or bisexual male	351 (77.1)	303 (77.1)	231 (74.0)	196 (73.4)	157 (70.1)	71 (68.3)
Injection drug user (IDU)	20 (4.4)	21 (5.3)	23 (7.4)	17 (6.4)	20 (8.9)	12 (11.5)
Gay or bisexual male IDU	47 (10.3)	45 (11.5)	38 (12.2)	24 (9.0)	21 (9.4)	7 (6.7)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	25 (5.5)	16 (4.1)	9 (2.9)	18 (6.7)	15 (6.7)	8 (7.7)
Perinatal	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Other/Unknown	12 (2.6)	8 (2.0)	11 (3.5)	12 (4.5)	11 (4.9)	6 (5.8)
Total	455 (100)	393 (100)	312 (100)	267 (100)	224 (100)	104 (100)

- (1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of positive HIV antibody test, positive HIV antigen/antibody combination test, detectable viral load, physician-documented diagnosis in absence of sufficient laboratory evidence, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.
- (2) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
- (3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2017

Table 8. Diagnoses of HIV infection stage 3 (AIDS) by transmission category and year of AIDS diagnosis (1), San Francisco, 2006-2017

Transmission category (2)	Year of AIDS diagnosis					
	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)
Gay or bisexual male	286 (63.6)	288 (64.6)	262 (60.2)	211 (65.1)	171 (56.8)	149 (59.4)
Injection drug user (IDU)	39 (8.7)	38 (8.5)	34 (7.8)	35 (10.8)	23 (7.6)	34 (13.5)
Gay or bisexual male IDU	99 (22.0)	87 (19.5)	97 (22.3)	53 (16.4)	84 (27.9)	47 (18.7)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.3)	0 (0.0)	0 (0.0)
Heterosexual (3)	20 (4.4)	29 (6.5)	32 (7.4)	18 (5.6)	18 (6.0)	17 (6.8)
Perinatal	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.3)	0 (0.0)	1 (0.4)
Other/Unknown	6 (1.3)	4 (0.9)	10 (2.3)	5 (1.5)	5 (1.7)	3 (1.2)
Total	450 (100)	446 (100)	435 (100)	324 (100)	301 (100)	251 (100)

Transmission category (2)	Year of AIDS diagnosis					
	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)
Gay or bisexual male	153 (63.2)	112 (59.3)	95 (67.9)	68 (55.3)	55 (57.9)	27 (51.9)
Injection drug user (IDU)	23 (9.5)	19 (10.1)	15 (10.7)	17 (13.8)	13 (13.7)	5 (9.6)
Gay or bisexual male IDU	39 (16.1)	45 (23.8)	21 (15.0)	23 (18.7)	16 (16.8)	11 (21.2)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	20 (8.3)	12 (6.3)	6 (4.3)	10 (8.1)	9 (9.5)	7 (13.5)
Perinatal	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Other/Unknown	7 (2.9)	1 (0.5)	3 (2.1)	5 (4.1)	2 (2.1)	2 (3.8)
Total	242 (100)	189 (100)	140 (100)	123 (100)	95 (100)	52 (100)

- (1) Data in recent years are incomplete due to delays in case reporting.
(2) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
(3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2017

Table 9. Diagnoses of HIV infection, HIV deaths and persons living with HIV by year, San Francisco, 2002-2017

Year	Number of cases diagnosed per year (1)	Number of deaths occurred per year (1)	Number of persons living with HIV infection (1)
2002 (2)	664	350	13363
2003	655	338	13680
2004	689	343	14026
2005	571	348	14249
2006	532	327	14454
2007	531	324	14661
2008	519	264	14916
2009	469	254	15131
2010	461	247	15345
2011	420	234	15531
2012	455	237	15749
2013	393	261	15881
2014	312	241	15952
2015	267	257	15962
2016	224	167	16019
2017	104	66	16057

(1) Data in recent years are incomplete due to delays in case/death reporting.

(2) Year of the mandatory reporting of HIV non-AIDS cases in California began.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2017

Table 10. Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year, San Francisco, 1980-2017

Year	Number of cases diagnosed per year (1)	Number of deaths occurred per year (1)	Number of persons living with AIDS (1)
1980	3	0	3
1981	26	8	21
1982	99	32	88
1983	274	111	251
1984	558	273	536
1985	860	534	862
1986	1236	807	1291
1987	1632	878	2045
1988	1763	1040	2768
1989	2161	1279	3650
1990	2046	1365	4331
1991	2285	1512	5104
1992	2329	1640	5793
1993	2069	1603	6259
1994	1781	1601	6439
1995	1556	1485	6510
1996	1076	995	6591
1997	803	424	6970
1998	693	402	7261
1999	577	353	7485
2000	557	350	7692
2001	512	324	7880
2002	493	323	8050
2003	556	303	8303
2004	483	309	8477
2005	480	312	8645
2006	450	289	8806
2007	446	271	8981
2008	435	229	9187
2009	324	211	9300
2010	301	195	9406
2011	251	188	9469
2012	242	184	9527
2013	189	192	9524
2014	140	189	9475
2015	123	204	9394
2016	95	138	9351
2017	52	53	9350
Total	29956	20606	

(1) Data in recent years are incomplete due to delays in case/death reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2017

Table 11. Persons living with HIV infection by age, gender and race/ethnicity, San Francisco

Current age (years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander/ Native American No. (%)	Multiple races No. (%)	Total (1) No. (%)
Male						
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	0 (0.0)	0 (0.0)	1 (0.0)	1 (0.1)	1 (0.2)	3 (0.0)
18 - 24	21 (0.2)	16 (1.1)	23 (0.8)	15 (1.6)	4 (0.9)	79 (0.5)
25 - 29	117 (1.3)	52 (3.5)	125 (4.5)	46 (5.0)	14 (3.2)	354 (2.4)
30 - 39	677 (7.4)	140 (9.5)	491 (17.8)	190 (20.5)	66 (15.2)	1565 (10.6)
40 - 49	1750 (19.1)	277 (18.9)	747 (27.2)	294 (31.7)	123 (28.3)	3195 (21.7)
50 - 59	3599 (39.3)	509 (34.7)	941 (34.2)	268 (28.9)	147 (33.9)	5465 (37.0)
60 +	2999 (32.7)	474 (32.3)	423 (15.4)	114 (12.3)	79 (18.2)	4091 (27.7)
Subtotal	9163 (100)	1468 (100)	2751 (100)	928 (100)	434 (100)	14752 (100)
Female						
0 - 12	0 (0.0)	0 (0.0)	1 (0.5)	0 (0.0)	0 (0.0)	1 (0.1)
13 - 17	0 (0.0)	1 (0.3)	0 (0.0)	0 (0.0)	1 (2.9)	2 (0.2)
18 - 24	3 (1.1)	6 (1.7)	8 (4.1)	1 (1.4)	3 (8.6)	21 (2.3)
25 - 29	5 (1.9)	7 (2.0)	8 (4.1)	2 (2.7)	0 (0.0)	22 (2.4)
30 - 39	36 (13.5)	33 (9.5)	38 (19.3)	13 (17.8)	4 (11.4)	124 (13.5)
40 - 49	59 (22.1)	74 (21.4)	41 (20.8)	21 (28.8)	12 (34.3)	208 (22.6)
50 - 59	110 (41.2)	133 (38.4)	55 (27.9)	23 (31.5)	11 (31.4)	333 (36.2)
60 +	54 (20.2)	92 (26.6)	46 (23.4)	13 (17.8)	4 (11.4)	209 (22.7)
Subtotal	267 (100)	346 (100)	197 (100)	73 (100)	35 (100)	920 (100)
Transgender	75 (100)	126 (100)	126 (100)	40 (100)	18 (100)	385 (100)
Total	9505 (100)	1940 (100)	3074 (100)	1041 (100)	487 (100)	16057 (100)

(1) Total includes persons with unknown race.

HIV Semi-Annual Surveillance Report Cases Reported as of 06/30/2017

Table 12. Persons living with HIV infection by transmission category, gender and race/ethnicity, San Francisco

Transmission category (1)	White		African American		Latino		Asian/ Pacific Islander/ Native American		Multiple races		Total (2)	
	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
Male												
Gay or bisexual male	7479	(81.6)	865	(58.9)	2256	(82.0)	791	(85.2)	312	(71.9)	11706	(79.4)
Injection drug user (IDU)	182	(2.0)	203	(13.8)	66	(2.4)	20	(2.2)	16	(3.7)	487	(3.3)
Gay or bisexual male IDU	1404	(15.3)	292	(19.9)	327	(11.9)	79	(8.5)	95	(21.9)	2197	(14.9)
Transfusion/Hemophiliac	6	(0.1)	2	(0.1)	2	(0.1)	3	(0.3)	1	(0.3)	14	(0.2)
Heterosexual (3)	30	(0.3)	65	(4.4)	53	(1.9)	18	(1.9)	5	(1.2)	171	(1.2)
Perinatal	1	(0.0)	3	(0.2)	2	(0.1)	1	(0.1)	2	(0.5)	9	(0.1)
Other/Unknown	61	(0.7)	38	(2.6)	45	(1.6)	16	(1.7)	3	(0.7)	168	(1.1)
Subtotal	9163	(100)	1468	(100)	2751	(100)	928	(100)	434	(100)	14752	(100)
Female												
Injection drug user (IDU)	150	(56.2)	177	(51.2)	62	(31.5)	15	(20.5)	23	(65.7)	427	(46.4)
Transfusion/Hemophilia	5	(1.9)	2	(0.6)	2	(1.0)	2	(2.7)	0	(0.0)	11	(1.2)
Heterosexual (3)	92	(34.5)	141	(40.8)	104	(52.8)	49	(67.1)	7	(20.0)	393	(42.7)
Perinatal	1	(0.4)	6	(1.7)	9	(4.6)	0	(0.0)	2	(5.7)	18	(2.0)
Other/Unknown	19	(7.1)	20	(5.8)	20	(10.2)	7	(9.6)	3	(8.6)	71	(7.7)
Subtotal	267	(100)	346	(100)	197	(100)	73	(100)	35	(100)	920	(100)
Transgender	75	(100)	126	(100)	126	(100)	40	(100)	18	(100)	385	(100)
Total	9505	(100)	1940	(100)	3074	(100)	1041	(100)	487	(100)	16057	(100)

(1) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.

(2) Total includes persons with unknown race.

(3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.