

HIV SEMI-ANNUAL SURVEILLANCE REPORT

San Francisco Department of Public Health

HIV Cases Reported Through December 2017

Contents	Page
Surveillance Summary.....	1
Figure 1: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year	2
Figure 2: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year	2
Table 1. Diagnoses of HIV infection by age and year of initial HIV diagnosis.....	3
Table 2: Diagnoses of HIV infection stage 3 (AIDS) by age and year of AIDS diagnosis.....	4
Table 3: Diagnoses of HIV infection by gender and year of initial HIV diagnosis	5
Table 4: Diagnoses of HIV infection stage 3 (AIDS) by gender and year of AIDS diagnosis...	6
Table 5: Diagnoses of HIV infection by race/ethnicity and year of initial HIV diagnosis.....	7
Table 6: Diagnoses of HIV infection stage 3 (AIDS) by race/ethnicity and year of AIDS diagnosis.....	8
Table 7: Diagnoses of HIV infection by transmission category and year of initial HIV diagnosis.....	9
Table 8: Diagnoses of HIV infection stage 3 (AIDS) by transmission category and year of AIDS diagnosis.....	10
Table 9: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year	11
Table 10: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year	12
Table 11: Persons living with HIV infection by age, gender and race/ethnicity.....	13
Table 12: Persons living with HIV infection by transmission category, gender and race/ethnicity.....	14

The HIV Surveillance Report is published semi-annually by the
San Francisco Department of Public Health, Population Health Division
Applied Research, Community Health Epidemiology, and Surveillance Branch (ARCHES)
25 Van Ness Avenue, Suite 500, San Francisco, CA 94102; Phone (415) 437-6336, FAX (415) 431-0353

Director of Health: Barbara A. Garcia, MPA
Director of Population Health Division: Tomas Aragon, MD, DrPH
Director of ARCHES: Wayne Enanoria, PhD, MPH
Directors of HIV Surveillance Section: Susan Scheer, PhD, MPH; Ling Hsu, MPH

***The HIV Surveillance Report is accessible via internet:
<http://www.sfdph.org/dph/files/reports/default.asp>***

HIV SURVEILLANCE SUMMARY

San Francisco¹ (as of 12/31/2017)

Persons living with HIV infection ²	16,011
Diagnoses of HIV infection stage 3 (AIDS), cumulative	30,013
Deaths among persons diagnosed with HIV infection stage 3 (AIDS), cumulative	20,730

California³ (as of 12/31/2015)

Persons living with HIV infection ²	128,415
Diagnoses of HIV infection stage 3 (AIDS), cumulative	NA
Deaths among persons diagnosed with HIV infection stage 3 (AIDS), cumulative	NA

United States⁴ (as of 12/31/2015)

Persons living with HIV infection ²	991,289
Diagnoses of HIV infection stage 3 (AIDS), cumulative (as of 12/31/2016)	1,268,595
Deaths among persons diagnosed with HIV infection stage 3 (AIDS), cumulative	718,906

¹ Residents of San Francisco at time of HIV or AIDS diagnosis.

² Persons diagnosed with HIV infection regardless of disease stage.


³ CA data source: California HIV Surveillance Report 2015.

https://www.cdph.ca.gov/Programs/CID/DOA/Pages/OA_case_surveillance_reports.aspx

⁴ US data source: CDC HIV Surveillance Report 2016, vol. 28 (Table 2b, 17b, 20b).


<https://www.cdc.gov/hiv/library/reports/hiv-surveillance.html>.

Figure 1: Diagnoses of HIV infection, HIV deaths and persons living with HIV by year, San Francisco, 2002-2017*



* Reporting for recent years is incomplete. See Table 9 for actual numbers per year.

Figure 2: Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year, San Francisco, 1980-2017 †



† Reporting for recent years is incomplete. See Table 10 for actual numbers per year.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2017

Table 1. Diagnoses of HIV infection by age and year of initial HIV diagnosis (1), San Francisco, 2006-2017

Age at initial HIV diagnosis (years)	Year of initial HIV diagnosis					
	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	0 (0.0)	3 (0.6)	4 (0.8)	1 (0.2)	3 (0.7)	0 (0.0)
18 - 24	66 (12.4)	53 (9.9)	55 (10.6)	61 (13.0)	56 (12.2)	44 (10.4)
25 - 29	64 (12.0)	102 (19.1)	80 (15.4)	65 (13.8)	63 (13.7)	69 (16.3)
30 - 39	180 (33.8)	184 (34.5)	182 (35.0)	141 (29.9)	144 (31.4)	113 (26.7)
40 - 49	150 (28.1)	132 (24.8)	150 (28.8)	123 (26.1)	128 (27.9)	131 (30.9)
50 - 59	54 (10.1)	46 (8.6)	39 (7.5)	63 (13.4)	49 (10.7)	52 (12.3)
60 +	19 (3.6)	13 (2.4)	10 (1.9)	17 (3.6)	16 (3.5)	15 (3.5)
Total	533 (100)	533 (100)	520 (100)	471 (100)	459 (100)	424 (100)

Age at initial HIV diagnosis (years)	Year of initial HIV diagnosis					
	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	0 (0.0)	0 (0.0)	1 (0.3)	3 (1.1)	0 (0.0)	4 (2.0)
18 - 24	59 (12.9)	55 (14.1)	38 (12.1)	35 (13.0)	31 (13.4)	22 (10.9)
25 - 29	74 (16.2)	78 (19.9)	55 (17.5)	61 (22.6)	56 (24.1)	36 (17.8)
30 - 39	141 (30.9)	113 (28.9)	91 (29.0)	82 (30.4)	75 (32.3)	67 (33.2)
40 - 49	129 (28.2)	97 (24.8)	74 (23.6)	57 (21.1)	35 (15.1)	41 (20.3)
50 - 59	42 (9.2)	35 (9.0)	41 (13.1)	18 (6.7)	24 (10.3)	23 (11.4)
60 +	12 (2.6)	13 (3.3)	14 (4.5)	14 (5.2)	11 (4.7)	9 (4.5)
Total	457 (100)	391 (100)	314 (100)	270 (100)	232 (100)	202 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of positive HIV antibody test, positive HIV antigen/antibody combination test, detectable viral load, physician-documented diagnosis in absence of sufficient laboratory evidence, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report

Cases Reported as of 12/31/2017

Table 2. Diagnoses of HIV infection stage 3 (AIDS) by age and year of AIDS diagnosis (1), San Francisco, 2006-2017

Age at AIDS diagnosis (years)	Year of AIDS diagnosis					
	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	0 (0.0)	1 (0.2)	1 (0.2)	1 (0.3)	0 (0.0)	0 (0.0)
18 - 24	13 (2.9)	15 (3.4)	17 (3.9)	15 (4.6)	15 (5.0)	12 (4.8)
25 - 29	25 (5.6)	39 (8.7)	31 (7.1)	23 (7.1)	27 (9.0)	20 (8.0)
30 - 39	125 (27.8)	134 (30.0)	123 (28.3)	75 (23.1)	74 (24.6)	62 (24.7)
40 - 49	185 (41.1)	167 (37.4)	167 (38.4)	120 (37.0)	100 (33.2)	99 (39.4)
50 - 59	81 (18.0)	65 (14.5)	75 (17.2)	69 (21.3)	62 (20.6)	49 (19.5)
60 +	21 (4.7)	26 (5.8)	21 (4.8)	21 (6.5)	23 (7.6)	9 (3.6)
Total	450 (100)	447 (100)	435 (100)	324 (100)	301 (100)	251 (100)

Age at AIDS diagnosis (years)	Year of AIDS diagnosis					
	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.9)
18 - 24	11 (4.5)	10 (5.3)	5 (3.6)	4 (3.3)	2 (2.1)	1 (0.9)
25 - 29	25 (10.3)	17 (9.0)	14 (10.0)	8 (6.5)	8 (8.3)	13 (11.6)
30 - 39	60 (24.7)	43 (22.8)	41 (29.3)	34 (27.6)	27 (28.1)	27 (24.1)
40 - 49	83 (34.2)	59 (31.2)	40 (28.6)	35 (28.5)	22 (22.9)	38 (33.9)
50 - 59	44 (18.1)	39 (20.6)	27 (19.3)	28 (22.8)	18 (18.8)	10 (8.9)
60 +	20 (8.2)	21 (11.1)	13 (9.3)	14 (11.4)	19 (19.8)	22 (19.6)
Total	243 (100)	189 (100)	140 (100)	123 (100)	96 (100)	112 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2017

Table 3. Diagnoses of HIV infection by gender and year of initial HIV diagnosis (1), San Francisco, 2006-2017

Gender	Year of initial HIV diagnosis					
	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)
Male	483 (90.6)	459 (86.1)	463 (89.0)	420 (89.2)	408 (88.9)	369 (87.0)
Female	36 (6.8)	49 (9.2)	40 (7.7)	30 (6.4)	36 (7.8)	41 (9.7)
Transgender	14 (2.6)	25 (4.7)	17 (3.3)	21 (4.5)	15 (3.3)	14 (3.3)
Total	533 (100)	533 (100)	520 (100)	471 (100)	459 (100)	424 (100)

Gender	Year of initial HIV diagnosis					
	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)
Male	426 (93.2)	357 (91.3)	293 (93.3)	237 (87.8)	201 (86.6)	176 (87.1)
Female	23 (5.0)	22 (5.6)	13 (4.1)	26 (9.6)	24 (10.3)	22 (10.9)
Transgender	8 (1.8)	12 (3.1)	8 (2.5)	7 (2.6)	7 (3.0)	4 (2.0)
Total	457 (100)	391 (100)	314 (100)	270 (100)	232 (100)	202 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of positive HIV antibody test, positive HIV antigen/antibody combination test, detectable viral load, physician-documented diagnosis in absence of sufficient laboratory evidence, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2017

Table 4. Diagnoses of HIV infection stage 3 (AIDS) by gender and year of AIDS diagnosis (1), San Francisco, 2006-2017

Gender	Year of AIDS diagnosis					
	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)
Male	402 (89.3)	402 (89.9)	375 (86.2)	288 (88.9)	270 (89.7)	213 (84.9)
Female	32 (7.1)	32 (7.2)	43 (9.9)	24 (7.4)	16 (5.3)	24 (9.6)
Transgender	16 (3.6)	13 (2.9)	17 (3.9)	12 (3.7)	15 (5.0)	14 (5.6)
Total	450 (100)	447 (100)	435 (100)	324 (100)	301 (100)	251 (100)

Gender	Year of AIDS diagnosis					
	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)
Male	211 (86.8)	166 (87.8)	126 (90.0)	99 (80.5)	83 (86.5)	93 (83.0)
Female	24 (9.9)	17 (9.0)	7 (5.0)	18 (14.6)	10 (10.4)	16 (14.3)
Transgender	8 (3.3)	6 (3.2)	7 (5.0)	6 (4.9)	3 (3.1)	3 (2.7)
Total	243 (100)	189 (100)	140 (100)	123 (100)	96 (100)	112 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2017

Table 5. Diagnoses of HIV infection by race/ethnicity and year of initial HIV diagnosis (1), San Francisco, 2006-2017

Race/ethnicity	Year of initial HIV diagnosis					
	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)
White	291 (54.6)	268 (50.3)	255 (49.0)	226 (48.0)	220 (47.9)	219 (51.7)
African American	77 (14.4)	81 (15.2)	80 (15.4)	64 (13.6)	62 (13.5)	64 (15.1)
Latino	114 (21.4)	107 (20.1)	118 (22.7)	114 (24.2)	109 (23.7)	85 (20.0)
Asian/Pacific Islander/ Native American	33 (6.2)	48 (9.0)	44 (8.5)	40 (8.5)	41 (8.9)	37 (8.7)
Multiple races	18 (3.4)	29 (5.4)	23 (4.4)	26 (5.5)	26 (5.7)	18 (4.2)
Total (2)	533 (100)	533 (100)	520 (100)	471 (100)	459 (100)	424 (100)

Race/ethnicity	Year of initial HIV diagnosis					
	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)
White	227 (49.7)	179 (45.8)	136 (43.3)	112 (41.5)	88 (37.9)	84 (41.6)
African American	46 (10.1)	49 (12.5)	30 (9.6)	40 (14.8)	33 (14.2)	32 (15.8)
Latino	111 (24.3)	97 (24.8)	89 (28.3)	70 (25.9)	68 (29.3)	48 (23.8)
Asian/Pacific Islander/ Native American	58 (12.7)	49 (12.5)	44 (14.0)	32 (11.9)	37 (15.9)	27 (13.4)
Multiple races	14 (3.1)	15 (3.8)	15 (4.8)	14 (5.2)	6 (2.6)	11 (5.4)
Total (2)	457 (100)	391 (100)	314 (100)	270 (100)	232 (100)	202 (100)

(1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of positive HIV antibody test, positive HIV antigen/antibody combination test, detectable viral load, physician-documented diagnosis in absence of sufficient laboratory evidence, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.

(2) Total includes persons with unknown race.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2017

Table 6. Diagnoses of HIV infection stage 3 (AIDS) by race/ethnicity and year of AIDS diagnosis (1), San Francisco, 2006-2017

Race/ethnicity	Year of AIDS diagnosis					
	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)
White	247 (54.9)	254 (56.8)	219 (50.3)	171 (52.8)	145 (48.2)	129 (51.4)
African American	82 (18.2)	74 (16.6)	80 (18.4)	60 (18.5)	50 (16.6)	45 (17.9)
Latino	80 (17.8)	78 (17.4)	77 (17.7)	66 (20.4)	64 (21.3)	46 (18.3)
Asian/Pacific Islander/ Native American	24 (5.3)	26 (5.8)	39 (9.0)	15 (4.6)	25 (8.3)	15 (6.0)
Multiple races	17 (3.8)	15 (3.4)	20 (4.6)	12 (3.7)	17 (5.6)	16 (6.4)
Total (2)	450 (100)	447 (100)	435 (100)	324 (100)	301 (100)	251 (100)

Race/ethnicity	Year of AIDS diagnosis					
	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)
White	122 (50.2)	93 (49.2)	52 (37.1)	56 (45.5)	46 (47.9)	49 (43.8)
African American	39 (16.0)	32 (16.9)	29 (20.7)	19 (15.4)	15 (15.6)	21 (18.8)
Latino	54 (22.2)	34 (18.0)	32 (22.9)	32 (26.0)	20 (20.8)	29 (25.9)
Asian/Pacific Islander/ Native American	15 (6.2)	18 (9.5)	17 (12.1)	5 (4.1)	10 (10.4)	10 (8.9)
Multiple races	13 (5.3)	12 (6.3)	10 (7.1)	10 (8.1)	5 (5.2)	3 (2.7)
Total (2)	243 (100)	189 (100)	140 (100)	123 (100)	96 (100)	112 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

(2) Total includes persons with unknown race.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2017

Table 7. Diagnoses of HIV infection by transmission category and year of initial HIV diagnosis (1), San Francisco, 2006-2017

Transmission category (2)	Year of initial HIV diagnosis					
	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)
Gay or bisexual male	367 (68.9)	342 (64.2)	370 (71.2)	327 (69.4)	299 (65.1)	301 (71.0)
Injection drug user (IDU)	38 (7.1)	40 (7.5)	33 (6.3)	25 (5.3)	37 (8.1)	28 (6.6)
Gay or bisexual male IDU	93 (17.4)	92 (17.3)	69 (13.3)	81 (17.2)	68 (14.8)	59 (13.9)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	26 (4.9)	43 (8.1)	36 (6.9)	25 (5.3)	37 (8.1)	29 (6.8)
Perinatal	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Other/Unknown	9 (1.7)	16 (3.0)	12 (2.3)	13 (2.8)	18 (3.9)	7 (1.7)
Total	533 (100)	533 (100)	520 (100)	471 (100)	459 (100)	424 (100)

Transmission category (2)	Year of initial HIV diagnosis					
	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)
Gay or bisexual male	353 (77.2)	300 (76.7)	231 (73.6)	200 (74.1)	164 (70.7)	123 (60.9)
Injection drug user (IDU)	20 (4.4)	21 (5.4)	23 (7.3)	17 (6.3)	20 (8.6)	23 (11.4)
Gay or bisexual male IDU	47 (10.3)	46 (11.8)	41 (13.1)	24 (8.9)	22 (9.5)	29 (14.4)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	25 (5.5)	16 (4.1)	9 (2.9)	18 (6.7)	15 (6.5)	13 (6.4)
Perinatal	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Other/Unknown	12 (2.6)	8 (2.0)	10 (3.2)	11 (4.1)	11 (4.7)	14 (6.9)
Total	457 (100)	391 (100)	314 (100)	270 (100)	232 (100)	202 (100)

- (1) Includes persons diagnosed with HIV infection regardless of disease stage. The initial year of HIV diagnosis was determined based on the earliest date of positive HIV antibody test, positive HIV antigen/antibody combination test, detectable viral load, physician-documented diagnosis in absence of sufficient laboratory evidence, or patient self-report of a positive HIV test. Data in recent years are incomplete due to delays in case reporting.
- (2) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
- (3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2017

Table 8. Diagnoses of HIV infection stage 3 (AIDS) by transmission category and year of AIDS diagnosis (1), San Francisco, 2006-2017

Transmission category (2)	Year of AIDS diagnosis					
	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)
Gay or bisexual male	286 (63.6)	287 (64.2)	261 (60.0)	210 (64.8)	171 (56.8)	149 (59.4)
Injection drug user (IDU)	39 (8.7)	39 (8.7)	34 (7.8)	35 (10.8)	23 (7.6)	34 (13.5)
Gay or bisexual male IDU	99 (22.0)	89 (19.9)	98 (22.5)	54 (16.7)	84 (27.9)	47 (18.7)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.3)	0 (0.0)	0 (0.0)
Heterosexual (3)	20 (4.4)	29 (6.5)	32 (7.4)	18 (5.6)	18 (6.0)	17 (6.8)
Perinatal	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.3)	0 (0.0)	1 (0.4)
Other/Unknown	6 (1.3)	3 (0.7)	10 (2.3)	5 (1.5)	5 (1.7)	3 (1.2)
Total	450 (100)	447 (100)	435 (100)	324 (100)	301 (100)	251 (100)

Transmission category (2)	Year of AIDS diagnosis					
	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)
Gay or bisexual male	152 (62.6)	112 (59.3)	95 (67.9)	69 (56.1)	56 (58.3)	53 (47.3)
Injection drug user (IDU)	23 (9.5)	19 (10.1)	16 (11.4)	17 (13.8)	13 (13.5)	9 (8.0)
Gay or bisexual male IDU	41 (16.9)	45 (23.8)	21 (15.0)	23 (18.7)	16 (16.7)	29 (25.9)
Transfusion/Hemophilia	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	20 (8.2)	12 (6.3)	6 (4.3)	10 (8.1)	9 (9.4)	15 (13.4)
Perinatal	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Other/Unknown	7 (2.9)	1 (0.5)	2 (1.4)	4 (3.3)	2 (2.1)	6 (5.4)
Total	243 (100)	189 (100)	140 (100)	123 (100)	96 (100)	112 (100)

(1) Data in recent years are incomplete due to delays in case reporting.

(2) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.

(3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2017

Table 9. Diagnoses of HIV infection, HIV deaths and persons living with HIV by year, San Francisco, 2002-2017

Year	Number of cases diagnosed per year (1)	Number of deaths occurred per year (1)	Number of persons living with HIV infection (1)
2002 (2)	665	350	13363
2003	655	338	13680
2004	690	343	14027
2005	568	348	14247
2006	533	327	14453
2007	533	325	14661
2008	520	265	14916
2009	471	254	15133
2010	459	247	15345
2011	424	235	15534
2012	457	237	15754
2013	391	261	15884
2014	314	241	15957
2015	270	256	15971
2016	232	236	15967
2017	202	158	16011

(1) Data in recent years are incomplete due to delays in case/death reporting.

(2) Year of the mandatory reporting of HIV non-AIDS cases in California began.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2017

Table 10. Diagnoses of HIV infection stage 3 (AIDS), AIDS deaths and persons living with AIDS by year, San Francisco, 1980-2017

Year	Number of cases diagnosed per year (1)	Number of deaths occurred per year (1)	Number of persons living with AIDS (1)
1980	3	0	3
1981	26	8	21
1982	99	32	88
1983	274	111	251
1984	557	273	535
1985	860	534	861
1986	1236	807	1290
1987	1632	878	2044
1988	1763	1040	2767
1989	2161	1279	3649
1990	2046	1365	4330
1991	2285	1512	5103
1992	2328	1640	5791
1993	2068	1603	6256
1994	1780	1601	6435
1995	1555	1485	6505
1996	1076	995	6586
1997	803	424	6965
1998	693	402	7256
1999	577	353	7480
2000	557	350	7687
2001	512	324	7875
2002	492	323	8044
2003	556	303	8297
2004	483	309	8471
2005	480	312	8639
2006	450	289	8800
2007	447	271	8976
2008	435	229	9182
2009	324	211	9295
2010	301	195	9401
2011	251	189	9463
2012	243	184	9522
2013	189	192	9519
2014	140	189	9470
2015	123	203	9390
2016	96	187	9299
2017	112	128	9283
Total	30013	20730	

(1) Data in recent years are incomplete due to delays in case/death reporting.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2017

Table 11. Persons living with HIV infection by age, gender and race/ethnicity, San Francisco

Current age (years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander/ Native American No. (%)	Multiple races No. (%)	Total (1) No. (%)
Male						
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.1)	1 (0.2)	2 (0.0)
18 - 24	23 (0.3)	15 (1.0)	23 (0.8)	11 (1.2)	5 (1.1)	77 (0.5)
25 - 29	108 (1.2)	47 (3.2)	123 (4.5)	45 (4.8)	15 (3.3)	338 (2.3)
30 - 39	666 (7.3)	148 (10.1)	475 (17.2)	191 (20.4)	64 (14.2)	1544 (10.5)
40 - 49	1635 (18.0)	266 (18.2)	741 (26.8)	289 (30.9)	129 (28.7)	3065 (20.8)
50 - 59	3577 (39.3)	501 (34.2)	951 (34.4)	273 (29.2)	144 (32.0)	5447 (37.0)
60 +	3093 (34.0)	486 (33.2)	449 (16.3)	126 (13.5)	92 (20.4)	4248 (28.9)
Subtotal	9102 (100)	1463 (100)	2762 (100)	936 (100)	450 (100)	14721 (100)
Female						
0 - 12	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
13 - 17	0 (0.0)	1 (0.3)	1 (0.5)	0 (0.0)	1 (2.8)	3 (0.3)
18 - 24	3 (1.1)	7 (2.1)	7 (3.6)	0 (0.0)	3 (8.3)	20 (2.2)
25 - 29	6 (2.3)	7 (2.1)	9 (4.6)	3 (4.2)	0 (0.0)	25 (2.8)
30 - 39	33 (12.5)	32 (9.4)	38 (19.6)	10 (13.9)	4 (11.1)	117 (12.9)
40 - 49	59 (22.3)	62 (18.3)	40 (20.6)	22 (30.6)	11 (30.6)	195 (21.5)
50 - 59	108 (40.8)	132 (38.9)	51 (26.3)	23 (31.9)	13 (36.1)	328 (36.1)
60 +	56 (21.1)	98 (28.9)	48 (24.7)	14 (19.4)	4 (11.1)	220 (24.2)
Subtotal	265 (100)	339 (100)	194 (100)	72 (100)	36 (100)	908 (100)
Transgender	72 (100)	123 (100)	129 (100)	39 (100)	19 (100)	382 (100)
Total	9439 (100)	1925 (100)	3085 (100)	1047 (100)	505 (100)	16011 (100)

(1) Total includes persons with unknown race.

HIV Semi-Annual Surveillance Report Cases Reported as of 12/31/2017

Table 12. Persons living with HIV infection by transmission category, gender and race/ethnicity, San Francisco

Transmission category (1)	White		African American		Latino		Asian/ Pacific Islander/ Native American		Multiple races		Total (2)	
	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
Male												
Gay or bisexual male	7427	(81.6)	862	(58.9)	2267	(82.1)	791	(84.5)	324	(72.0)	11674	(79.3)
Injection drug user (IDU)	178	(2.0)	198	(13.5)	67	(2.4)	20	(2.1)	16	(3.6)	479	(3.3)
Gay or bisexual male IDU	1404	(15.4)	293	(20.0)	328	(11.9)	83	(8.9)	99	(22.0)	2207	(15.0)
Transfusion/Hemophiliac	6	(0.1)	2	(0.1)	2	(0.1)	3	(0.3)	1	(0.3)	14	(0.2)
Heterosexual (3)	30	(0.3)	66	(4.5)	53	(1.9)	20	(2.1)	5	(1.1)	174	(1.2)
Perinatal	1	(0.0)	3	(0.2)	2	(0.1)	1	(0.1)	2	(0.4)	9	(0.1)
Other/Unknown	56	(0.6)	39	(2.7)	43	(1.6)	18	(1.9)	3	(0.7)	164	(1.1)
Subtotal	9102	(100)	1463	(100)	2762	(100)	936	(100)	450	(100)	14721	(100)
Female												
Injection drug user (IDU)	148	(55.8)	172	(50.7)	59	(30.4)	15	(20.8)	24	(66.7)	418	(46.0)
Transfusion/Hemophilia	5	(1.9)	2	(0.6)	2	(1.0)	2	(2.8)	0	(0.0)	11	(1.2)
Heterosexual (3)	90	(34.0)	139	(41.0)	105	(54.1)	48	(66.7)	7	(19.4)	389	(42.8)
Perinatal	1	(0.4)	6	(1.8)	9	(4.6)	0	(0.0)	2	(5.6)	18	(2.0)
Other/Unknown	21	(7.9)	20	(5.9)	19	(9.8)	7	(9.7)	3	(8.3)	72	(7.9)
Subtotal	265	(100)	339	(100)	194	(100)	72	(100)	36	(100)	908	(100)
Transgender	72	(100)	123	(100)	129	(100)	39	(100)	19	(100)	382	(100)
Total	9439	(100)	1925	(100)	3085	(100)	1047	(100)	505	(100)	16011	(100)

(1) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.

(2) Total includes persons with unknown race.

(3) Includes persons who have had heterosexual contact with a person with HIV or with a person who is at risk for HIV.