

QUARTERLY HIV/AIDS SURVEILLANCE REPORT

San Francisco Department of Public Health HIV/AIDS Cases Reported Through June 2011

Contents	Page
Surveillance Summary.....	1
Table 1: Adult/Adolescent HIV/AIDS Cases by Transmission Category.....	2
Table 2: AIDS Cases by Gender and Year of AIDS Diagnosis.....	3
Table 3: HIV/AIDS Cases by Gender and Year of Initial HIV Diagnosis.....	3
Table 4: AIDS Cases by Transmission Category and Race/Ethnicity.....	4
Table 5: HIV Non-AIDS Cases by Transmission Category and Race/Ethnicity.....	4
Table 6: AIDS Cases by Transmission Category and Year of AIDS Diagnosis.....	5
Table 7: HIV/AIDS Cases by Transmission Category and Year of Initial HIV Diagnosis.....	6
Table 8: AIDS Cases by Gender, Age Group and Race/Ethnicity.....	7
Table 9: HIV Non-AIDS Cases by Gender, Age Group and Race/Ethnicity.....	8
Table 10: AIDS Cases by Race/Ethnicity and Year of AIDS Diagnosis.....	9
Table 11: HIV/AIDS Cases by Race/Ethnicity and Year of Initial HIV Diagnosis.....	9
Table 12: AIDS Incidence, Mortality, and Prevalence by Year.....	10
Table 13: AIDS Cases by Initial AIDS-Defining Condition and Year of AIDS Diagnosis.....	11
Table 14: Cumulative AIDS Indicator Conditions among Persons with AIDS.....	12
Table 15: Living Adult/Adolescent HIV/AIDS Cases by Transmission Category.....	13
Table 16: Living HIV/AIDS Cases by Transmission Category and Race/Ethnicity.....	13
Table 17: Living HIV/AIDS Cases by Gender, Age Group and Race/Ethnicity.....	14

The HIV/AIDS Surveillance Report is published quarterly by the
San Francisco Department of Public Health, HIV Epidemiology Section
25 Van Ness Avenue, Suite 500, San Francisco, CA 94102; Phone (415) 554-9050, FAX (415) 431-0353
Director of Health: Barbara A. Garcia, MPA;
Health Officer and Director of Population Health and Prevention. Tomas Aragon, MD, DrPH;
Section Director: Susan Scheer, PhD, MPH; Surveillance Unit Director: Ling Hsu, MPH;
Program Coordinators: Maree Kay Parisi, Viva Delgado, MPH;
Epidemiologists: Mia Miao-Jung Chen, PhD, MPH, Alison Hughes, MPH,
Priscilla Lee Chu, DrPH, MPH, Sharon Pipkin, MPH, Annie Vu, MPH, Kara O'Keefe, PhD, MPH
The HIV/AIDS Surveillance Report is accessible via internet:
<http://www.sfdph.org/dph/files/reports/default.asp>

HIV/AIDS Surveillance Summary

San Francisco (as of 06/30/2011)

Cumulative cases ¹ :	
AIDS	28,948
HIV non-AIDS ²	5,985
Cumulative AIDS deaths:	
	19,433
Living HIV/AIDS cases	
	15,430


California³ (as of 12/31/2010)

Cumulative cases:	
AIDS	159,329
HIV non-AIDS	41,892
Cumulative AIDS deaths:	
	88,842
Living HIV/AIDS cases:	
	110,994

United States⁴

Cumulative AIDS cases (as of 12/31/2009):	1,113,971
Cumulative AIDS deaths (as of 12/31/2008):	601,415
Living HIV/AIDS cases (as of 12/31/2008):	670,903

San Francisco AIDS Incidence, Mortality, and Prevalence by Year, 1980-2011⁵



1. Includes SF residents diagnosed in SF and SF residents diagnosed in other jurisdictions.
2. Includes HIV non-AIDS cases reported by name. HIV non-AIDS cases reported by a non-name code between July 2002 and April 2006 whose names have not been ascertained are not included.
3. CA data source: www.cdph.ca.gov/data/statistics/Pages/OAHIVAIDSStatistics.aspx.
4. US data source: CDC HIV Surveillance Report, 2009, vol. 21. www.cdc.gov/hiv/topics/surveillance/resources/reports/.
5. Reporting for recent year is incomplete. See Table 12 for actual numbers per year.

**HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 06/30/2011**

Table 1. Adult/Adolescent HIV/AIDS Cases (>12 years) by Transmission Category, San Francisco, 1980-2011

Transmission Category	AIDS		HIV non-AIDS#	
	No.	(%)	No.	(%)
Gay or bisexual male	21147	(73.1)	4323	(72.4)
Heterosexual male injection drug user	1408	(4.9)	220	(3.7)
Heterosexual female injection drug user	716	(2.5)	158	(2.6)
Gay or bisexual male injection drug user	4273	(14.8)	696	(11.7)
Lesbian or bisexual injection drug user	60	(0.2)	15	(0.3)
Transgender (1)	424	(1.5)	146	(2.4)
Hemophiliac	16	(0.1)	2	(0.0)
Heterosexual contact male (2)	159	(0.5)	51	(0.9)
Heterosexual contact female (2)	318	(1.1)	131	(2.2)
Transfusion recipient	143	(0.5)	1	(0.0)
Risk not reported/Other (3)	246	(0.9)	228	(3.8)
Total	28910	(100)	5971	(100)

* Residents of San Francisco at time of HIV/AIDS diagnosis.

Includes HIV non-AIDS cases reported by name. Excludes HIV non-AIDS cases reported by a non-name code between July 2002 and April 2006 whose names have not been ascertained.

- (1) Transgender information was collected since September 1996. Data prior to this are incomplete.
- (2) Includes persons who have had heterosexual contact with a person with HIV/AIDS or with a person who is at risk for HIV.
- (3) Includes persons for whom risk information is incomplete (due to death, refusal to be interviewed or loss to follow-up), cases still under investigation, or interviewed patients who offered no plausible risk for HIV.

**HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 06/30/2011**

Table 2. AIDS Cases by Gender and Year of AIDS Diagnosis, San Francisco, 1980-2011

Gender	Year of AIDS Diagnosis					
	< 2001 No. (%)	2001 No. (%)	2002 No. (%)	2003 No. (%)	2004 No. (%)	2005 No. (%)
Male	23254 (95.3)	454 (88.3)	443 (89.1)	499 (88.6)	423 (88.3)	423 (89.1)
Female	875 (3.6)	45 (8.8)	36 (7.2)	40 (7.1)	39 (8.1)	37 (7.8)
Transgender (1)	275 (1.1)	15 (2.9)	18 (3.6)	24 (4.3)	17 (3.5)	15 (3.2)
Total	24404 (100)	514 (100)	497 (100)	563 (100)	479 (100)	475 (100)

Gender	Year of AIDS Diagnosis					
	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)
Male	401 (90.1)	403 (90.2)	370 (88.1)	285 (89.6)	252 (90.0)	94 (88.7)
Female	30 (6.7)	32 (7.2)	40 (9.5)	24 (7.5)	15 (5.4)	10 (9.4)
Transgender (1)	14 (3.1)	12 (2.7)	10 (2.4)	9 (2.8)	13 (4.6)	2 (1.9)
Total	445 (100)	447 (100)	420 (100)	318 (100)	280 (100)	106 (100)

Table 3. HIV/AIDS Cases by Gender and Year of Initial HIV Diagnosis#, San Francisco, 2006-2011

Gender	Year of Initial HIV Diagnosis					
	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)
Male	467 (91.4)	461 (87.8)	446 (89.6)	418 (89.9)	383 (90.1)	143 (88.8)
Female	35 (6.8)	42 (8.0)	40 (8.0)	28 (6.0)	31 (7.3)	16 (9.9)
Transgender (1)	9 (1.8)	22 (4.2)	12 (2.4)	19 (4.1)	11 (2.6)	2 (1.2)
Total	511 (100)	525 (100)	498 (100)	465 (100)	425 (100)	161 (100)

* Residents of San Francisco at time of HIV/AIDS diagnosis.

Includes persons with a diagnosis of HIV (not AIDS), an initial diagnosis of HIV (not AIDS) and later diagnosed with AIDS, and concurrent diagnosis of HIV and AIDS. The initial year of HIV diagnosis was determined based on the earliest date of HIV antibody test, viral load or CD4 test, initiation of antiretroviral therapy, or patient self-report of a positive HIV test.

(1) Transgender information was collected since September 1996. Data prior to this are incomplete.

HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 06/30/2011

Table 4. AIDS Cases by Transmission Category and Race/Ethnicity, San Francisco San Francisco, 1980-2011

Transmission Category (1)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander No. (%)	Native American No. (%)
Adult/Adolescent					
Gay or bisexual male	16222 (79.6)	1567 (42.6)	2616 (73.2)	708 (76.8)	55 (43.7)
Injection drug user (IDU)	738 (3.6)	1047 (28.5)	251 (7.0)	40 (4.3)	19 (15.1)
Gay or bisexual male IDU	3105 (15.2)	728 (19.8)	485 (13.6)	77 (8.4)	45 (35.7)
Lesbian or bisexual IDU	26 (0.1)	22 (0.6)	7 (0.2)	2 (0.2)	2 (1.6)
Hemophiliac	8 (0.0)	2 (0.1)	5 (0.1)	1 (0.1)	0 (0.0)
Heterosexual (2)	120 (0.6)	195 (5.3)	114 (3.2)	46 (5.0)	4 (3.2)
Transfusion recipient	68 (0.3)	23 (0.6)	27 (0.8)	19 (2.1)	0 (0.0)
Risk not reported/Other (3)	82 (0.4)	78 (2.1)	58 (1.6)	24 (2.6)	1 (0.8)
Pediatric (0-12 years) (4)	8 (0.0)	13 (0.4)	10 (0.3)	5 (0.5)	0 (0.0)
Total	20377 (100)	3675 (100)	3573 (100)	922 (100)	126 (100)

Table 5. HIV Non-AIDS Cases# by Transmission Category and Race/Ethnicity, San Francisco

Transmission Category (1)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander No. (%)	Native American No. (%)
Adult/Adolescent					
Gay or bisexual male	2943 (78.7)	367 (46.3)	729 (75.9)	256 (80.5)	19 (57.6)
Injection drug user (IDU)	154 (4.1)	170 (21.4)	39 (4.1)	4 (1.3)	5 (15.2)
Gay or bisexual male IDU	499 (13.3)	103 (13.0)	97 (10.1)	26 (8.2)	9 (27.3)
Lesbian or bisexual IDU	5 (0.1)	5 (0.6)	4 (0.4)	0 (0.0)	0 (0.0)
Hemophiliac	2 (0.1)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (2)	34 (0.9)	88 (11.1)	40 (4.2)	18 (5.7)	0 (0.0)
Transfusion recipient	0 (0.0)	1 (0.1)	0 (0.0)	0 (0.0)	0 (0.0)
Risk not reported/Other (3)	101 (2.7)	56 (7.1)	45 (4.7)	13 (4.1)	0 (0.0)
Pediatric (0-12 years) (4)	2 (0.1)	3 (0.4)	6 (0.6)	1 (0.3)	0 (0.0)
Total	3740 (100)	793 (100)	960 (100)	318 (100)	33 (100)

* Residents of San Francisco at time of HIV/AIDS diagnosis.

Includes HIV non-AIDS cases reported by name. Excludes HIV non-AIDS cases reported by a non-name code between July 2002 and April 2006 whose names have not been ascertained.

- (1) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
- (2) Includes persons who have had heterosexual contact with a person with HIV/AIDS or with a person who is at risk for HIV.
- (3) Includes persons for whom risk information is incomplete (due to death, refusal to be interviewed or loss to follow-up), cases still under investigation, or interviewed patients who offered no plausible risk for HIV.
- (4) Includes children who have hemophilia or other coagulation disorder, have received a blood transfusion, or who have acquired their infection from an infected mother during the perinatal period.

HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 06/30/2011

Table 6. AIDS Cases by Transmission Category and Year of AIDS Diagnosis, San Francisco, 1980-2011

Transmission Category (1)	Year of AIDS Diagnosis					
	< 2001 No. (%)	2001 No. (%)	2002 No. (%)	2003 No. (%)	2004 No. (%)	2005 No. (%)
Adult/Adolescent						
Gay or bisexual male	18485 (75.7)	310 (60.3)	304 (61.2)	362 (64.3)	316 (66.0)	293 (61.7)
Injection drug user (IDU)	1631 (6.7)	66 (12.8)	63 (12.7)	75 (13.3)	51 (10.6)	55 (11.6)
Gay or bisexual male IDU	3653 (15.0)	96 (18.7)	103 (20.7)	92 (16.3)	86 (18.0)	93 (19.6)
Lesbian or bisexual IDU	41 (0.2)	3 (0.6)	2 (0.4)	3 (0.5)	1 (0.2)	3 (0.6)
Hemophiliac	15 (0.1)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (2)	276 (1.1)	19 (3.7)	16 (3.2)	21 (3.7)	14 (2.9)	19 (4.0)
Transfusion recipient	141 (0.6)	1 (0.2)	1 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)
Risk not reported/Other (3)	126 (0.5)	19 (3.7)	7 (1.4)	9 (1.6)	11 (2.3)	12 (2.5)
Pediatric (0-12 years) (4)	36 (0.1)	0 (0.0)	1 (0.2)	1 (0.2)	0 (0.0)	0 (0.0)
Total	24404 (100)	514 (100)	497 (100)	563 (100)	479 (100)	475 (100)

Transmission Category (1)	Year of AIDS Diagnosis					
	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)
Adult/Adolescent						
Gay or bisexual male	288 (64.7)	290 (64.9)	263 (62.6)	206 (64.8)	160 (57.1)	55 (51.9)
Injection drug user (IDU)	40 (9.0)	42 (9.4)	31 (7.4)	35 (11.0)	24 (8.6)	15 (14.2)
Gay or bisexual male IDU	86 (19.3)	75 (16.8)	80 (19.0)	45 (14.2)	69 (24.6)	25 (23.6)
Lesbian or bisexual IDU	2 (0.4)	0 (0.0)	3 (0.7)	2 (0.6)	0 (0.0)	0 (0.0)
Hemophiliac	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.3)	0 (0.0)	0 (0.0)
Heterosexual (2)	20 (4.5)	32 (7.2)	27 (6.4)	17 (5.3)	15 (5.4)	5 (4.7)
Transfusion recipient	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Risk not reported/Other (3)	9 (2.0)	8 (1.8)	16 (3.8)	12 (3.8)	12 (4.3)	6 (5.7)
Pediatric (0-12 years) (4)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Total	445 (100)	447 (100)	420 (100)	318 (100)	280 (100)	106 (100)

* Residents of San Francisco at time of HIV/AIDS diagnosis.

- (1) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
- (2) Includes persons who have had heterosexual contact with a person with HIV/AIDS or with a person who is at risk for HIV.
- (3) Includes persons for whom risk information is incomplete (due to death, refusal to be interviewed or loss to follow-up), cases still under investigation, or interviewed patients who offered no plausible risk for HIV.
- (4) Includes children who have hemophilia or other coagulation disorder, have received a blood transfusion, or who have acquired their infection from an infected mother during the perinatal period.

**HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 06/30/2011**

Table 7. HIV/AIDS Cases by Transmission Category and Year of Initial HIV Diagnosis#, San Francisco, 2006-2011

Transmission Category (1)	Year of Initial HIV Diagnosis					
	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)
Adult/Adolescent						
Gay or bisexual male	360 (70.5)	345 (65.7)	357 (71.7)	336 (72.3)	272 (64.0)	113 (70.2)
Injection drug user (IDU)	40 (7.8)	36 (6.9)	27 (5.4)	22 (4.7)	31 (7.3)	9 (5.6)
Gay or bisexual male IDU	68 (13.3)	74 (14.1)	49 (9.8)	59 (12.7)	58 (13.6)	18 (11.2)
Lesbian or bisexual IDU	1 (0.2)	1 (0.2)	3 (0.6)	3 (0.6)	0 (0.0)	0 (0.0)
Hemophiliac	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (2)	23 (4.5)	42 (8.0)	32 (6.4)	19 (4.1)	31 (7.3)	9 (5.6)
Transfusion recipient	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Risk not reported/Other (3)	19 (3.7)	27 (5.1)	30 (6.0)	26 (5.6)	33 (7.8)	12 (7.5)
Pediatric (0-12 years) (4)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Total	511 (100)	525 (100)	498 (100)	465 (100)	425 (100)	161 (100)

* Residents of San Francisco at time of HIV/AIDS diagnosis.

Includes persons with a diagnosis of HIV (not AIDS), an initial diagnosis of HIV (not AIDS) and later diagnosed with AIDS, and concurrent diagnosis of HIV and AIDS. The initial year of HIV diagnosis was determined based on the earliest date of HIV antibody test, viral load or CD4 test, initiation of antiretroviral therapy, or patient self-report of a positive HIV test.

- (1) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
- (2) Includes persons who have had heterosexual contact with a person with HIV/AIDS or with a person who is at risk for HIV.
- (3) Includes persons for whom risk information is incomplete (due to death, refusal to be interviewed or loss to follow-up), cases still under investigation, or interviewed patients who offered no plausible risk for HIV.
- (4) Includes children who have hemophilia or other coagulation disorder, have received a blood transfusion, or who have acquired their infection from an infected mother during the perinatal period.

HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 06/30/2011

Table 8. AIDS Cases by Gender, Age Group and Race/Ethnicity, San Francisco, 1980-2011

Male Age at AIDS Diagnosis (Years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander No. (%)	Native American No. (%)	Total (1) No. (%)
0 - 12	4 (0.0)	4 (0.1)	5 (0.2)	4 (0.5)	0 (0.0)	19 (0.1)
13 - 19	12 (0.1)	1 (0.0)	14 (0.4)	1 (0.1)	2 (1.8)	30 (0.1)
20 - 24	310 (1.6)	71 (2.4)	124 (3.8)	26 (3.2)	5 (4.6)	540 (2.0)
25 - 29	1665 (8.4)	251 (8.4)	464 (14.2)	88 (10.8)	21 (19.3)	2519 (9.2)
30 - 39	8840 (44.5)	1168 (39.1)	1538 (47.0)	361 (44.1)	51 (46.8)	12072 (44.2)
40 - 49	6537 (32.9)	998 (33.4)	820 (25.1)	252 (30.8)	25 (22.9)	8695 (31.8)
50 - 59	2007 (10.1)	385 (12.9)	247 (7.5)	64 (7.8)	4 (3.7)	2732 (10.0)
60 +	494 (2.5)	111 (3.7)	61 (1.9)	22 (2.7)	1 (0.9)	694 (2.5)
Male subtotal	19869 (100)	2989 (100)	3273 (100)	818 (100)	109 (100)	27301 (100)
Female Age at AIDS Diagnosis (Years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander No. (%)	Native American No. (%)	Total (1) No. (%)
0 - 12	4 (1.0)	9 (1.6)	5 (2.8)	1 (1.5)	0 (0.0)	19 (1.6)
13 - 19	1 (0.3)	2 (0.4)	2 (1.1)	0 (0.0)	0 (0.0)	5 (0.4)
20 - 24	15 (3.8)	10 (1.8)	10 (5.6)	4 (6.2)	1 (7.1)	41 (3.4)
25 - 29	40 (10.2)	46 (8.3)	26 (14.7)	11 (16.9)	1 (7.1)	126 (10.3)
30 - 39	154 (39.1)	204 (37.0)	60 (33.9)	25 (38.5)	8 (57.1)	458 (37.4)
40 - 49	107 (27.2)	189 (34.3)	44 (24.9)	17 (26.2)	4 (28.6)	366 (29.9)
50 - 59	42 (10.7)	68 (12.3)	17 (9.6)	4 (6.2)	0 (0.0)	135 (11.0)
60 +	31 (7.9)	23 (4.2)	13 (7.3)	3 (4.6)	0 (0.0)	73 (6.0)
Female subtotal	394 (100)	551 (100)	177 (100)	65 (100)	14 (100)	1223 (100)
Transgender (2) Age at AIDS Diagnosis (Years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander Native American No. (%)	Total (1) No. (%)	
13 - 29	23 (20.2)	28 (20.7)	32 (26.0)	13 (31.0)	98 (23.1)	
30 - 39	55 (48.2)	49 (36.3)	57 (46.3)	19 (45.2)	185 (43.6)	
40 +	36 (31.6)	58 (43.0)	34 (27.6)	10 (23.8)	141 (33.3)	
Transgender subtotal	114 (100)	135 (100)	123 (100)	42 (100)	424 (100)	

* Residents of San Francisco at time of HIV/AIDS diagnosis.

(1) Total includes persons with multiple or unknown race.

(2) Transgender information was collected since September 1996. Certain age or race/ethnic groups are combined for transgender cases because of small number.

**HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 06/30/2011**

Table 9. HIV Non-AIDS Cases# by Gender, Age Group and Race/Ethnicity, San Francisco

Male Age at Initial HIV Diagnosis (Years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander No. (%)	Native American No. (%)	Total (1) No. (%)
0 - 12	1 (0.0)	2 (0.3)	1 (0.1)	1 (0.4)	0 (0.0)	5 (0.1)
13 - 19	30 (0.8)	21 (3.6)	22 (2.6)	7 (2.5)	0 (0.0)	82 (1.5)
20 - 24	307 (8.5)	55 (9.4)	108 (12.8)	33 (11.7)	5 (17.9)	523 (9.6)
25 - 29	563 (15.6)	83 (14.1)	187 (22.2)	63 (22.3)	5 (17.9)	928 (17.0)
30 - 39	1578 (43.8)	203 (34.6)	361 (42.9)	132 (46.8)	14 (50.0)	2335 (42.7)
40 - 49	833 (23.1)	159 (27.1)	138 (16.4)	40 (14.2)	4 (14.3)	1200 (22.0)
50 - 59	237 (6.6)	49 (8.3)	21 (2.5)	6 (2.1)	0 (0.0)	315 (5.8)
60 +	54 (1.5)	15 (2.6)	4 (0.5)	0 (0.0)	0 (0.0)	74 (1.4)
Male subtotal	3603 (100)	587 (100)	842 (100)	282 (100)	28 (100)	5462 (100)
Female Age at Initial HIV Diagnosis (Years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander No. (%)	Native American No. (%)	Total (1) No. (%)
0 - 12	1 (0.9)	1 (0.6)	5 (7.0)	0 (0.0)	0 (0.0)	9 (2.4)
13 - 19	2 (1.9)	5 (3.2)	2 (2.8)	0 (0.0)	0 (0.0)	9 (2.4)
20 - 24	12 (11.2)	13 (8.4)	13 (18.3)	1 (4.0)	0 (0.0)	41 (10.9)
25 - 29	22 (20.6)	14 (9.0)	13 (18.3)	6 (24.0)	0 (0.0)	56 (14.9)
30 - 39	32 (29.9)	49 (31.6)	23 (32.4)	7 (28.0)	3 (75.0)	120 (31.8)
40 - 49	23 (21.5)	52 (33.5)	11 (15.5)	7 (28.0)	0 (0.0)	95 (25.2)
50 - 59	15 (14.0)	18 (11.6)	4 (5.6)	2 (8.0)	1 (25.0)	42 (11.1)
60 +	0 (0.0)	3 (1.9)	0 (0.0)	2 (8.0)	0 (0.0)	5 (1.3)
Female subtotal	107 (100)	155 (100)	71 (100)	25 (100)	4 (100)	377 (100)
Transgender (2) Age at Initial HIV Diagnosis (Years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander Native American No. (%)	Total (1) No. (%)	
13 - 29	13 (43.3)	25 (49.0)	23 (48.9)	4 (33.3)	68 (46.6)	
30 +	17 (56.7)	26 (51.0)	24 (51.1)	8 (66.7)	78 (53.4)	
Transgender subtotal	30 (100)	51 (100)	47 (100)	12 (100)	146 (100)	

* Residents of San Francisco at time of HIV/AIDS diagnosis.

Includes HIV non-AIDS cases reported by name. Excludes HIV non-AIDS cases reported by a non-name code between July 2002 and April 2006 whose names have not been ascertained.

(1) Total includes persons with multiple or unknown race.

(2) Transgender information was collected since September 1996. Certain age or race/ethnic groups are combined for transgender cases because of small number.

**HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 06/30/2011**

Table 10. AIDS Cases by Race/Ethnicity and Year of AIDS Diagnosis, San Francisco, 1980-2011

Race/Ethnicity	Year of AIDS Diagnosis					
	< 2001 No. (%)	2001 No. (%)	2002 No. (%)	2003 No. (%)	2004 No. (%)	2005 No. (%)
White	17880 (73.3)	295 (57.4)	293 (59.0)	285 (50.6)	260 (54.3)	273 (57.5)
African American	2849 (11.7)	98 (19.1)	89 (17.9)	105 (18.7)	78 (16.3)	84 (17.7)
Latino	2745 (11.2)	74 (14.4)	76 (15.3)	126 (22.4)	110 (23.0)	86 (18.1)
Asian/Pacific Islander	642 (2.6)	34 (6.6)	30 (6.0)	34 (6.0)	20 (4.2)	24 (5.1)
Native American	92 (0.4)	5 (1.0)	2 (0.4)	6 (1.1)	4 (0.8)	3 (0.6)
Total (1)	24404 (100)	514 (100)	497 (100)	563 (100)	479 (100)	475 (100)

Race/Ethnicity	Year of AIDS Diagnosis					
	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)
White	247 (55.5)	260 (58.2)	218 (51.9)	166 (52.2)	137 (48.9)	63 (59.4)
African American	82 (18.4)	78 (17.4)	80 (19.0)	61 (19.2)	55 (19.6)	16 (15.1)
Latino	81 (18.2)	73 (16.3)	70 (16.7)	62 (19.5)	54 (19.3)	16 (15.1)
Asian/Pacific Islander	22 (4.9)	28 (6.3)	41 (9.8)	14 (4.4)	25 (8.9)	8 (7.5)
Native American	6 (1.3)	0 (0.0)	3 (0.7)	3 (0.9)	2 (0.7)	0 (0.0)
Total (1)	445 (100)	447 (100)	420 (100)	318 (100)	280 (100)	106 (100)

Table 11. HIV/AIDS Cases by Race/Ethnicity and Year of Initial HIV Diagnosis#, San Francisco, 2006-2011

Race/Ethnicity	Year of Initial HIV Diagnosis					
	2006 No. (%)	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)
White	289 (56.6)	276 (52.6)	248 (49.8)	236 (50.8)	215 (50.6)	79 (49.1)
African American	76 (14.9)	80 (15.2)	80 (16.1)	74 (15.9)	61 (14.4)	23 (14.3)
Latino	103 (20.2)	103 (19.6)	112 (22.5)	96 (20.6)	89 (20.9)	38 (23.6)
Asian/Pacific Islander	32 (6.3)	47 (9.0)	42 (8.4)	41 (8.8)	43 (10.1)	19 (11.8)
Native American	4 (0.8)	0 (0.0)	4 (0.8)	2 (0.4)	3 (0.7)	0 (0.0)
Total (1)	511 (100)	525 (100)	498 (100)	465 (100)	425 (100)	161 (100)

* Residents of San Francisco at time of HIV/AIDS diagnosis.

Includes persons with a diagnosis of HIV (not AIDS), an initial diagnosis of HIV (not AIDS) and later diagnosed with AIDS, and concurrent diagnosis of HIV and AIDS. The initial year of HIV diagnosis was determined based on the earliest date of HIV antibody test, viral load or CD4 test, initiation of antiretroviral therapy, or patient self-report of a positive HIV test.

(1) Total includes persons with multiple or unknown race.

**HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 06/30/2011**

Table 12. AIDS Incidence, Mortality, and Prevalence by Year, San Francisco, 1980-2011

Year	Number of AIDS Cases Reported per Year	Number of AIDS Cases Diagnosed per Year (1)	Number of AIDS Deaths Occurred per Year (1)	Number of Persons Living with AIDS (1)
1980	0	3	0	3
1981	21	26	8	21
1982	75	99	32	88
1983	197	274	111	251
1984	451	557	273	535
1985	673	859	534	860
1986	981	1236	807	1289
1987	1287	1629	878	2040
1988	1408	1763	1039	2764
1989	1584	2161	1276	3649
1990	1686	2046	1366	4329
1991	1685	2285	1508	5106
1992	1638	2329	1641	5794
1993	4271	2073	1600	6267
1994	1915	1788	1595	6460
1995	1633	1563	1483	6540
1996	1241	1080	992	6628
1997	1060	805	422	7011
1998	794	695	402	7304
1999	723	576	356	7524
2000	625	557	349	7732
2001	496	514	324	7922
2002	440	497	321	8098
2003	533	563	303	8358
2004	557	479	307	8530
2005	499	475	313	8692
2006	423	445	288	8849
2007	537	447	269	9027
2008	558	420	231	9216
2009	355	318	206	9328
2010	408	280	140	9468
2011	194	106	59	9515
Total	28948	28948	19433	

* Residents of San Francisco at time of HIV/AIDS diagnosis.

(1) Data in recent years is incomplete due to delay in cases/deaths reporting.

HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 06/30/2011

Table 13. AIDS Cases by Initial AIDS-Defining Condition and Year of AIDS Diagnosis, San Francisco, 1980-2011

Initial AIDS-Defining Condition	Year of AIDS Diagnosis		
	1980-1989 No. (%)	1990-1995 No. (%)	1996-2011 No. (%)
Bacterial infections, recurrent, <13 years [HIV+(1)]	1 (0.0)	4 (0.0)	3 (0.0)
Candidiasis of bronchi, trachea, or lungs	15 (0.2)	6 (0.0)	4 (0.0)
Candidiasis of esophagus	221 (2.6)	217 (1.8)	100 (1.2)
Cervical cancer, invasive [HIV+]	1 (0.0)	1 (0.0)	4 (0.0)
Coccidioidomycosis, disseminated or extrapulmonary [HIV+]	4 (0.0)	2 (0.0)	1 (0.0)
Cryptococcosis, extrapulmonary	197 (2.3)	130 (1.1)	53 (0.6)
Cryptosporidiosis, intestinal (>1 mo. duration)	105 (1.2)	176 (1.5)	54 (0.7)
Cytomegalovirus (except liver, spleen, lymph nodes), >1 month of age	101 (1.2)	93 (0.8)	11 (0.1)
CMV retinitis with loss of vision [HIV+]	34 (0.4)	67 (0.6)	12 (0.1)
HIV encephalopathy [HIV+]	181 (2.1)	163 (1.3)	53 (0.6)
Herpes simplex: chronic (>1 mo.), bronchitis, pneumonitis, esophagitis	49 (0.6)	39 (0.3)	13 (0.2)
Histoplasmosis, disseminated or extrapulmonary [HIV+]	10 (0.1)	8 (0.1)	8 (0.1)
Isosporiasis, intestinal (>1 mo. duration) [HIV+]	8 (0.1)	8 (0.1)	5 (0.1)
Kaposi's sarcoma (2)	1852 (21.5)	767 (6.3)	238 (2.9)
Lymphoid interstitial pneumonia/ pulmonary lymphoid hyperplasia, <13 years	3 (0.0)	1 (0.0)	0 (0.0)
Lymphoma, Burkitt's (non-Hodgkin's) [HIV+]	55 (0.6)	27 (0.2)	30 (0.4)
Lymphoma, immunoblastic (non-Hodgkin's) [HIV+]	200 (2.3)	153 (1.3)	68 (0.8)
Lymphoma, primary in brain (2)	19 (0.2)	12 (0.1)	10 (0.1)
Mycobacterium avium complex or M. kansasii, disseminated or extrapulmonary	139 (1.6)	103 (0.9)	18 (0.2)
Mycobacterium tuberculosis, pulmonary [HIV+]	43 (0.5)	117 (1.0)	59 (0.7)
Mycobacterium tuberculosis, disseminated or extrapulmonary [HIV+]	62 (0.7)	62 (0.5)	26 (0.3)
Mycobacterium other species, disseminated or extrapulmonary [HIV+]	12 (0.1)	14 (0.1)	1 (0.0)
Pneumocystis carinii pneumonia	3611 (42.0)	1587 (13.1)	514 (6.2)
Pneumonia, recurrent [HIV+]	33 (0.4)	123 (1.0)	96 (1.2)
Progressive multifocal leukoencephalopathy	37 (0.4)	8 (0.1)	5 (0.1)
Salmonella sepsis, recurrent [HIV+]	6 (0.1)	1 (0.0)	1 (0.0)
Toxoplasmosis of brain, >1 month of age	147 (1.7)	74 (0.6)	22 (0.3)
Wasting syndrome [HIV+]	223 (2.6)	333 (2.8)	132 (1.6)
CD4 T lymphocyte count <200 or percent <14 [HIV+]	1235 (14.3)	7787 (64.4)	6716 (81.3)
Any AIDS indicator condition, HIV-negative and CD4 count <400 (3)	3 (0.0)	1 (0.0)	0 (0.0)
Total	8607 (100)	12084 (100)	8257 (100)

* Residents of San Francisco at time of HIV/AIDS diagnosis.

(1) [HIV+]: Indicator conditions that require laboratory evidence of HIV infection.

(2) Laboratory evidence of HIV infection in persons > 60 years of age.

(3) In the absence of other causes of immunocompromise.

**HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 06/30/2011**

Table 14. Cumulative AIDS Indicator Conditions among Persons with AIDS, San Francisco, 1980-2011

AIDS Indicator Condition (1)	Total No. (%)
Bacterial infections, recurrent, <13 years [HIV+(2)]	11 (0.0)
Candidiasis of bronchi, trachea, or lungs	145 (0.5)
Candidiasis of esophagus	2632 (9.1)
Cervical cancer, invasive [HIV+]	13 (0.0)
Coccidioidomycosis, disseminated or extrapulmonary [HIV+]	62 (0.2)
Cryptococcosis, extrapulmonary	1876 (6.5)
Cryptosporidiosis, intestinal (>1 mo. duration)	1381 (4.8)
Cytomegalovirus (except liver, spleen, lymph nodes), >1 month of age	2469 (8.5)
CMV retinitis with loss of vision [HIV+]	2493 (8.6)
HIV encephalopathy [HIV+]	2496 (8.6)
Herpes simplex: chronic (>1 mo.), bronchitis, pneumonitis, esophagitis	442 (1.5)
Histoplasmosis, disseminated or extrapulmonary [HIV+]	159 (0.5)
Isosporiasis, intestinal (>1 mo. duration) [HIV+]	68 (0.2)
Kaposi's sarcoma (3)	6747 (23.3)
Lymphoid interstitial pneumonia/pulmonary lymphoid hyperplasia, <13 years	6 (0.0)
Lymphoma, Burkitt's (non-Hodgkin's) [HIV+]	555 (1.9)
Lymphoma, immunoblastic (non-Hodgkin's) [HIV+]	1135 (3.9)
Lymphoma, primary in brain (3)	392 (1.4)
Mycobacterium avium complex or M. kansasii, disseminated or extrapulmonary	5097 (17.6)
Mycobacterium tuberculosis, pulmonary [HIV+]	658 (2.3)
Mycobacterium tuberculosis, disseminated or extrapulmonary [HIV+]	480 (1.7)
Mycobacterium other species, disseminated or extrapulmonary [HIV+]	336 (1.2)
Pneumocystis carinii pneumonia	11048 (38.2)
Pneumonia, recurrent [HIV+]	1026 (3.5)
Progressive multifocal leukoencephalopathy	311 (1.1)
Salmonella sepsis, recurrent [HIV+]	58 (0.2)
Toxoplasmosis of brain, >1 month of age	1163 (4.0)
Wasting syndrome [HIV+]	4354 (15.0)

* Residents of San Francisco at time of HIV/AIDS diagnosis.

(1) Cases may have more than one condition.

(2) [HIV+]: Indicator conditions that require laboratory evidence of HIV infection.

(3) Laboratory evidence of HIV infection in persons > 60 years of age.

**HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 06/30/2011**

Table 15. Living Adult/Adolescent HIV/AIDS Cases (>12 years) by Transmission Category, San Francisco

Transmission Category	No.	(%)
Gay or bisexual male	11116	(72.2)
Heterosexual male injection drug user	606	(3.9)
Heterosexual female injection drug user	418	(2.7)
Gay or bisexual male injection drug user	2063	(13.4)
Lesbian or bisexual injection drug user	41	(0.3)
Transgender (1)	340	(2.2)
Hemophiliac	5	(0.0)
Heterosexual contact male (2)	150	(1.0)
Heterosexual contact female (2)	308	(2.0)
Transfusion recipient	17	(0.1)
Risk not reported/Other (3)	329	(2.1)
Total	15393	(100)

Table 16. Living HIV/AIDS Cases by Transmission Category and Race/Ethnicity, San Francisco

Transmission Category	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander No. (%)	Native American No. (%)
Adult/Adolescent					
Gay or bisexual male	7659 (78.8)	910 (44.2)	1910 (75.3)	626 (77.8)	43 (50.0)
Injection drug user (IDU)	379 (3.9)	473 (23.0)	119 (4.7)	24 (3.0)	16 (18.6)
Gay or bisexual male IDU	1422 (14.6)	366 (17.8)	294 (11.6)	74 (9.2)	22 (25.6)
Lesbian or bisexual IDU	16 (0.2)	17 (0.8)	5 (0.2)	1 (0.1)	2 (2.3)
Hemophiliac	3 (0.0)	2 (0.1)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (2)	92 (0.9)	193 (9.4)	122 (4.8)	47 (5.8)	3 (3.5)
Transfusion recipient	6 (0.1)	3 (0.1)	5 (0.2)	3 (0.4)	0 (0.0)
Risk not reported/Other (3)	137 (1.4)	84 (4.1)	70 (2.8)	24 (3.0)	0 (0.0)
Pediatric (0-12 years) (4)	5 (0.1)	9 (0.4)	13 (0.5)	6 (0.7)	0 (0.0)
Total	9719 (100)	2057 (100)	2538 (100)	805 (100)	86 (100)

* Residents of San Francisco at time of HIV/AIDS diagnosis.

- (1) Transgender information was collected since September 1996. Data prior to this are incomplete.
- (2) Includes persons who have had heterosexual contact with a person with HIV/AIDS or with a person who is at risk for HIV.
- (3) Includes persons for whom risk information is incomplete (due to death, refusal to be interviewed or loss to follow-up), cases still under investigation, or interviewed patients who offered no plausible risk for HIV.
- (4) Includes children who have hemophilia or other coagulation disorder, have received a blood transfusion, or who have acquired their infection from an infected mother during the perinatal period.

HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 06/30/2011

Table 17. Living HIV/AIDS Cases by Gender, Age Group and Race/Ethnicity, San Francisco

Male Current Age (Years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander No. (%)	Native American No. (%)	Total (1) No. (%)
0 - 12	0 (0.0)	0 (0.0)	1 (0.0)	0 (0.0)	0 (0.0)	2 (0.0)
13 - 19	2 (0.0)	4 (0.3)	1 (0.0)	0 (0.0)	0 (0.0)	7 (0.0)
20 - 24	42 (0.4)	27 (1.7)	38 (1.7)	13 (1.8)	0 (0.0)	125 (0.9)
25 - 29	169 (1.8)	45 (2.9)	98 (4.3)	33 (4.7)	2 (2.8)	358 (2.5)
30 - 39	922 (9.8)	166 (10.6)	455 (20.0)	165 (23.3)	14 (19.4)	1781 (12.5)
40 - 49	3338 (35.6)	520 (33.1)	931 (41.0)	288 (40.6)	37 (51.4)	5189 (36.5)
50 - 59	3305 (35.2)	571 (36.4)	561 (24.7)	147 (20.7)	18 (25.0)	4637 (32.6)
60 +	1609 (17.1)	237 (15.1)	186 (8.2)	63 (8.9)	1 (1.4)	2109 (14.8)
Male subtotal	9387 (100)	1570 (100)	2271 (100)	709 (100)	72 (100)	14208 (100)
Female Current Age (Years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander No. (%)	Native American No. (%)	Total (1) No. (%)
0 - 12	0 (0.0)	1 (0.3)	2 (1.2)	0 (0.0)	0 (0.0)	3 (0.3)
13 - 19	1 (0.4)	1 (0.3)	6 (3.7)	1 (1.7)	0 (0.0)	11 (1.2)
20 - 24	2 (0.8)	6 (1.6)	3 (1.9)	0 (0.0)	0 (0.0)	11 (1.2)
25 - 29	6 (2.3)	15 (4.1)	15 (9.3)	4 (6.7)	0 (0.0)	41 (4.6)
30 - 39	43 (16.6)	42 (11.4)	24 (14.8)	12 (20.0)	2 (16.7)	129 (14.6)
40 - 49	106 (40.9)	108 (29.3)	52 (32.1)	21 (35.0)	4 (33.3)	296 (33.6)
50 - 59	76 (29.3)	154 (41.7)	40 (24.7)	17 (28.3)	6 (50.0)	299 (33.9)
60 +	25 (9.7)	42 (11.4)	20 (12.3)	5 (8.3)	0 (0.0)	92 (10.4)
Female subtotal	259 (100)	369 (100)	162 (100)	60 (100)	12 (100)	882 (100)
Transgender (2) Current Age (Years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander Native American No. (%)	Total (1) No. (%)	
13 - 39	19 (26.0)	36 (30.5)	47 (44.8)	12 (31.6)	119 (35.0)	
40 +	54 (74.0)	82 (69.5)	58 (55.2)	26 (68.4)	221 (65.0)	
Transgender subtotal	73 (100)	118 (100)	105 (100)	38 (100)	340 (100)	

* Residents of San Francisco at time of HIV/AIDS diagnosis.

(1) Total includes persons with multiple or unknown race.

(2) Transgender information was collected since September 1996. Certain age or race/ethnic groups are combined for transgender cases because of small number.