

QUARTERLY HIV/AIDS SURVEILLANCE REPORT

San Francisco Department of Public Health

HIV/AIDS Cases Reported Through September 2012

| Contents | Page |
|---|------|
| Surveillance Summary..... | 1 |
| Table 1: Adult/Adolescent HIV/AIDS Cases by Transmission Category..... | 2 |
| Table 2: AIDS Cases by Gender and Year of AIDS Diagnosis..... | 3 |
| Table 3: HIV/AIDS Cases by Gender and Year of Initial HIV Diagnosis..... | 3 |
| Table 4: AIDS Cases by Transmission Category and Race/Ethnicity..... | 4 |
| Table 5: HIV Non-AIDS Cases by Transmission Category and Race/Ethnicity..... | 4 |
| Table 6: AIDS Cases by Transmission Category and Year of AIDS Diagnosis..... | 5 |
| Table 7: HIV/AIDS Cases by Transmission Category and Year of Initial HIV Diagnosis..... | 6 |
| Table 8: AIDS Cases by Gender, Age Group and Race/Ethnicity..... | 7 |
| Table 9: HIV Non-AIDS Cases by Gender, Age Group and Race/Ethnicity..... | 8 |
| Table 10: AIDS Cases by Race/Ethnicity and Year of AIDS Diagnosis..... | 9 |
| Table 11: HIV/AIDS Cases by Race/Ethnicity and Year of Initial HIV Diagnosis..... | 9 |
| Table 12: AIDS Incidence, Mortality, and Prevalence by Year..... | 10 |
| Table 13: AIDS Cases by Initial AIDS-Defining Condition and Year of AIDS Diagnosis..... | 11 |
| Table 14: Cumulative AIDS Indicator Conditions among Persons with AIDS..... | 12 |
| Table 15: Living Adult/Adolescent HIV/AIDS Cases by Transmission Category..... | 13 |
| Table 16: Living HIV/AIDS Cases by Transmission Category and Race/Ethnicity..... | 13 |
| Table 17: Living HIV/AIDS Cases by Gender, Age Group and Race/Ethnicity..... | 14 |

The HIV/AIDS Surveillance Report is published quarterly by the
San Francisco Department of Public Health, HIV Epidemiology Section
25 Van Ness Avenue, Suite 500, San Francisco, CA 94102; Phone (415) 554-9050, FAX (415) 431-0353
Director of Health: Barbara A. Garcia, MPA;
Health Officer and Director of Population Health and Prevention. Tomas Aragon, MD, DrPH;
Section Director: Susan Scheer, PhD, MPH; Surveillance Unit Director: Ling Hsu, MPH;
Program Coordinators: Maree Kay Parisi, Viva Delgado, MPH;
Epidemiologists: Mia Miao-Jung Chen, PhD, MPH, Priscilla Lee Chu, DrPH, MPH, Alison Hughes, MPH,
Kara O'Keefe, PhD, MPH, Sharon Pipkin, MPH, Tony Su, MPH, Annie Vu, MPH
The HIV/AIDS Surveillance Report is accessible via internet:
<http://www.sfdph.org/dph/files/reports/default.asp>

HIV/AIDS Surveillance Summary

San Francisco (as of 09/30/2012)

| | |
|---------------------------------|--------|
| Cumulative cases ¹ : | |
| AIDS | 29,271 |
| HIV non-AIDS ² | 6,042 |
| Cumulative AIDS deaths: | |
| | 19,695 |
| Living HIV/AIDS cases | |
| | 15,631 |


California³ (as of 12/31/2011)

| | |
|-------------------------|---------|
| Cumulative cases: | |
| AIDS | 161,847 |
| HIV non-AIDS | 44,554 |
| Cumulative AIDS deaths: | |
| | 91,906 |
| Living HIV/AIDS cases: | |
| | 112,602 |

United States⁴

| | |
|--|-----------|
| Cumulative AIDS cases (as of 12/31/2010): | 1,140,203 |
| Cumulative AIDS deaths (as of 12/31/2009): | 624,184 |
| Living HIV/AIDS cases (as of 12/31/2009): | 808,090 |

San Francisco AIDS Incidence, Mortality, and Prevalence by Year, 1980-2012⁵



1. Includes SF residents diagnosed in SF and SF residents diagnosed in other jurisdictions.
2. Includes HIV non-AIDS cases reported by name. HIV non-AIDS cases reported by a non-name code between July 2002 and April 2006 whose names have not been ascertained are not included.
3. CA data source: www.cdph.ca.gov/data/statistics/Pages/OAHIVAIDSStatistics.aspx.
4. US data source: CDC HIV Surveillance Report, 2010 vol. 22. www.cdc.gov/hiv/topics/surveillance/resources/reports/.
5. Reporting for recent years is incomplete. See Table 12 for actual numbers per year.

**HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 09/30/2012**

Table 1. Adult/Adolescent HIV/AIDS Cases (>12 years) by Transmission Category, San Francisco, 1980-2012

| Transmission Category | AIDS | | HIV non-AIDS# | |
|--|--------------|---------------|---------------|---------------|
| | No. | (%) | No. | (%) |
| Gay or bisexual male | 21312 | (72.9) | 4423 | (73.3) |
| Heterosexual male injection drug user | 1393 | (4.8) | 199 | (3.3) |
| Heterosexual female injection drug user | 725 | (2.5) | 141 | (2.3) |
| Gay or bisexual male injection drug user | 4396 | (15.0) | 723 | (12.0) |
| Lesbian or bisexual injection drug user | 69 | (0.2) | 25 | (0.4) |
| Transgender (1) | 445 | (1.5) | 132 | (2.2) |
| Hemophiliac | 16 | (0.1) | 2 | (0.0) |
| Heterosexual contact male (2) | 164 | (0.6) | 54 | (0.9) |
| Heterosexual contact female (2) | 330 | (1.1) | 136 | (2.3) |
| Transfusion recipient | 140 | (0.5) | 1 | (0.0) |
| Risk not reported/Other (3) | 243 | (0.8) | 194 | (3.2) |
| Total | 29233 | (100) | 6030 | (100) |

* Residents of San Francisco at time of HIV/AIDS diagnosis.

Includes HIV non-AIDS cases reported by name. Excludes HIV non-AIDS cases reported by a non-name code between July 2002 and April 2006 whose names have not been ascertained.

- (1) Transgender information was collected since September 1996. Data prior to this are incomplete.
- (2) Includes persons who have had heterosexual contact with a person with HIV/AIDS or with a person who is at risk for HIV.
- (3) Includes persons for whom risk information is incomplete (due to death, refusal to be interviewed or loss to follow-up), cases still under investigation, or interviewed patients who offered no plausible risk for HIV.

**HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 09/30/2012**

Table 2. AIDS Cases by Gender and Year of AIDS Diagnosis, San Francisco, 1980-2012

| Gender | Year of AIDS Diagnosis (1) | | | | | |
|-----------------|----------------------------|------------------|------------------|------------------|------------------|------------------|
| | < 2002 No. (%) | 2002 No. (%) | 2003 No. (%) | 2004 No. (%) | 2005 No. (%) | 2006 No. (%) |
| Male | 23721 (95.1) | 442 (88.9) | 495 (88.4) | 428 (88.1) | 425 (89.3) | 401 (90.1) |
| Female | 920 (3.7) | 35 (7.0) | 40 (7.1) | 41 (8.4) | 38 (8.0) | 30 (6.7) |
| Transgender (2) | 291 (1.2) | 20 (4.0) | 25 (4.5) | 17 (3.5) | 13 (2.7) | 14 (3.1) |
| Total | 24932 (100) | 497 (100) | 560 (100) | 486 (100) | 476 (100) | 445 (100) |

| Gender | Year of AIDS Diagnosis (1) | | | | | |
|-----------------|----------------------------|------------------|------------------|------------------|------------------|------------------|
| | 2007 No. (%) | 2008 No. (%) | 2009 No. (%) | 2010 No. (%) | 2011 No. (%) | 2012 No. (%) |
| Male | 406 (90.2) | 375 (86.8) | 286 (89.4) | 263 (89.8) | 209 (85.3) | 117 (86.7) |
| Female | 33 (7.3) | 42 (9.7) | 24 (7.5) | 16 (5.5) | 24 (9.8) | 15 (11.1) |
| Transgender (2) | 11 (2.4) | 15 (3.5) | 10 (3.1) | 14 (4.8) | 12 (4.9) | 3 (2.2) |
| Total | 450 (100) | 432 (100) | 320 (100) | 293 (100) | 245 (100) | 135 (100) |

Table 3. HIV/AIDS Cases by Gender and Year of Initial HIV Diagnosis#, San Francisco, 2007-2012

| Gender | Year of Initial HIV Diagnosis (1) | | | | | |
|-----------------|-----------------------------------|------------------|------------------|------------------|------------------|------------------|
| | 2007 No. (%) | 2008 No. (%) | 2009 No. (%) | 2010 No. (%) | 2011 No. (%) | 2012 No. (%) |
| Male | 467 (87.5) | 457 (89.4) | 414 (90.8) | 380 (90.0) | 350 (87.9) | 272 (94.4) |
| Female | 44 (8.2) | 38 (7.4) | 24 (5.3) | 33 (7.8) | 41 (10.3) | 15 (5.2) |
| Transgender (2) | 23 (4.3) | 16 (3.1) | 18 (3.9) | 9 (2.1) | 7 (1.8) | 1 (0.3) |
| Total | 534 (100) | 511 (100) | 456 (100) | 422 (100) | 398 (100) | 288 (100) |

* Residents of San Francisco at time of HIV/AIDS diagnosis.

Includes persons with a diagnosis of HIV (not AIDS), an initial diagnosis of HIV (not AIDS) and later diagnosed with AIDS, and concurrent diagnosis of HIV and AIDS. The initial year of HIV diagnosis was determined based on the earliest date of HIV antibody test, viral load or CD4 test, initiation of antiretroviral therapy, or patient self-report of a positive HIV test.

(1) Data in recent years are incomplete due to delays in case reporting.

(2) Transgender information was collected since September 1996. Data prior to this are incomplete.

**HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 09/30/2012**

Table 4. AIDS Cases by Transmission Category and Race/Ethnicity, San Francisco San Francisco, 1980-2012

| Transmission Category (1) | White No. (%) | African American No. (%) | Latino No. (%) | Asian/ Pacific Islander No. (%) | Native American No. (%) |
|-----------------------------|------------------|--------------------------------|-------------------|--|-------------------------------|
| Adult/Adolescent | | | | | |
| Gay or bisexual male | 16288 (79.4) | 1579 (42.6) | 2678 (72.9) | 709 (76.3) | 53 (42.7) |
| Injection drug user (IDU) | 732 (3.6) | 1046 (28.2) | 250 (6.8) | 36 (3.9) | 18 (14.5) |
| Gay or bisexual male IDU | 3173 (15.5) | 740 (20.0) | 515 (14.0) | 86 (9.3) | 46 (37.1) |
| Lesbian or bisexual IDU | 31 (0.2) | 24 (0.6) | 9 (0.2) | 2 (0.2) | 2 (1.6) |
| Hemophiliac | 8 (0.0) | 2 (0.1) | 5 (0.1) | 1 (0.1) | 0 (0.0) |
| Heterosexual (2) | 122 (0.6) | 204 (5.5) | 119 (3.2) | 47 (5.1) | 4 (3.2) |
| Transfusion recipient | 66 (0.3) | 22 (0.6) | 26 (0.7) | 20 (2.2) | 0 (0.0) |
| Risk not reported/Other (3) | 81 (0.4) | 75 (2.0) | 61 (1.7) | 23 (2.5) | 1 (0.8) |
| Pediatric (0-12 years) (4) | 8 (0.0) | 13 (0.4) | 10 (0.3) | 5 (0.5) | 0 (0.0) |
| Total | 20509 (100) | 3705 (100) | 3673 (100) | 929 (100) | 124 (100) |

Table 5. HIV Non-AIDS Cases# by Transmission Category and Race/Ethnicity, San Francisco

| Transmission Category (1) | White No. (%) | African American No. (%) | Latino No. (%) | Asian/ Pacific Islander No. (%) | Native American No. (%) |
|-----------------------------|------------------|--------------------------------|-------------------|--|-------------------------------|
| Adult/Adolescent | | | | | |
| Gay or bisexual male | 2956 (79.3) | 371 (47.6) | 778 (76.9) | 269 (82.3) | 19 (55.9) |
| Injection drug user (IDU) | 132 (3.5) | 159 (20.4) | 35 (3.5) | 4 (1.2) | 4 (11.8) |
| Gay or bisexual male IDU | 507 (13.6) | 106 (13.6) | 101 (10.0) | 26 (8.0) | 10 (29.4) |
| Lesbian or bisexual IDU | 11 (0.3) | 6 (0.8) | 4 (0.4) | 0 (0.0) | 0 (0.0) |
| Hemophiliac | 2 (0.1) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) |
| Heterosexual (2) | 42 (1.1) | 85 (10.9) | 46 (4.5) | 14 (4.3) | 0 (0.0) |
| Transfusion recipient | 0 (0.0) | 1 (0.1) | 0 (0.0) | 0 (0.0) | 0 (0.0) |
| Risk not reported/Other (3) | 75 (2.0) | 49 (6.3) | 42 (4.2) | 14 (4.3) | 1 (2.9) |
| Pediatric (0-12 years) (4) | 2 (0.1) | 2 (0.3) | 6 (0.6) | 0 (0.0) | 0 (0.0) |
| Total | 3727 (100) | 779 (100) | 1012 (100) | 327 (100) | 34 (100) |

* Residents of San Francisco at time of HIV/AIDS diagnosis.

Includes HIV non-AIDS cases reported by name. Excludes HIV non-AIDS cases reported by a non-name code between July 2002 and April 2006 whose names have not been ascertained.

- (1) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
- (2) Includes persons who have had heterosexual contact with a person with HIV/AIDS or with a person who is at risk for HIV.
- (3) Includes persons for whom risk information is incomplete (due to death, refusal to be interviewed or loss to follow-up), cases still under investigation, or interviewed patients who offered no plausible risk for HIV.
- (4) Includes children who have hemophilia or other coagulation disorder, have received a blood transfusion, or who have acquired their infection from an infected mother during the perinatal period.

HIV/AIDS Quarterly Surveillance Report

Summary of San Francisco Residents with HIV/AIDS*

Reported as of 09/30/2012

Table 6. AIDS Cases by Transmission Category and Year of AIDS Diagnosis, San Francisco, 1980-2012

| Transmission Category (2) | Year of AIDS Diagnosis (1) | | | | | |
|-----------------------------|----------------------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | < 2002 No. (%) | 2002 No. (%) | 2003 No. (%) | 2004 No. (%) | 2005 No. (%) | 2006 No. (%) |
| Adult/Adolescent | | | | | | |
| Gay or bisexual male | 18787 (75.4) | 304 (61.2) | 353 (63.0) | 317 (65.2) | 293 (61.6) | 287 (64.5) |
| Injection drug user (IDU) | 1688 (6.8) | 58 (11.7) | 70 (12.5) | 51 (10.5) | 50 (10.5) | 38 (8.5) |
| Gay or bisexual male IDU | 3781 (15.2) | 107 (21.5) | 105 (18.8) | 91 (18.7) | 98 (20.6) | 90 (20.2) |
| Lesbian or bisexual IDU | 48 (0.2) | 4 (0.8) | 3 (0.5) | 1 (0.2) | 3 (0.6) | 2 (0.4) |
| Hemophiliac | 15 (0.1) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) |
| Heterosexual (3) | 290 (1.2) | 15 (3.0) | 22 (3.9) | 16 (3.3) | 19 (4.0) | 20 (4.5) |
| Transfusion recipient | 138 (0.6) | 1 (0.2) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) |
| Risk not reported/Other (4) | 149 (0.6) | 7 (1.4) | 6 (1.1) | 10 (2.1) | 13 (2.7) | 8 (1.8) |
| Pediatric (0-12 years) (5) | 36 (0.1) | 1 (0.2) | 1 (0.2) | 0 (0.0) | 0 (0.0) | 0 (0.0) |
| Total | 24932 (100) | 497 (100) | 560 (100) | 486 (100) | 476 (100) | 445 (100) |

| Transmission Category (2) | Year of AIDS Diagnosis (1) | | | | | |
|-----------------------------|----------------------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | 2007 No. (%) | 2008 No. (%) | 2009 No. (%) | 2010 No. (%) | 2011 No. (%) | 2012 No. (%) |
| Adult/Adolescent | | | | | | |
| Gay or bisexual male | 293 (65.1) | 265 (61.3) | 207 (64.7) | 170 (58.0) | 144 (58.8) | 88 (65.2) |
| Injection drug user (IDU) | 37 (8.2) | 30 (6.9) | 33 (10.3) | 23 (7.8) | 34 (13.9) | 11 (8.1) |
| Gay or bisexual male IDU | 82 (18.2) | 91 (21.1) | 51 (15.9) | 74 (25.3) | 43 (17.6) | 21 (15.6) |
| Lesbian or bisexual IDU | 0 (0.0) | 4 (0.9) | 2 (0.6) | 0 (0.0) | 1 (0.4) | 1 (0.7) |
| Hemophiliac | 0 (0.0) | 0 (0.0) | 1 (0.3) | 0 (0.0) | 0 (0.0) | 0 (0.0) |
| Heterosexual (3) | 30 (6.7) | 30 (6.9) | 17 (5.3) | 17 (5.8) | 15 (6.1) | 8 (5.9) |
| Transfusion recipient | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 1 (0.4) | 0 (0.0) |
| Risk not reported/Other (4) | 8 (1.8) | 12 (2.8) | 9 (2.8) | 9 (3.1) | 7 (2.9) | 6 (4.4) |
| Pediatric (0-12 years) (5) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) |
| Total | 450 (100) | 432 (100) | 320 (100) | 293 (100) | 245 (100) | 135 (100) |

* Residents of San Francisco at time of HIV/AIDS diagnosis.

- (1) Data in recent years are incomplete due to delays in case reporting.
- (2) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
- (3) Includes persons who have had heterosexual contact with a person with HIV/AIDS or with a person who is at risk for HIV.
- (4) Includes persons for whom risk information is incomplete (due to death, refusal to be interviewed or loss to follow-up), cases still under investigation, or interviewed patients who offered no plausible risk for HIV.
- (5) Includes children who have hemophilia or other coagulation disorder, have received a blood transfusion, or who have acquired their infection from an infected mother during the perinatal period.

**HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 09/30/2012**

Table 7. HIV/AIDS Cases by Transmission Category and Year of Initial HIV Diagnosis#, San Francisco, 2007-2012

| Transmission Category (2) | Year of Initial HIV Diagnosis (1) | | | | | |
|-----------------------------|-----------------------------------|-----------------|-----------------|-----------------|-----------------|-----------------|
| | 2007 No. (%) | 2008 No. (%) | 2009 No. (%) | 2010 No. (%) | 2011 No. (%) | 2012 No. (%) |
| Adult/Adolescent | | | | | | |
| Gay or bisexual male | 354 (66.3) | 374 (73.2) | 329 (72.1) | 271 (64.2) | 283 (71.1) | 230 (79.9) |
| Injection drug user (IDU) | 33 (6.2) | 26 (5.1) | 21 (4.6) | 32 (7.6) | 22 (5.5) | 3 (1.0) |
| Gay or bisexual male IDU | 82 (15.4) | 56 (11.0) | 66 (14.5) | 62 (14.7) | 43 (10.8) | 26 (9.0) |
| Lesbian or bisexual IDU | 2 (0.4) | 4 (0.8) | 4 (0.9) | 1 (0.2) | 3 (0.8) | 1 (0.3) |
| Hemophiliac | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) |
| Heterosexual (3) | 42 (7.9) | 33 (6.5) | 19 (4.2) | 33 (7.8) | 23 (5.8) | 14 (4.9) |
| Transfusion recipient | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) |
| Risk not reported/Other (4) | 21 (3.9) | 18 (3.5) | 17 (3.7) | 23 (5.5) | 24 (6.0) | 14 (4.9) |
| Pediatric (0-12 years) (5) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) |
| Total | 534 (100) | 511 (100) | 456 (100) | 422 (100) | 398 (100) | 288 (100) |

* Residents of San Francisco at time of HIV/AIDS diagnosis.

Includes persons with a diagnosis of HIV (not AIDS), an initial diagnosis of HIV (not AIDS) and later diagnosed with AIDS, and concurrent diagnosis of HIV and AIDS. The initial year of HIV diagnosis was determined based on the earliest date of HIV antibody test, viral load or CD4 test, initiation of antiretroviral therapy, or patient self-report of a positive HIV test.

- (1) Data in recent years are incomplete due to delays in case reporting.
- (2) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
- (3) Includes persons who have had heterosexual contact with a person with HIV/AIDS or with a person who is at risk for HIV.
- (4) Includes persons for whom risk information is incomplete (due to death, refusal to be interviewed or loss to follow-up), cases still under investigation, or interviewed patients who offered no plausible risk for HIV.
- (5) Includes children who have hemophilia or other coagulation disorder, have received a blood transfusion, or who have acquired their infection from an infected mother during the perinatal period.

HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 09/30/2012

Table 8. AIDS Cases by Gender, Age Group and Race/Ethnicity, San Francisco, 1980-2012

| Male Age at AIDS Diagnosis (Years) | White No. (%) | African American No. (%) | Latino No. (%) | Asian/ Pacific Islander No. (%) | Native American No. (%) | Total (1) No. (%) |
|---|---------------------|--------------------------------|--------------------|--|-------------------------------|----------------------|
| 0 - 12 | 4 (0.0) | 4 (0.1) | 5 (0.1) | 4 (0.5) | 0 (0.0) | 19 (0.1) |
| 13 - 19 | 12 (0.1) | 1 (0.0) | 15 (0.4) | 1 (0.1) | 2 (1.9) | 31 (0.1) |
| 20 - 24 | 311 (1.6) | 70 (2.3) | 132 (3.9) | 26 (3.2) | 4 (3.8) | 550 (2.0) |
| 25 - 29 | 1673 (8.4) | 250 (8.3) | 472 (14.1) | 92 (11.2) | 21 (19.8) | 2544 (9.2) |
| 30 - 39 | 8858 (44.3) | 1164 (38.8) | 1571 (46.8) | 369 (44.8) | 51 (48.1) | 12141 (44.0) |
| 40 - 49 | 6591 (33.0) | 1004 (33.5) | 844 (25.2) | 244 (29.6) | 24 (22.6) | 8791 (31.9) |
| 50 - 59 | 2031 (10.2) | 394 (13.1) | 254 (7.6) | 66 (8.0) | 4 (3.8) | 2779 (10.1) |
| 60 + | 509 (2.5) | 111 (3.7) | 62 (1.8) | 22 (2.7) | 0 (0.0) | 713 (2.6) |
| Male subtotal | 19989 (100) | 2998 (100) | 3355 (100) | 824 (100) | 106 (100) | 27568 (100) |
| Female Age at AIDS Diagnosis (Years) | White No. (%) | African American No. (%) | Latino No. (%) | Asian/ Pacific Islander No. (%) | Native American No. (%) | Total (1) No. (%) |
| 0 - 12 | 4 (1.0) | 9 (1.6) | 5 (2.7) | 1 (1.5) | 0 (0.0) | 19 (1.5) |
| 13 - 19 | 1 (0.3) | 2 (0.4) | 2 (1.1) | 0 (0.0) | 0 (0.0) | 5 (0.4) |
| 20 - 24 | 16 (4.0) | 10 (1.8) | 11 (5.9) | 4 (6.1) | 1 (7.1) | 43 (3.4) |
| 25 - 29 | 40 (10.0) | 47 (8.3) | 26 (14.1) | 11 (16.7) | 1 (7.1) | 127 (10.1) |
| 30 - 39 | 157 (39.3) | 205 (36.0) | 61 (33.0) | 25 (37.9) | 8 (57.1) | 463 (36.8) |
| 40 - 49 | 106 (26.5) | 195 (34.3) | 48 (25.9) | 18 (27.3) | 4 (28.6) | 377 (30.0) |
| 50 - 59 | 45 (11.3) | 76 (13.4) | 18 (9.7) | 4 (6.1) | 0 (0.0) | 148 (11.8) |
| 60 + | 31 (7.8) | 25 (4.4) | 14 (7.6) | 3 (4.5) | 0 (0.0) | 76 (6.0) |
| Female subtotal | 400 (100) | 569 (100) | 185 (100) | 66 (100) | 14 (100) | 1258 (100) |
| Transgender (2) Age at AIDS Diagnosis (Years) | White No. (%) | African American No. (%) | Latino No. (%) | Asian/ Pacific Islander Native American No. (%) | Total (1) No. (%) | |
| 13 - 29 | 25 (20.8) | 31 (22.5) | 35 (26.3) | 12 (27.9) | 106 (23.8) | |
| 30 - 39 | 57 (47.5) | 50 (36.2) | 62 (46.6) | 20 (46.5) | 194 (43.6) | |
| 40 + | 38 (31.7) | 57 (41.3) | 36 (27.1) | 11 (25.6) | 145 (32.6) | |
| Transgender subtotal | 120 (100) | 138 (100) | 133 (100) | 43 (100) | 445 (100) | |

* Residents of San Francisco at time of HIV/AIDS diagnosis.

(1) Total includes persons with multiple or unknown race.

(2) Transgender information was collected since September 1996. Certain age or race/ethnic groups are combined for transgender cases because of small number.

HIV/AIDS Quarterly Surveillance Report

Summary of San Francisco Residents with HIV/AIDS*

Reported as of 09/30/2012

Table 9. HIV Non-AIDS Cases# by Gender, Age Group and Race/Ethnicity, San Francisco

| Male Age at Initial HIV Diagnosis (Years) | White No. (%) | African American No. (%) | Latino No. (%) | Asian/ Pacific Islander No. (%) | Native American No. (%) | Total (1) No. (%) |
|--|--------------------|--------------------------------|-------------------|--|-------------------------------|----------------------|
| 0 - 12 | 1 (0.0) | 1 (0.2) | 1 (0.1) | 0 (0.0) | 0 (0.0) | 3 (0.1) |
| 13 - 19 | 26 (0.7) | 19 (3.3) | 21 (2.3) | 9 (3.1) | 0 (0.0) | 76 (1.4) |
| 20 - 24 | 301 (8.4) | 56 (9.6) | 120 (13.4) | 31 (10.5) | 4 (13.3) | 531 (9.6) |
| 25 - 29 | 545 (15.2) | 88 (15.1) | 198 (22.0) | 68 (23.1) | 6 (20.0) | 935 (16.9) |
| 30 - 39 | 1564 (43.5) | 192 (32.9) | 374 (41.6) | 141 (47.8) | 15 (50.0) | 2342 (42.3) |
| 40 - 49 | 861 (23.9) | 163 (27.9) | 152 (16.9) | 40 (13.6) | 5 (16.7) | 1253 (22.6) |
| 50 - 59 | 243 (6.8) | 48 (8.2) | 27 (3.0) | 6 (2.0) | 0 (0.0) | 326 (5.9) |
| 60 + | 54 (1.5) | 17 (2.9) | 5 (0.6) | 0 (0.0) | 0 (0.0) | 77 (1.4) |
| Male subtotal | 3595 (100) | 584 (100) | 898 (100) | 295 (100) | 30 (100) | 5543 (100) |
| Female Age at Initial HIV Diagnosis (Years) | White No. (%) | African American No. (%) | Latino No. (%) | Asian/ Pacific Islander No. (%) | Native American No. (%) | Total (1) No. (%) |
| 0 - 12 | 1 (0.9) | 1 (0.7) | 5 (6.8) | 0 (0.0) | 0 (0.0) | 9 (2.5) |
| 13 - 19 | 2 (1.9) | 2 (1.4) | 2 (2.7) | 0 (0.0) | 0 (0.0) | 7 (1.9) |
| 20 - 24 | 11 (10.4) | 13 (8.8) | 14 (19.2) | 0 (0.0) | 0 (0.0) | 40 (10.9) |
| 25 - 29 | 20 (18.9) | 15 (10.1) | 11 (15.1) | 5 (23.8) | 0 (0.0) | 52 (14.2) |
| 30 - 39 | 27 (25.5) | 46 (31.1) | 22 (30.1) | 6 (28.6) | 3 (100) | 109 (29.7) |
| 40 - 49 | 30 (28.3) | 51 (34.5) | 14 (19.2) | 7 (33.3) | 0 (0.0) | 105 (28.6) |
| 50 - 59 | 15 (14.2) | 17 (11.5) | 4 (5.5) | 2 (9.5) | 0 (0.0) | 40 (10.9) |
| 60 + | 0 (0.0) | 3 (2.0) | 1 (1.4) | 1 (4.8) | 0 (0.0) | 5 (1.4) |
| Female subtotal | 106 (100) | 148 (100) | 73 (100) | 21 (100) | 3 (100) | 367 (100) |
| Transgender (2) Age at Initial HIV Diagnosis (Years) | White No. (%) | African American No. (%) | Latino No. (%) | Asian/ Pacific Islander Native American No. (%) | Total (1) No. (%) | |
| 13 - 29 | 8 (30.8) | 23 (48.9) | 17 (41.5) | 4 (33.3) | 55 (41.7) | |
| 30 + | 18 (69.2) | 24 (51.1) | 24 (58.5) | 8 (66.7) | 77 (58.3) | |
| Transgender subtotal | 26 (100) | 47 (100) | 41 (100) | 12 (100) | 132 (100) | |

* Residents of San Francisco at time of HIV/AIDS diagnosis.

Includes HIV non-AIDS cases reported by name. Excludes HIV non-AIDS cases reported by a non-name code between July 2002 and April 2006 whose names have not been ascertained.

(1) Total includes persons with multiple or unknown race.

(2) Transgender information was collected since September 1996. Certain age or race/ethnic groups are combined for transgender cases because of small number.

HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 09/30/2012

Table 10. AIDS Cases by Race/Ethnicity and Year of AIDS Diagnosis, San Francisco, 1980-2012

| Race/Ethnicity | Year of AIDS Diagnosis (1) | | | | | |
|------------------------|----------------------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | < 2002 No. (%) | 2002 No. (%) | 2003 No. (%) | 2004 No. (%) | 2005 No. (%) | 2006 No. (%) |
| White | 18165 (72.9) | 289 (58.1) | 282 (50.4) | 260 (53.5) | 269 (56.5) | 250 (56.2) |
| African American | 2941 (11.8) | 88 (17.7) | 105 (18.8) | 79 (16.3) | 83 (17.4) | 79 (17.8) |
| Latino | 2837 (11.4) | 77 (15.5) | 127 (22.7) | 113 (23.3) | 89 (18.7) | 80 (18.0) |
| Asian/Pacific Islander | 672 (2.7) | 28 (5.6) | 33 (5.9) | 22 (4.5) | 25 (5.3) | 22 (4.9) |
| Native American | 95 (0.4) | 2 (0.4) | 5 (0.9) | 4 (0.8) | 3 (0.6) | 4 (0.9) |
| Total (2) | 24932 (100) | 497 (100) | 560 (100) | 486 (100) | 476 (100) | 445 (100) |

| Race/Ethnicity | Year of AIDS Diagnosis& | | | | | |
|------------------------|-------------------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | 2007 No. (%) | 2008 No. (%) | 2009 No. (%) | 2010 No. (%) | 2011 No. (%) | 2012 No. (%) |
| White | 260 (57.8) | 223 (51.6) | 168 (52.5) | 145 (49.5) | 130 (53.1) | 68 (50.4) |
| African American | 76 (16.9) | 81 (18.8) | 60 (18.8) | 54 (18.4) | 40 (16.3) | 19 (14.1) |
| Latino | 76 (16.9) | 73 (16.9) | 65 (20.3) | 60 (20.5) | 44 (18.0) | 32 (23.7) |
| Asian/Pacific Islander | 27 (6.0) | 40 (9.3) | 13 (4.1) | 24 (8.2) | 15 (6.1) | 8 (5.9) |
| Native American | 0 (0.0) | 3 (0.7) | 3 (0.9) | 2 (0.7) | 2 (0.8) | 1 (0.7) |
| Total (2) | 450 (100) | 432 (100) | 320 (100) | 293 (100) | 245 (100) | 135 (100) |

Table 11. HIV/AIDS Cases by Race/Ethnicity and Year of Initial HIV Diagnosis#, San Francisco, 2007-2012

| Race/Ethnicity | Year of Initial HIV Diagnosis (1) | | | | | |
|------------------------|-----------------------------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | 2007 No. (%) | 2008 No. (%) | 2009 No. (%) | 2010 No. (%) | 2011 No. (%) | 2012 No. (%) |
| White | 275 (51.5) | 258 (50.5) | 234 (51.3) | 201 (47.6) | 206 (51.8) | 141 (49.0) |
| African American | 79 (14.8) | 82 (16.0) | 70 (15.4) | 60 (14.2) | 66 (16.6) | 29 (10.1) |
| Latino | 105 (19.7) | 115 (22.5) | 93 (20.4) | 102 (24.2) | 79 (19.8) | 72 (25.0) |
| Asian/Pacific Islander | 50 (9.4) | 40 (7.8) | 38 (8.3) | 41 (9.7) | 32 (8.0) | 34 (11.8) |
| Native American | 1 (0.2) | 3 (0.6) | 1 (0.2) | 3 (0.7) | 2 (0.5) | 3 (1.0) |
| Total (2) | 534 (100) | 511 (100) | 456 (100) | 422 (100) | 398 (100) | 288 (100) |

* Residents of San Francisco at time of HIV/AIDS diagnosis.

Includes persons with a diagnosis of HIV (not AIDS), an initial diagnosis of HIV (not AIDS) and later diagnosed with AIDS, and concurrent diagnosis of HIV and AIDS. The initial year of HIV diagnosis was determined based on the earliest date of HIV antibody test, viral load or CD4 test, initiation of antiretroviral therapy, or patient self-report of a positive HIV test.

(1) Data in recent years are incomplete due to delays in case reporting.

(2) Total includes persons with multiple or unknown race.

**HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 09/30/2012**

Table 12. AIDS Incidence, Mortality, and Prevalence by Year, San Francisco, 1980-2012

| Year | Number of AIDS Cases Reported per Year | Number of AIDS Cases Diagnosed per Year (1) | Number of AIDS Deaths Occurred per Year (1) | Number of Persons Living with AIDS (1) |
|-------|--|---|---|--|
| 1980 | 0 | 3 | 0 | 3 |
| 1981 | 21 | 26 | 8 | 21 |
| 1982 | 75 | 99 | 32 | 88 |
| 1983 | 197 | 274 | 111 | 251 |
| 1984 | 451 | 558 | 273 | 536 |
| 1985 | 673 | 860 | 534 | 862 |
| 1986 | 981 | 1236 | 807 | 1291 |
| 1987 | 1287 | 1629 | 878 | 2042 |
| 1988 | 1407 | 1763 | 1039 | 2766 |
| 1989 | 1584 | 2161 | 1276 | 3651 |
| 1990 | 1686 | 2046 | 1366 | 4331 |
| 1991 | 1685 | 2287 | 1509 | 5109 |
| 1992 | 1638 | 2330 | 1641 | 5798 |
| 1993 | 4267 | 2071 | 1600 | 6269 |
| 1994 | 1913 | 1789 | 1595 | 6463 |
| 1995 | 1632 | 1565 | 1483 | 6545 |
| 1996 | 1238 | 1082 | 992 | 6635 |
| 1997 | 1060 | 807 | 423 | 7019 |
| 1998 | 793 | 696 | 402 | 7313 |
| 1999 | 723 | 578 | 353 | 7538 |
| 2000 | 624 | 559 | 349 | 7748 |
| 2001 | 495 | 513 | 324 | 7937 |
| 2002 | 439 | 497 | 321 | 8113 |
| 2003 | 531 | 560 | 303 | 8370 |
| 2004 | 553 | 486 | 309 | 8547 |
| 2005 | 498 | 476 | 312 | 8711 |
| 2006 | 421 | 445 | 289 | 8867 |
| 2007 | 536 | 450 | 270 | 9047 |
| 2008 | 554 | 432 | 229 | 9250 |
| 2009 | 348 | 320 | 210 | 9360 |
| 2010 | 402 | 293 | 193 | 9460 |
| 2011 | 333 | 245 | 165 | 9540 |
| 2012 | 226 | 135 | 99 | 9576 |
| Total | 29271 | 29271 | 19695 | |

* Residents of San Francisco at time of HIV/AIDS diagnosis.

(1) Data in recent years are incomplete due to delays in case/death reporting.

HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 09/30/2012

Table 13. AIDS Cases by Initial AIDS-Defining Condition and Year of AIDS Diagnosis, San Francisco, 1980-2012

| Initial AIDS-Defining Condition (1) | Year of AIDS Diagnosis | | |
|---|------------------------|----------------------|----------------------|
| | 1980-1989 No. (%) | 1990-1995 No. (%) | 1996-2012 No. (%) |
| Bacterial infections, recurrent, <13 years | 1 (0.0) | 5 (0.0) | 3 (0.0) |
| Candidiasis of bronchi, trachea, or lungs | 25 (0.3) | 13 (0.1) | 21 (0.2) |
| Candidiasis of esophagus | 301 (3.5) | 307 (2.5) | 180 (2.1) |
| Invasive cervical cancer | 1 (0.0) | 1 (0.0) | 7 (0.1) |
| Coccidioidomycosis, disseminated or extrapulmonary | 8 (0.1) | 3 (0.0) | 4 (0.0) |
| Cryptococcosis, extrapulmonary | 247 (2.9) | 157 (1.3) | 92 (1.1) |
| Cryptosporidiosis, intestinal (>1 mo. duration) | 137 (1.6) | 203 (1.7) | 65 (0.8) |
| Cytomegalovirus (except liver, spleen, lymph nodes), CMV retinitis with loss of vision | 175 (2.0) | 139 (1.1) | 51 (0.6) |
| HIV encephalopathy | 54 (0.6) | 92 (0.8) | 21 (0.2) |
| Herpes simplex: chronic (>1 mo.), bronchitis, pneumonitis, esophagitis | 225 (2.6) | 196 (1.6) | 70 (0.8) |
| Histoplasmosis | 75 (0.9) | 53 (0.4) | 22 (0.3) |
| Isosporiasis, intestinal (>1 mo. duration) | 14 (0.2) | 15 (0.1) | 13 (0.2) |
| Kaposi's sarcoma | 10 (0.1) | 11 (0.1) | 8 (0.1) |
| Lymphoid interstitial pneumonia/ pulmonary lymphoid hyperplasia, <13 years | 1981 (23.0) | 870 (7.2) | 305 (3.6) |
| Lymphoma, Burkitt's (non-Hodgkin's) | 3 (0.0) | 1 (0.0) | 0 (0.0) |
| Lymphoma, immunoblastic (non-Hodgkin's) | 59 (0.7) | 32 (0.3) | 41 (0.5) |
| Lymphoma, primary in brain | 216 (2.5) | 163 (1.3) | 81 (0.9) |
| Mycobacterium avium complex or M. kansasii, disseminated or extrapulmonary | 27 (0.3) | 16 (0.1) | 11 (0.1) |
| Mycobacterium tuberculosis, pulmonary | 217 (2.5) | 166 (1.4) | 55 (0.6) |
| Mycobacterium tuberculosis, disseminated or extrapulmonary | 46 (0.5) | 162 (1.3) | 98 (1.1) |
| Mycobacterium other species, disseminated or extrapulmonary | 72 (0.8) | 93 (0.8) | 36 (0.4) |
| Pneumocystis carinii pneumonia | 19 (0.2) | 27 (0.2) | 8 (0.1) |
| Pneumonia, recurrent | 3794 (44.1) | 1809 (15.0) | 790 (9.2) |
| Progressive multifocal leukoencephalopathy | 37 (0.4) | 144 (1.2) | 120 (1.4) |
| Salmonella sepsis, recurrent | 40 (0.5) | 15 (0.1) | 8 (0.1) |
| Toxoplasmosis of brain, >1 month of age | 10 (0.1) | 2 (0.0) | 2 (0.0) |
| Wasting syndrome | 178 (2.1) | 102 (0.8) | 39 (0.5) |
| CD4 T lymphocyte count <200 or percent <14 (2) | 286 (3.3) | 448 (3.7) | 187 (2.2) |
| | 1217 (14.1) | 7487 (61.9) | 6512 (76.0) |
| Total (3) | 8609 (100) | 12088 (100) | 8574 (100) |

* Residents of San Francisco at time of HIV/AIDS diagnosis.

(1) Based on AIDS-defining conditions diagnosed in the same calendar month and year.

Cases may have more than one initial AIDS-defining condition.

(2) Cases whose initial AIDS defining condition is low CD4 count (or percent) only.

(3) Total represents unique AIDS cases diagnosed during the time period specified.

**HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 09/30/2012**

Table 14. Cumulative AIDS Indicator Conditions among Persons with AIDS, San Francisco, 1980-2012

| AIDS Indicator Condition (1) | Total No. (%) |
|--|------------------|
| Bacterial infections, recurrent, <13 years | 11 (0.0) |
| Candidiasis of bronchi, trachea, or lungs | 164 (0.6) |
| Candidiasis of esophagus | 2694 (9.2) |
| Invasive cervical cancer | 15 (0.1) |
| Coccidioidomycosis, disseminated or extrapulmonary | 63 (0.2) |
| Cryptococcosis, extrapulmonary | 1897 (6.5) |
| Cryptosporidiosis, intestinal (>1 mo. duration) | 1387 (4.7) |
| Cytomegalovirus (except liver, spleen, lymph nodes) | 2497 (8.5) |
| CMV retinitis with loss of vision | 2506 (8.6) |
| HIV encephalopathy | 2513 (8.6) |
| Herpes simplex: chronic (>1 mo.), bronchitis, pneumonitis, esophagitis | 469 (1.6) |
| Histoplasmosis | 161 (0.6) |
| Isosporiasis, intestinal (>1 mo. duration) | 67 (0.2) |
| Kaposi's sarcoma | 6800 (23.2) |
| Lymphoid interstitial pneumonia/pulmonary lymphoid hyperplasia, <13 years | 6 (0.0) |
| Lymphoma, Burkitt's (non-Hodgkin's) | 560 (1.9) |
| Lymphoma, immunoblastic (non-Hodgkin's) | 1145 (3.9) |
| Lymphoma, primary in brain | 393 (1.3) |
| Mycobacterium avium complex or M. kansasii, disseminated or extrapulmonary | 5117 (17.5) |
| Mycobacterium tuberculosis, pulmonary | 666 (2.3) |
| Mycobacterium tuberculosis, disseminated or extrapulmonary | 487 (1.7) |
| Mycobacterium other species, disseminated or extrapulmonary | 342 (1.2) |
| Pneumocystis carinii pneumonia | 11129 (38.0) |
| Pneumonia, recurrent | 1045 (3.6) |
| Progressive multifocal leukoencephalopathy | 315 (1.1) |
| Salmonella sepsis, recurrent | 62 (0.2) |
| Toxoplasmosis of brain, >1 month of age | 1180 (4.0) |
| Wasting syndrome | 4390 (15.0) |

* Residents of San Francisco at time of HIV/AIDS diagnosis.

(1) Includes initial and subsequent diagnoses of AIDS indicator conditions. Cases may have more than one condition.

**HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 09/30/2012**

Table 15. Living Adult/Adolescent HIV/AIDS Cases (>12 years) by Transmission Category, San Francisco

| Transmission Category | No. | (%) |
|--|--------------|---------------|
| Gay or bisexual male | 11294 | (72.4) |
| Heterosexual male injection drug user | 542 | (3.5) |
| Heterosexual female injection drug user | 399 | (2.6) |
| Gay or bisexual male injection drug user | 2166 | (13.9) |
| Lesbian or bisexual injection drug user | 57 | (0.4) |
| Transgender (1) | 344 | (2.2) |
| Hemophiliac | 4 | (0.0) |
| Heterosexual contact male (2) | 155 | (1.0) |
| Heterosexual contact female (2) | 331 | (2.1) |
| Transfusion recipient | 16 | (0.1) |
| Risk not reported/Other (3) | 284 | (1.8) |
| Total | 15592 | (100) |

Table 16. Living HIV/AIDS Cases by Transmission Category and Race/Ethnicity, San Francisco

| Transmission Category | White No. (%) | African American No. (%) | Latino No. (%) | Asian/ Pacific Islander No. (%) | Native American No. (%) |
|-----------------------------------|--------------------|--------------------------------|--------------------|--|-------------------------------|
| Adult/Adolescent | | | | | |
| Gay or bisexual male | 7661 (78.8) | 924 (45.1) | 2009 (75.4) | 650 (78.4) | 44 (49.4) |
| Injection drug user (IDU) | 341 (3.5) | 443 (21.6) | 112 (4.2) | 18 (2.2) | 14 (15.7) |
| Gay or bisexual male IDU | 1466 (15.1) | 376 (18.4) | 318 (11.9) | 82 (9.9) | 24 (27.0) |
| Lesbian or bisexual IDU | 26 (0.3) | 18 (0.9) | 7 (0.3) | 1 (0.1) | 2 (2.2) |
| Hemophiliac | 3 (0.0) | 1 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) |
| Heterosexual (2) | 103 (1.1) | 201 (9.8) | 131 (4.9) | 46 (5.5) | 4 (4.5) |
| Transfusion recipient | 6 (0.1) | 3 (0.1) | 4 (0.2) | 3 (0.4) | 0 (0.0) |
| Risk not reported/Other (3) | 108 (1.1) | 71 (3.5) | 68 (2.6) | 23 (2.8) | 1 (1.1) |
| Pediatric (0-12 years) (4) | 5 (0.1) | 10 (0.5) | 14 (0.5) | 6 (0.7) | 0 (0.0) |
| Total | 9719 (100) | 2047 (100) | 2663 (100) | 829 (100) | 89 (100) |

* Residents of San Francisco at time of HIV/AIDS diagnosis.

- (1) Transgender information was collected since September 1996. Data prior to this are incomplete.
- (2) Includes persons who have had heterosexual contact with a person with HIV/AIDS or with a person who is at risk for HIV.
- (3) Includes persons for whom risk information is incomplete (due to death, refusal to be interviewed or loss to follow-up), cases still under investigation, or interviewed patients who offered no plausible risk for HIV.
- (4) Includes children who have hemophilia or other coagulation disorder, have received a blood transfusion, or who have acquired their infection from an infected mother during the perinatal period.

HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 09/30/2012

Table 17. Living HIV/AIDS Cases by Gender, Age Group and Race/Ethnicity, San Francisco

| Male Current Age (Years) | White No. (%) | African American No. (%) | Latino No. (%) | Asian/ Pacific Islander No. (%) | Native American No. (%) | Total (1) No. (%) |
|---|--------------------|--------------------------------|--------------------|--|-------------------------------|----------------------|
| 0 - 12 | 0 (0.0) | 0 (0.0) | 1 (0.0) | 0 (0.0) | 0 (0.0) | 2 (0.0) |
| 13 - 19 | 2 (0.0) | 3 (0.2) | 0 (0.0) | 3 (0.4) | 0 (0.0) | 8 (0.1) |
| 20 - 24 | 33 (0.4) | 27 (1.7) | 34 (1.4) | 8 (1.1) | 0 (0.0) | 107 (0.7) |
| 25 - 29 | 167 (1.8) | 49 (3.2) | 110 (4.6) | 40 (5.4) | 1 (1.4) | 377 (2.6) |
| 30 - 39 | 833 (8.9) | 137 (8.8) | 457 (19.2) | 168 (22.8) | 17 (23.0) | 1680 (11.7) |
| 40 - 49 | 3036 (32.4) | 469 (30.2) | 915 (38.4) | 283 (38.3) | 35 (47.3) | 4836 (33.6) |
| 50 - 59 | 3408 (36.3) | 585 (37.7) | 643 (27.0) | 165 (22.4) | 19 (25.7) | 4878 (33.9) |
| 60 + | 1905 (20.3) | 282 (18.2) | 224 (9.4) | 71 (9.6) | 2 (2.7) | 2500 (17.4) |
| Male subtotal | 9384 (100) | 1552 (100) | 2384 (100) | 738 (100) | 74 (100) | 14388 (100) |
| Female Current Age (Years) | White No. (%) | African American No. (%) | Latino No. (%) | Asian/ Pacific Islander No. (%) | Native American No. (%) | Total (1) No. (%) |
| 0 - 12 | 0 (0.0) | 1 (0.3) | 1 (0.6) | 0 (0.0) | 0 (0.0) | 2 (0.2) |
| 13 - 19 | 1 (0.4) | 1 (0.3) | 6 (3.5) | 1 (1.8) | 0 (0.0) | 11 (1.2) |
| 20 - 24 | 2 (0.8) | 4 (1.1) | 5 (2.9) | 0 (0.0) | 0 (0.0) | 11 (1.2) |
| 25 - 29 | 9 (3.4) | 11 (2.9) | 13 (7.5) | 0 (0.0) | 0 (0.0) | 34 (3.8) |
| 30 - 39 | 32 (12.3) | 41 (10.9) | 30 (17.3) | 15 (26.3) | 2 (16.7) | 127 (14.1) |
| 40 - 49 | 102 (39.1) | 111 (29.5) | 49 (28.3) | 20 (35.1) | 4 (33.3) | 290 (32.3) |
| 50 - 59 | 86 (33.0) | 151 (40.2) | 40 (23.1) | 15 (26.3) | 6 (50.0) | 304 (33.8) |
| 60 + | 29 (11.1) | 56 (14.9) | 29 (16.8) | 6 (10.5) | 0 (0.0) | 120 (13.3) |
| Female subtotal | 261 (100) | 376 (100) | 173 (100) | 57 (100) | 12 (100) | 899 (100) |
| Transgender (2) Current Age (Years) | White No. (%) | African American No. (%) | Latino No. (%) | Asian/ Pacific Islander Native American No. (%) | Total (1) No. (%) | |
| 13 - 39 | 14 (18.9) | 35 (29.4) | 46 (43.4) | 10 (27.0) | 111 (32.3) | |
| 40 + | 60 (81.1) | 84 (70.6) | 60 (56.6) | 27 (73.0) | 233 (67.7) | |
| Transgender subtotal | 74 (100) | 119 (100) | 106 (100) | 37 (100) | 344 (100) | |

* Residents of San Francisco at time of HIV/AIDS diagnosis.

(1) Total includes persons with multiple or unknown race.

(2) Transgender information was collected since September 1996. Certain age or race/ethnic groups are combined for transgender cases because of small number.