

QUARTERLY HIV/AIDS SURVEILLANCE REPORT

San Francisco Department of Public Health HIV/AIDS Cases Reported Through March 2012

Contents	Page
Surveillance Summary.....	1
Table 1: Adult/Adolescent HIV/AIDS Cases by Transmission Category.....	2
Table 2: AIDS Cases by Gender and Year of AIDS Diagnosis.....	3
Table 3: HIV/AIDS Cases by Gender and Year of Initial HIV Diagnosis.....	3
Table 4: AIDS Cases by Transmission Category and Race/Ethnicity.....	4
Table 5: HIV Non-AIDS Cases by Transmission Category and Race/Ethnicity.....	4
Table 6: AIDS Cases by Transmission Category and Year of AIDS Diagnosis.....	5
Table 7: HIV/AIDS Cases by Transmission Category and Year of Initial HIV Diagnosis.....	6
Table 8: AIDS Cases by Gender, Age Group and Race/Ethnicity.....	7
Table 9: HIV Non-AIDS Cases by Gender, Age Group and Race/Ethnicity.....	8
Table 10: AIDS Cases by Race/Ethnicity and Year of AIDS Diagnosis.....	9
Table 11: HIV/AIDS Cases by Race/Ethnicity and Year of Initial HIV Diagnosis.....	9
Table 12: AIDS Incidence, Mortality, and Prevalence by Year.....	10
Table 13: AIDS Cases by Initial AIDS-Defining Condition and Year of AIDS Diagnosis.....	11
Table 14: Cumulative AIDS Indicator Conditions among Persons with AIDS.....	12
Table 15: Living Adult/Adolescent HIV/AIDS Cases by Transmission Category.....	13
Table 16: Living HIV/AIDS Cases by Transmission Category and Race/Ethnicity.....	13
Table 17: Living HIV/AIDS Cases by Gender, Age Group and Race/Ethnicity.....	14

The HIV/AIDS Surveillance Report is published quarterly by the
San Francisco Department of Public Health, HIV Epidemiology Section
25 Van Ness Avenue, Suite 500, San Francisco, CA 94102; Phone (415) 554-9050, FAX (415) 431-0353
Director of Health: Barbara A. Garcia, MPA;
Health Officer and Director of Population Health and Prevention. Tomas Aragon, MD, DrPH;
Section Director: Susan Scheer, PhD, MPH; Surveillance Unit Director: Ling Hsu, MPH;
Program Coordinators: Maree Kay Parisi, Viva Delgado, MPH;
Epidemiologists: Mia Miao-Jung Chen, PhD, MPH, Priscilla Lee Chu, DrPH, MPH, Alison Hughes, MPH,
Kara O'Keefe, PhD, MPH, Sharon Pipkin, MPH, Tony Su, MPH, Annie Vu, MPH
The HIV/AIDS Surveillance Report is accessible via internet:
<http://www.sfdph.org/dph/files/reports/default.asp>

HIV/AIDS Surveillance Summary

San Francisco (as of 03/31/2012)

Cumulative cases ¹ :	
AIDS	29,131
HIV non-AIDS ²	5,996
Cumulative AIDS deaths:	
	19,606
Living HIV/AIDS cases	
	15,523


California³ (as of 06/30/2011)

Cumulative cases:	
AIDS	160,760
HIV non-AIDS	43,501
Cumulative AIDS deaths:	
	91,371
Living HIV/AIDS cases:	
	111,100

United States⁴

Cumulative AIDS cases (as of 12/31/2010):	1,140,203
Cumulative AIDS deaths (as of 12/31/2009):	624,184
Living HIV/AIDS cases (as of 12/31/2009):	808,090

San Francisco AIDS Incidence, Mortality, and Prevalence by Year, 1980-2012⁵



1. Includes SF residents diagnosed in SF and SF residents diagnosed in other jurisdictions.
2. Includes HIV non-AIDS cases reported by name. HIV non-AIDS cases reported by a non-name code between July 2002 and April 2006 whose names have not been ascertained are not included.
3. CA data source: www.cdph.ca.gov/data/statistics/Pages/OAHIVAIDSStatistics.aspx.
4. US data source: CDC HIV Surveillance Report, 2010 vol. 22. www.cdc.gov/hiv/topics/surveillance/resources/reports/.
5. Reporting for recent years is incomplete. See Table 12 for actual numbers per year.

**HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 03/31/2012**

Table 1. Adult/Adolescent HIV/AIDS Cases (>12 years) by Transmission Category, San Francisco, 1980-2012

Transmission Category	AIDS		HIV non-AIDS#	
	No.	(%)	No.	(%)
Gay or bisexual male	21228	(73.0)	4363	(72.9)
Heterosexual male injection drug user	1400	(4.8)	211	(3.5)
Heterosexual female injection drug user	725	(2.5)	155	(2.6)
Gay or bisexual male injection drug user	4347	(14.9)	711	(11.9)
Lesbian or bisexual injection drug user	61	(0.2)	15	(0.3)
Transgender (1)	436	(1.5)	136	(2.3)
Hemophiliac	16	(0.1)	2	(0.0)
Heterosexual contact male (2)	166	(0.6)	53	(0.9)
Heterosexual contact female (2)	328	(1.1)	136	(2.3)
Transfusion recipient	144	(0.5)	1	(0.0)
Risk not reported/Other (3)	242	(0.8)	201	(3.4)
Total	29093	(100)	5984	(100)

* Residents of San Francisco at time of HIV/AIDS diagnosis.

Includes HIV non-AIDS cases reported by name. Excludes HIV non-AIDS cases reported by a non-name code between July 2002 and April 2006 whose names have not been ascertained.

- (1) Transgender information was collected since September 1996. Data prior to this are incomplete.
- (2) Includes persons who have had heterosexual contact with a person with HIV/AIDS or with a person who is at risk for HIV.
- (3) Includes persons for whom risk information is incomplete (due to death, refusal to be interviewed or loss to follow-up), cases still under investigation, or interviewed patients who offered no plausible risk for HIV.

**HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 03/31/2012**

Table 2. AIDS Cases by Gender and Year of AIDS Diagnosis, San Francisco, 1980-2012

Gender	Year of AIDS Diagnosis (1)					
	< 2002 No. (%)	2002 No. (%)	2003 No. (%)	2004 No. (%)	2005 No. (%)	2006 No. (%)
Male	23721 (95.2)	442 (89.1)	495 (88.4)	428 (88.2)	422 (89.0)	402 (90.1)
Female	920 (3.7)	35 (7.1)	40 (7.1)	40 (8.2)	38 (8.0)	30 (6.7)
Transgender (2)	289 (1.2)	19 (3.8)	25 (4.5)	17 (3.5)	14 (3.0)	14 (3.1)
Total	24930 (100)	496 (100)	560 (100)	485 (100)	474 (100)	446 (100)

Gender	Year of AIDS Diagnosis (1)					
	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)
Male	403 (90.0)	371 (86.9)	285 (89.3)	263 (90.1)	205 (86.1)	13 (81.3)
Female	34 (7.6)	42 (9.8)	24 (7.5)	16 (5.5)	23 (9.7)	3 (18.8)
Transgender (2)	11 (2.5)	14 (3.3)	10 (3.1)	13 (4.5)	10 (4.2)	0 (0.0)
Total	448 (100)	427 (100)	319 (100)	292 (100)	238 (100)	16 (100)

Table 3. HIV/AIDS Cases by Gender and Year of Initial HIV Diagnosis#, San Francisco, 2007-2012

Gender	Year of Initial HIV Diagnosis (1)					
	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)
Male	461 (87.5)	455 (89.6)	411 (90.3)	389 (90.5)	351 (88.2)	64 (91.4)
Female	44 (8.3)	38 (7.5)	26 (5.7)	32 (7.4)	40 (10.1)	5 (7.1)
Transgender (2)	22 (4.2)	15 (3.0)	18 (4.0)	9 (2.1)	7 (1.8)	1 (1.4)
Total	527 (100)	508 (100)	455 (100)	430 (100)	398 (100)	70 (100)

* Residents of San Francisco at time of HIV/AIDS diagnosis.

Includes persons with a diagnosis of HIV (not AIDS), an initial diagnosis of HIV (not AIDS) and later diagnosed with AIDS, and concurrent diagnosis of HIV and AIDS. The initial year of HIV diagnosis was determined based on the earliest date of HIV antibody test, viral load or CD4 test, initiation of antiretroviral therapy, or patient self-report of a positive HIV test.

(1) Data in recent years are incomplete due to delays in case reporting.

(2) Transgender information was collected since September 1996. Data prior to this are incomplete.

**HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 03/31/2012**

Table 4. AIDS Cases by Transmission Category and Race/Ethnicity, San Francisco San Francisco, 1980-2012

Transmission Category (1)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander No. (%)	Native American No. (%)
Adult/Adolescent					
Gay or bisexual male	16249 (79.5)	1573 (42.6)	2652 (72.9)	702 (76.4)	54 (43.2)
Injection drug user (IDU)	736 (3.6)	1046 (28.3)	252 (6.9)	37 (4.0)	18 (14.4)
Gay or bisexual male IDU	3145 (15.4)	731 (19.8)	507 (13.9)	80 (8.7)	46 (36.8)
Lesbian or bisexual IDU	26 (0.1)	23 (0.6)	7 (0.2)	2 (0.2)	2 (1.6)
Hemophiliac	8 (0.0)	2 (0.1)	5 (0.1)	1 (0.1)	0 (0.0)
Heterosexual (2)	120 (0.6)	203 (5.5)	121 (3.3)	47 (5.1)	4 (3.2)
Transfusion recipient	68 (0.3)	23 (0.6)	27 (0.7)	20 (2.2)	0 (0.0)
Risk not reported/Other (3)	82 (0.4)	76 (2.1)	55 (1.5)	25 (2.7)	1 (0.8)
Pediatric (0-12 years) (4)	8 (0.0)	13 (0.4)	10 (0.3)	5 (0.5)	0 (0.0)
Total	20442 (100)	3690 (100)	3636 (100)	919 (100)	125 (100)

Table 5. HIV Non-AIDS Cases# by Transmission Category and Race/Ethnicity, San Francisco

Transmission Category (1)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander No. (%)	Native American No. (%)
Adult/Adolescent					
Gay or bisexual male	2946 (79.1)	365 (46.6)	749 (76.7)	256 (81.0)	19 (54.3)
Injection drug user (IDU)	145 (3.9)	167 (21.3)	39 (4.0)	4 (1.3)	5 (14.3)
Gay or bisexual male IDU	502 (13.5)	104 (13.3)	95 (9.7)	27 (8.5)	10 (28.6)
Lesbian or bisexual IDU	5 (0.1)	5 (0.6)	4 (0.4)	0 (0.0)	0 (0.0)
Hemophiliac	2 (0.1)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (2)	40 (1.1)	85 (10.9)	46 (4.7)	16 (5.1)	0 (0.0)
Transfusion recipient	0 (0.0)	1 (0.1)	0 (0.0)	0 (0.0)	0 (0.0)
Risk not reported/Other (3)	83 (2.2)	54 (6.9)	38 (3.9)	13 (4.1)	1 (2.9)
Pediatric (0-12 years) (4)	2 (0.1)	2 (0.3)	6 (0.6)	0 (0.0)	0 (0.0)
Total	3725 (100)	783 (100)	977 (100)	316 (100)	35 (100)

* Residents of San Francisco at time of HIV/AIDS diagnosis.

Includes HIV non-AIDS cases reported by name. Excludes HIV non-AIDS cases reported by a non-name code between July 2002 and April 2006 whose names have not been ascertained.

- (1) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
- (2) Includes persons who have had heterosexual contact with a person with HIV/AIDS or with a person who is at risk for HIV.
- (3) Includes persons for whom risk information is incomplete (due to death, refusal to be interviewed or loss to follow-up), cases still under investigation, or interviewed patients who offered no plausible risk for HIV.
- (4) Includes children who have hemophilia or other coagulation disorder, have received a blood transfusion, or who have acquired their infection from an infected mother during the perinatal period.

HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 03/31/2012

Table 6. AIDS Cases by Transmission Category and Year of AIDS Diagnosis, San Francisco, 1980-2012

Transmission Category (2)	Year of AIDS Diagnosis (1)					
	< 2002 No. (%)	2002 No. (%)	2003 No. (%)	2004 No. (%)	2005 No. (%)	2006 No. (%)
Adult/Adolescent						
Gay or bisexual male	18790 (75.4)	303 (61.1)	357 (63.8)	319 (65.8)	293 (61.8)	288 (64.6)
Injection drug user (IDU)	1692 (6.8)	61 (12.3)	70 (12.5)	52 (10.7)	50 (10.5)	38 (8.5)
Gay or bisexual male IDU	3771 (15.1)	106 (21.4)	101 (18.0)	88 (18.1)	96 (20.3)	89 (20.0)
Lesbian or bisexual IDU	45 (0.2)	2 (0.4)	3 (0.5)	1 (0.2)	3 (0.6)	2 (0.4)
Hemophiliac	15 (0.1)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	295 (1.2)	15 (3.0)	22 (3.9)	15 (3.1)	19 (4.0)	20 (4.5)
Transfusion recipient	142 (0.6)	1 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Risk not reported/Other (4)	144 (0.6)	7 (1.4)	6 (1.1)	10 (2.1)	13 (2.7)	9 (2.0)
Pediatric (0-12 years) (5)	36 (0.1)	1 (0.2)	1 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)
Total	24930 (100)	496 (100)	560 (100)	485 (100)	474 (100)	446 (100)

Transmission Category (2)	Year of AIDS Diagnosis (1)					
	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)
Adult/Adolescent						
Gay or bisexual male	286 (63.8)	262 (61.4)	208 (65.2)	168 (57.5)	138 (58.0)	9 (56.3)
Injection drug user (IDU)	40 (8.9)	31 (7.3)	34 (10.7)	24 (8.2)	36 (15.1)	1 (6.3)
Gay or bisexual male IDU	79 (17.6)	87 (20.4)	48 (15.0)	73 (25.0)	41 (17.2)	2 (12.5)
Lesbian or bisexual IDU	0 (0.0)	3 (0.7)	2 (0.6)	0 (0.0)	0 (0.0)	0 (0.0)
Hemophiliac	0 (0.0)	0 (0.0)	1 (0.3)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	32 (7.1)	29 (6.8)	17 (5.3)	16 (5.5)	15 (6.3)	3 (18.8)
Transfusion recipient	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.4)	0 (0.0)
Risk not reported/Other (4)	11 (2.5)	15 (3.5)	9 (2.8)	11 (3.8)	7 (2.9)	1 (6.3)
Pediatric (0-12 years) (5)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Total	448 (100)	427 (100)	319 (100)	292 (100)	238 (100)	16 (100)

* Residents of San Francisco at time of HIV/AIDS diagnosis.

(1) Data in recent years are incomplete due to delays in case reporting.

(2) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.

(3) Includes persons who have had heterosexual contact with a person with HIV/AIDS or with a person who is at risk for HIV.

(4) Includes persons for whom risk information is incomplete (due to death, refusal to be interviewed or loss to follow-up), cases still under investigation, or interviewed patients who offered no plausible risk for HIV.

(5) Includes children who have hemophilia or other coagulation disorder, have received a blood transfusion, or who have acquired their infection from an infected mother during the perinatal period.

**HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 03/31/2012**

Table 7. HIV/AIDS Cases by Transmission Category and Year of Initial HIV Diagnosis#, San Francisco, 2007-2012

Transmission Category (2)	Year of Initial HIV Diagnosis (1)					
	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)
Adult/Adolescent						
Gay or bisexual male	344 (65.3)	371 (73.0)	327 (71.9)	279 (64.9)	282 (70.9)	55 (78.6)
Injection drug user (IDU)	35 (6.6)	27 (5.3)	23 (5.1)	32 (7.4)	25 (6.3)	0 (0.0)
Gay or bisexual male IDU	79 (15.0)	53 (10.4)	64 (14.1)	57 (13.3)	44 (11.1)	6 (8.6)
Lesbian or bisexual IDU	1 (0.2)	3 (0.6)	3 (0.7)	0 (0.0)	0 (0.0)	0 (0.0)
Hemophiliac	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (3)	44 (8.3)	33 (6.5)	16 (3.5)	32 (7.4)	25 (6.3)	5 (7.1)
Transfusion recipient	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Risk not reported/Other (4)	24 (4.6)	21 (4.1)	22 (4.8)	30 (7.0)	22 (5.5)	4 (5.7)
Pediatric (0-12 years) (5)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Total	527 (100)	508 (100)	455 (100)	430 (100)	398 (100)	70 (100)

* Residents of San Francisco at time of HIV/AIDS diagnosis.

Includes persons with a diagnosis of HIV (not AIDS), an initial diagnosis of HIV (not AIDS) and later diagnosed with AIDS, and concurrent diagnosis of HIV and AIDS. The initial year of HIV diagnosis was determined based on the earliest date of HIV antibody test, viral load or CD4 test, initiation of antiretroviral therapy, or patient self-report of a positive HIV test.

- (1) Data in recent years are incomplete due to delays in case reporting.
- (2) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
- (3) Includes persons who have had heterosexual contact with a person with HIV/AIDS or with a person who is at risk for HIV.
- (4) Includes persons for whom risk information is incomplete (due to death, refusal to be interviewed or loss to follow-up), cases still under investigation, or interviewed patients who offered no plausible risk for HIV.
- (5) Includes children who have hemophilia or other coagulation disorder, have received a blood transfusion, or who have acquired their infection from an infected mother during the perinatal period.

HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 03/31/2012

Table 8. AIDS Cases by Gender, Age Group and Race/Ethnicity, San Francisco, 1980-2012

Male Age at AIDS Diagnosis (Years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander No. (%)	Native American No. (%)	Total (1) No. (%)
0 - 12	4 (0.0)	4 (0.1)	5 (0.2)	4 (0.5)	0 (0.0)	19 (0.1)
13 - 19	12 (0.1)	1 (0.0)	14 (0.4)	1 (0.1)	2 (1.9)	30 (0.1)
20 - 24	309 (1.6)	70 (2.3)	130 (3.9)	26 (3.2)	4 (3.7)	546 (2.0)
25 - 29	1671 (8.4)	251 (8.4)	469 (14.1)	91 (11.2)	21 (19.6)	2535 (9.2)
30 - 39	8841 (44.4)	1162 (38.8)	1564 (47.0)	364 (44.7)	52 (48.6)	12109 (44.1)
40 - 49	6569 (33.0)	1003 (33.5)	835 (25.1)	242 (29.7)	24 (22.4)	8753 (31.9)
50 - 59	2021 (10.1)	390 (13.0)	248 (7.5)	64 (7.9)	4 (3.7)	2756 (10.0)
60 +	500 (2.5)	110 (3.7)	61 (1.8)	22 (2.7)	0 (0.0)	702 (2.6)
Male subtotal	19927 (100)	2991 (100)	3326 (100)	814 (100)	107 (100)	27450 (100)
Female Age at AIDS Diagnosis (Years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander No. (%)	Native American No. (%)	Total (1) No. (%)
0 - 12	4 (1.0)	9 (1.6)	5 (2.7)	1 (1.5)	0 (0.0)	19 (1.5)
13 - 19	1 (0.3)	2 (0.4)	2 (1.1)	0 (0.0)	0 (0.0)	5 (0.4)
20 - 24	16 (4.0)	10 (1.8)	10 (5.5)	4 (6.1)	1 (7.1)	42 (3.4)
25 - 29	40 (10.1)	48 (8.5)	26 (14.3)	11 (16.7)	1 (7.1)	128 (10.3)
30 - 39	156 (39.3)	205 (36.4)	60 (33.0)	25 (37.9)	8 (57.1)	461 (37.0)
40 - 49	107 (27.0)	193 (34.3)	47 (25.8)	18 (27.3)	4 (28.6)	375 (30.1)
50 - 59	42 (10.6)	72 (12.8)	18 (9.9)	4 (6.1)	0 (0.0)	140 (11.2)
60 +	31 (7.8)	24 (4.3)	14 (7.7)	3 (4.5)	0 (0.0)	75 (6.0)
Female subtotal	397 (100)	563 (100)	182 (100)	66 (100)	14 (100)	1245 (100)
Transgender (2) Age at AIDS Diagnosis (Years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander Native American No. (%)	Total (1) No. (%)	
13 - 29	25 (21.2)	30 (22.1)	32 (25.0)	12 (27.9)	102 (23.4)	
30 - 39	56 (47.5)	49 (36.0)	61 (47.7)	20 (46.5)	191 (43.8)	
40 +	37 (31.4)	57 (41.9)	35 (27.3)	11 (25.6)	143 (32.8)	
Transgender subtotal	118 (100)	136 (100)	128 (100)	43 (100)	436 (100)	

* Residents of San Francisco at time of HIV/AIDS diagnosis.

(1) Total includes persons with multiple or unknown race.

(2) Transgender information was collected since September 1996. Certain age or race/ethnic groups are combined for transgender cases because of small number.

HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 03/31/2012

Table 9. HIV Non-AIDS Cases# by Gender, Age Group and Race/Ethnicity, San Francisco

Male Age at Initial HIV Diagnosis (Years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander No. (%)	Native American No. (%)	Total (1) No. (%)
0 - 12	1 (0.0)	1 (0.2)	1 (0.1)	0 (0.0)	0 (0.0)	3 (0.1)
13 - 19	25 (0.7)	18 (3.1)	20 (2.3)	8 (2.8)	0 (0.0)	72 (1.3)
20 - 24	303 (8.4)	52 (8.9)	110 (12.8)	30 (10.6)	4 (13.3)	517 (9.4)
25 - 29	543 (15.1)	89 (15.2)	191 (22.2)	65 (23.0)	6 (20.0)	925 (16.9)
30 - 39	1564 (43.6)	196 (33.6)	364 (42.3)	128 (45.2)	15 (50.0)	2325 (42.4)
40 - 49	859 (23.9)	159 (27.2)	148 (17.2)	45 (15.9)	5 (16.7)	1244 (22.7)
50 - 59	241 (6.7)	52 (8.9)	22 (2.6)	7 (2.5)	0 (0.0)	324 (5.9)
60 +	55 (1.5)	17 (2.9)	5 (0.6)	0 (0.0)	0 (0.0)	78 (1.4)
Male subtotal	3591 (100)	584 (100)	861 (100)	283 (100)	30 (100)	5488 (100)

Female Age at Initial HIV Diagnosis (Years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander No. (%)	Native American No. (%)	Total (1) No. (%)
0 - 12	1 (0.9)	1 (0.7)	5 (6.8)	0 (0.0)	0 (0.0)	9 (2.4)
13 - 19	2 (1.9)	2 (1.3)	2 (2.7)	0 (0.0)	0 (0.0)	7 (1.9)
20 - 24	12 (11.1)	13 (8.7)	13 (17.8)	0 (0.0)	0 (0.0)	40 (10.8)
25 - 29	21 (19.4)	15 (10.0)	11 (15.1)	6 (27.3)	0 (0.0)	54 (14.5)
30 - 39	28 (25.9)	47 (31.3)	24 (32.9)	6 (27.3)	3 (75.0)	113 (30.4)
40 - 49	28 (25.9)	51 (34.0)	13 (17.8)	7 (31.8)	0 (0.0)	101 (27.2)
50 - 59	16 (14.8)	17 (11.3)	4 (5.5)	2 (9.1)	1 (25.0)	42 (11.3)
60 +	0 (0.0)	4 (2.7)	1 (1.4)	1 (4.5)	0 (0.0)	6 (1.6)
Female subtotal	108 (100)	150 (100)	73 (100)	22 (100)	4 (100)	372 (100)

Transgender (2) Age at Initial HIV Diagnosis (Years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander Native American No. (%)	Total (1) No. (%)
13 - 29	8 (30.8)	23 (46.9)	18 (41.9)	4 (33.3)	56 (41.2)
30 +	18 (69.2)	26 (53.1)	25 (58.1)	8 (66.7)	80 (58.8)
Transgender subtotal	26 (100)	49 (100)	43 (100)	12 (100)	136 (100)

* Residents of San Francisco at time of HIV/AIDS diagnosis.

Includes HIV non-AIDS cases reported by name. Excludes HIV non-AIDS cases reported by a non-name code between July 2002 and April 2006 whose names have not been ascertained.

(1) Total includes persons with multiple or unknown race.

(2) Transgender information was collected since September 1996. Certain age or race/ethnic groups are combined for transgender cases because of small number.

HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 03/31/2012

Table 10. AIDS Cases by Race/Ethnicity and Year of AIDS Diagnosis, San Francisco, 1980-2012

Race/Ethnicity	Year of AIDS Diagnosis (1)					
	< 2002 No. (%)	2002 No. (%)	2003 No. (%)	2004 No. (%)	2005 No. (%)	2006 No. (%)
White	18167 (72.9)	289 (58.3)	283 (50.5)	261 (53.8)	269 (56.8)	250 (56.1)
African American	2943 (11.8)	88 (17.7)	105 (18.8)	78 (16.1)	83 (17.5)	79 (17.7)
Latino	2832 (11.4)	77 (15.5)	127 (22.7)	112 (23.1)	89 (18.8)	81 (18.2)
Asian/Pacific Islander	671 (2.7)	28 (5.6)	33 (5.9)	22 (4.5)	23 (4.9)	22 (4.9)
Native American	96 (0.4)	2 (0.4)	5 (0.9)	4 (0.8)	3 (0.6)	4 (0.9)
Total (2)	24930 (100)	496 (100)	560 (100)	485 (100)	474 (100)	446 (100)

Race/Ethnicity	Year of AIDS Diagnosis&					
	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)
White	260 (58.0)	221 (51.8)	166 (52.0)	145 (49.7)	125 (52.5)	6 (37.5)
African American	78 (17.4)	82 (19.2)	59 (18.5)	54 (18.5)	40 (16.8)	1 (6.3)
Latino	73 (16.3)	72 (16.9)	65 (20.4)	58 (19.9)	43 (18.1)	7 (43.8)
Asian/Pacific Islander	27 (6.0)	40 (9.4)	13 (4.1)	24 (8.2)	15 (6.3)	1 (6.3)
Native American	0 (0.0)	3 (0.7)	3 (0.9)	2 (0.7)	2 (0.8)	1 (6.3)
Total (2)	448 (100)	427 (100)	319 (100)	292 (100)	238 (100)	16 (100)

Table 11. HIV/AIDS Cases by Race/Ethnicity and Year of Initial HIV Diagnosis#, San Francisco, 2007-2012

Race/Ethnicity	Year of Initial HIV Diagnosis (1)					
	2007 No. (%)	2008 No. (%)	2009 No. (%)	2010 No. (%)	2011 No. (%)	2012 No. (%)
White	273 (51.8)	259 (51.0)	235 (51.6)	209 (48.6)	207 (52.0)	30 (42.9)
African American	78 (14.8)	81 (15.9)	70 (15.4)	61 (14.2)	68 (17.1)	4 (5.7)
Latino	104 (19.7)	113 (22.2)	91 (20.0)	97 (22.6)	76 (19.1)	24 (34.3)
Asian/Pacific Islander	48 (9.1)	40 (7.9)	40 (8.8)	43 (10.0)	34 (8.5)	8 (11.4)
Native American	2 (0.4)	4 (0.8)	1 (0.2)	3 (0.7)	2 (0.5)	2 (2.9)
Total (2)	527 (100)	508 (100)	455 (100)	430 (100)	398 (100)	70 (100)

* Residents of San Francisco at time of HIV/AIDS diagnosis.

Includes persons with a diagnosis of HIV (not AIDS), an initial diagnosis of HIV (not AIDS) and later diagnosed with AIDS, and concurrent diagnosis of HIV and AIDS. The initial year of HIV diagnosis was determined based on the earliest date of HIV antibody test, viral load or CD4 test, initiation of antiretroviral therapy, or patient self-report of a positive HIV test.

(1) Data in recent years are incomplete due to delays in case reporting.

(2) Total includes persons with multiple or unknown race.

HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 03/31/2012

Table 12. AIDS Incidence, Mortality, and Prevalence by Year, San Francisco, 1980-2012

Year	Number of AIDS Cases Reported per Year	Number of AIDS Cases Diagnosed per Year (1)	Number of AIDS Deaths Occurred per Year (1)	Number of Persons Living with AIDS (1)
1980	0	3	0	3
1981	21	26	8	21
1982	75	99	32	88
1983	197	274	111	251
1984	451	558	273	536
1985	673	861	534	863
1986	981	1236	807	1292
1987	1287	1629	878	2043
1988	1408	1763	1039	2767
1989	1584	2161	1276	3652
1990	1686	2046	1366	4332
1991	1685	2287	1509	5110
1992	1638	2331	1641	5800
1993	4268	2071	1600	6271
1994	1914	1789	1595	6465
1995	1632	1565	1483	6547
1996	1239	1083	992	6638
1997	1060	806	423	7021
1998	793	695	402	7314
1999	723	578	353	7539
2000	624	557	349	7747
2001	495	512	324	7935
2002	439	496	321	8110
2003	531	560	302	8368
2004	553	485	309	8544
2005	499	474	312	8706
2006	421	446	290	8862
2007	536	448	270	9040
2008	555	427	230	9237
2009	347	319	210	9346
2010	402	292	192	9446
2011	338	238	144	9540
2012	76	16	31	9525
Total	29131	29131	19606	

* Residents of San Francisco at time of HIV/AIDS diagnosis.

(1) Data in recent years are incomplete due to delays in case/death reporting.

HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 03/31/2012

Table 13. AIDS Cases by Initial AIDS-Defining Condition and Year of AIDS Diagnosis, San Francisco, 1980-2012

Initial AIDS-Defining Condition (1)	Year of AIDS Diagnosis		
	1980-1989 No. (%)	1990-1995 No. (%)	1996-2012 No. (%)
Bacterial infections, recurrent, <13 years	1 (0.0)	5 (0.0)	3 (0.0)
Candidiasis of bronchi, trachea, or lungs	25 (0.3)	13 (0.1)	17 (0.2)
Candidiasis of esophagus	301 (3.5)	307 (2.5)	177 (2.1)
Invasive cervical cancer	1 (0.0)	1 (0.0)	6 (0.1)
Coccidioidomycosis, disseminated or extrapulmonary	8 (0.1)	3 (0.0)	4 (0.0)
Cryptococcosis, extrapulmonary	247 (2.9)	157 (1.3)	91 (1.1)
Cryptosporidiosis, intestinal (>1 mo. duration)	137 (1.6)	203 (1.7)	65 (0.8)
Cytomegalovirus (except liver, spleen, lymph nodes), CMV retinitis with loss of vision	175 (2.0)	139 (1.1)	49 (0.6)
HIV encephalopathy	54 (0.6)	92 (0.8)	21 (0.2)
Herpes simplex: chronic (>1 mo.), bronchitis, pneumonitis, esophagitis	225 (2.6)	196 (1.6)	70 (0.8)
Histoplasmosis	75 (0.9)	53 (0.4)	20 (0.2)
Isosporiasis, intestinal (>1 mo. duration)	14 (0.2)	15 (0.1)	13 (0.2)
Kaposi's sarcoma	10 (0.1)	11 (0.1)	8 (0.1)
Lymphoid interstitial pneumonia/ pulmonary lymphoid hyperplasia, <13 years	1982 (23.0)	869 (7.2)	303 (3.6)
Lymphoma, Burkitt's (non-Hodgkin's)	3 (0.0)	1 (0.0)	0 (0.0)
Lymphoma, immunoblastic (non-Hodgkin's)	59 (0.7)	32 (0.3)	39 (0.5)
Lymphoma, primary in brain	216 (2.5)	163 (1.3)	81 (1.0)
Mycobacterium avium complex or M. kansasii, disseminated or extrapulmonary	27 (0.3)	16 (0.1)	11 (0.1)
Mycobacterium tuberculosis, pulmonary	217 (2.5)	165 (1.4)	53 (0.6)
Mycobacterium tuberculosis, disseminated or extrapulmonary	46 (0.5)	163 (1.3)	97 (1.2)
Mycobacterium other species, disseminated or extrapulmonary	72 (0.8)	93 (0.8)	36 (0.4)
Pneumocystis carinii pneumonia	19 (0.2)	27 (0.2)	8 (0.1)
Pneumonia, recurrent	3794 (44.1)	1809 (15.0)	780 (9.3)
Progressive multifocal leukoencephalopathy	37 (0.4)	144 (1.2)	117 (1.4)
Salmonella sepsis, recurrent	40 (0.5)	15 (0.1)	8 (0.1)
Toxoplasmosis of brain, >1 month of age	10 (0.1)	2 (0.0)	2 (0.0)
Wasting syndrome	178 (2.1)	102 (0.8)	38 (0.5)
CD4 T lymphocyte count <200 or percent <14 (2)	286 (3.3)	448 (3.7)	183 (2.2)
	1217 (14.1)	7489 (61.9)	6399 (75.9)
Total (3)	8610 (100)	12089 (100)	8432 (100)

* Residents of San Francisco at time of HIV/AIDS diagnosis.

(1) Based on AIDS-defining conditions diagnosed in the same calendar month and year.

Cases may have more than one initial AIDS-defining condition.

(2) Cases whose initial AIDS defining condition is low CD4 count (or percent) only.

(3) Total represents unique AIDS cases diagnosed during the time period specified.

**HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 03/31/2012**

Table 14. Cumulative AIDS Indicator Conditions among Persons with AIDS, San Francisco, 1980-2012

AIDS Indicator Condition (1)	Total No. (%)
Bacterial infections, recurrent, <13 years	11 (0.0)
Candidiasis of bronchi, trachea, or lungs	152 (0.5)
Candidiasis of esophagus	2672 (9.2)
Invasive cervical cancer	14 (0.0)
Coccidioidomycosis, disseminated or extrapulmonary	63 (0.2)
Cryptococcosis, extrapulmonary	1888 (6.5)
Cryptosporidiosis, intestinal (>1 mo. duration)	1385 (4.8)
Cytomegalovirus (except liver, spleen, lymph nodes)	2487 (8.5)
CMV retinitis with loss of vision	2503 (8.6)
HIV encephalopathy	2511 (8.6)
Herpes simplex: chronic (>1 mo.), bronchitis, pneumonitis, esophagitis	459 (1.6)
Histoplasmosis	161 (0.6)
Isosporiasis, intestinal (>1 mo. duration)	67 (0.2)
Kaposi's sarcoma	6777 (23.3)
Lymphoid interstitial pneumonia/pulmonary lymphoid hyperplasia, <13 years	6 (0.0)
Lymphoma, Burkitt's (non-Hodgkin's)	558 (1.9)
Lymphoma, immunoblastic (non-Hodgkin's)	1142 (3.9)
Lymphoma, primary in brain	393 (1.3)
Mycobacterium avium complex or M. kansasii, disseminated or extrapulmonary	5108 (17.5)
Mycobacterium tuberculosis, pulmonary	664 (2.3)
Mycobacterium tuberculosis, disseminated or extrapulmonary	485 (1.7)
Mycobacterium other species, disseminated or extrapulmonary	340 (1.2)
Pneumocystis carinii pneumonia	11106 (38.1)
Pneumonia, recurrent	1037 (3.6)
Progressive multifocal leukoencephalopathy	314 (1.1)
Salmonella sepsis, recurrent	60 (0.2)
Toxoplasmosis of brain, >1 month of age	1178 (4.0)
Wasting syndrome	4374 (15.0)

* Residents of San Francisco at time of HIV/AIDS diagnosis.

(1) Includes initial and subsequent diagnoses of AIDS indicator conditions. Cases may have more than one condition.

**HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 03/31/2012**

Table 15. Living Adult/Adolescent HIV/AIDS Cases (>12 years) by Transmission Category, San Francisco

Transmission Category	No.	(%)
Gay or bisexual male	11185	(72.2)
Heterosexual male injection drug user	572	(3.7)
Heterosexual female injection drug user	418	(2.7)
Gay or bisexual male injection drug user	2125	(13.7)
Lesbian or bisexual injection drug user	40	(0.3)
Transgender (1)	341	(2.2)
Hemophiliac	4	(0.0)
Heterosexual contact male (2)	158	(1.0)
Heterosexual contact female (2)	327	(2.1)
Transfusion recipient	17	(0.1)
Risk not reported/Other (3)	298	(1.9)
Total	15485	(100)

Table 16. Living HIV/AIDS Cases by Transmission Category and Race/Ethnicity, San Francisco

Transmission Category	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander No. (%)	Native American No. (%)
Adult/Adolescent					
Gay or bisexual male	7638 (78.8)	916 (44.6)	1961 (75.3)	630 (77.8)	45 (49.5)
Injection drug user (IDU)	365 (3.8)	458 (22.3)	118 (4.5)	20 (2.5)	15 (16.5)
Gay or bisexual male IDU	1445 (14.9)	371 (18.1)	309 (11.9)	78 (9.6)	24 (26.4)
Lesbian or bisexual IDU	16 (0.2)	16 (0.8)	5 (0.2)	1 (0.1)	2 (2.2)
Hemophiliac	3 (0.0)	1 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (2)	98 (1.0)	201 (9.8)	134 (5.1)	47 (5.8)	4 (4.4)
Transfusion recipient	6 (0.1)	3 (0.1)	5 (0.2)	3 (0.4)	0 (0.0)
Risk not reported/Other (3)	120 (1.2)	79 (3.8)	60 (2.3)	25 (3.1)	1 (1.1)
Pediatric (0-12 years) (4)	5 (0.1)	10 (0.5)	13 (0.5)	6 (0.7)	0 (0.0)
Total	9696 (100)	2055 (100)	2605 (100)	810 (100)	91 (100)

* Residents of San Francisco at time of HIV/AIDS diagnosis.

- (1) Transgender information was collected since September 1996. Data prior to this are incomplete.
- (2) Includes persons who have had heterosexual contact with a person with HIV/AIDS or with a person who is at risk for HIV.
- (3) Includes persons for whom risk information is incomplete (due to death, refusal to be interviewed or loss to follow-up), cases still under investigation, or interviewed patients who offered no plausible risk for HIV.
- (4) Includes children who have hemophilia or other coagulation disorder, have received a blood transfusion, or who have acquired their infection from an infected mother during the perinatal period.

HIV/AIDS Quarterly Surveillance Report
Summary of San Francisco Residents with HIV/AIDS*
Reported as of 03/31/2012

Table 17. Living HIV/AIDS Cases by Gender, Age Group and Race/Ethnicity, San Francisco

Male Current Age (Years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander No. (%)	Native American No. (%)	Total (1) No. (%)
0 - 12	0 (0.0)	0 (0.0)	1 (0.0)	0 (0.0)	0 (0.0)	2 (0.0)
13 - 19	1 (0.0)	4 (0.3)	0 (0.0)	2 (0.3)	0 (0.0)	7 (0.0)
20 - 24	38 (0.4)	26 (1.7)	29 (1.2)	9 (1.3)	0 (0.0)	106 (0.7)
25 - 29	160 (1.7)	46 (2.9)	104 (4.5)	38 (5.3)	2 (2.7)	361 (2.5)
30 - 39	862 (9.2)	155 (9.9)	463 (19.9)	160 (22.3)	16 (21.3)	1721 (12.0)
40 - 49	3164 (33.8)	488 (31.3)	916 (39.3)	286 (39.9)	37 (49.3)	4987 (34.9)
50 - 59	3352 (35.8)	579 (37.1)	609 (26.1)	152 (21.2)	18 (24.0)	4758 (33.3)
60 +	1784 (19.1)	263 (16.8)	210 (9.0)	70 (9.8)	2 (2.7)	2344 (16.4)
Male subtotal	9361 (100)	1561 (100)	2332 (100)	717 (100)	75 (100)	14286 (100)
Female Current Age (Years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander No. (%)	Native American No. (%)	Total (1) No. (%)
0 - 12	0 (0.0)	1 (0.3)	1 (0.6)	0 (0.0)	0 (0.0)	2 (0.2)
13 - 19	1 (0.4)	1 (0.3)	6 (3.6)	1 (1.7)	0 (0.0)	11 (1.2)
20 - 24	3 (1.1)	6 (1.6)	3 (1.8)	0 (0.0)	0 (0.0)	12 (1.3)
25 - 29	9 (3.4)	13 (3.5)	14 (8.3)	3 (5.2)	0 (0.0)	40 (4.5)
30 - 39	37 (14.1)	39 (10.4)	29 (17.2)	12 (20.7)	2 (15.4)	126 (14.1)
40 - 49	102 (38.9)	114 (30.4)	52 (30.8)	22 (37.9)	5 (38.5)	299 (33.4)
50 - 59	84 (32.1)	149 (39.7)	39 (23.1)	15 (25.9)	6 (46.2)	298 (33.3)
60 +	26 (9.9)	52 (13.9)	25 (14.8)	5 (8.6)	0 (0.0)	108 (12.1)
Female subtotal	262 (100)	375 (100)	169 (100)	58 (100)	13 (100)	896 (100)
Transgender (2) Current Age (Years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/ Pacific Islander Native American No. (%)	Total (1) No. (%)	
13 - 39	17 (23.3)	36 (30.3)	43 (41.3)	11 (28.9)	113 (33.1)	
40 +	56 (76.7)	83 (69.7)	61 (58.7)	27 (71.1)	228 (66.9)	
Transgender subtotal	73 (100)	119 (100)	104 (100)	38 (100)	341 (100)	

* Residents of San Francisco at time of HIV/AIDS diagnosis.

(1) Total includes persons with multiple or unknown race.

(2) Transgender information was collected since September 1996. Certain age or race/ethnic groups are combined for transgender cases because of small number.