

QUARTERLY AIDS SURVEILLANCE REPORT

San Francisco Department of Public Health

AIDS Cases Reported Through September 2005

Contents	Page
Surveillance Summary.....	1
Table 1: Adult/Adolescent AIDS Cases by Transmission Category.....	2
Table 2: AIDS Cases by Gender and Year of Diagnosis.....	2
Table 3: AIDS Cases by Transmission Category and Race/Ethnicity.....	3
Table 4: AIDS Cases by Transmission Category and Asian/Pacific Islander Ethnicity.....	3
Table 5: AIDS Cases by Transmission Category and Year of Diagnosis.....	4
Table 6: AIDS Cases by Gender, Age Group and Race/Ethnicity.....	5
Table 7: AIDS Cases by Race/Ethnicity and Year of Diagnosis.....	6
Table 8: AIDS Cases and Cumulative Rates per 100,000 by Race/Ethnicity and Gender.....	6
Table 9: AIDS Incidence, Mortality, and Prevalence by Year.....	7
Table 10: Cases by Initial AIDS-Defining Condition.....	8
Table 11: Cumulative AIDS Indicator Conditions among Persons with AIDS.....	9
Table 12: Living Adult/Adolescent AIDS Cases by Transmission Category.....	10
Table 13: Living AIDS Cases by Transmission Category and Race/Ethnicity.....	10
Table 14: Living AIDS Cases by Gender, Age Group and Race/Ethnicity.....	11
Table 15: Living AIDS Cases by Initial AIDS-Defining Condition.....	12
Table 16: Cumulative AIDS Indicator Conditions among Persons Living with AIDS.....	13

The AIDS Surveillance Report is published quarterly by the San Francisco Department of Public Health
HIV/AIDS Statistics, Epidemiology & Intervention Research Section
25 Van Ness Avenue, Suite 500, San Francisco, CA 94102; Phone (415) 554-9050, FAX (415) 431-0353
Director of Health: Mitchell Katz, MD; Section Director: Sandra Schwarcz, MD, MPH
Section Assistant Director: Ling Hsu, MPH; Program Coordinators: Maree Kay Parisi, Viva Delgado
Epidemiologists: Susan Scheer, PhD, Priscilla Lee Chu, MPH, Sharon Pipkin, MPH, Hilary Spindler, MPH

The AIDS Surveillance Report is accessible via internet: www.dph.sf.ca.us/PHP/AIDSSurvUnit.htm

October 12, 2005


Dear Colleague,

We want to call your attention to a recent change in how we present the San Francisco AIDS surveillance data. Beginning in 2005, publications of our data were changed to include only persons who were *residents* of San Francisco at the time they were diagnosed with AIDS. Prior to this change, our reports also included persons who were diagnosed with AIDS at a San Francisco medical facility, but who were residents elsewhere at the time of their diagnosis.

We have decided to exclude non-San Francisco residents from the data presentations because AIDS case reporting practices result in our having a complete count of San Francisco residents with AIDS, but an incomplete record of non-San Francisco residents with AIDS. In order to provide consistent AIDS surveillance data, our surveillance quarterly and annual reports will include only reports of persons diagnosed with HIV or AIDS who were residents of San Francisco at the time of their diagnosis. Consequently, the numbers in reports from 2005 and on will differ from reports published prior to 2005.

You may contact the HIV/AIDS Statistics, Epidemiology, and Intervention Research Section of the San Francisco Department of Public Health at (415) 554-9050 if you have questions concerning our reports.

Sincerely,

A handwritten signature in black ink, appearing to read "Sandy Schwarcz". The signature is fluid and cursive, written over a faint, dotted grid background.

Sandy Schwarcz, M.D., M.P.H.
Director of HIV/AIDS Surveillance

A handwritten signature in black ink, appearing to read "Ling Hsu". The signature is fluid and cursive, written over a faint, dotted grid background.

Ling Hsu, M.P.H.
Assistant Director

AIDS Surveillance Summary

San Francisco (as of 09/30/05)

Cumulative cases ¹ :	26,410
Cumulative deaths:	17,788


California (as of 08/31/05)

Cumulative cases:	138,198
Cumulative deaths:	80,439

United States² (as of 12/31/03)

Cumulative cases:	929,985
Cumulative deaths:	524,060

San Francisco AIDS Incidence, Mortality, and Prevalence by Year, 1980-2005³



1. Includes SF residents diagnosed in SF and SF residents diagnosed in other jurisdictions. Excludes persons diagnosed in SF who resided in other jurisdictions at the time of their AIDS diagnosis.
2. The US numbers do not represent actual cases or deaths reported. Rather, these numbers are estimated and adjusted for reporting delays.
3. Reporting for recent year is incomplete. See Table 9 for actual numbers per year.

Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report
 Summary of San Francisco Residents with AIDS*
 Reported as of 09/30/2005

Table 1. Adult/Adolescent AIDS Cases (>12 years) by Transmission Category, San Francisco, 1980-2005

Transmission Category	No.	(%)
Gay or bisexual male	19723	(74.8)
Heterosexual male injection drug user	1317	(5.0)
Heterosexual female injection drug user	623	(2.4)
Gay or bisexual male injection drug user	3627	(13.8)
Lesbian or bisexual injection drug user	42	(0.2)
Transgender (1)	326	(1.2)
Hemophiliac	17	(0.1)
Heterosexual contact male (2)	89	(0.3)
Heterosexual contact female (2)	251	(1.0)
Transfusion recipient	143	(0.5)
None of the above/Other (3)	214	(0.8)
Total	26372	(100)

Table 2. AIDS Cases by Gender and Year of Diagnosis, San Francisco, 1980-2005

Gender	Year of Diagnosis					
	< 1995 No. (%)	1995 No. (%)	1996 No. (%)	1997 No. (%)	1998 No. (%)	1999 No. (%)
Male	18595 (97.2)	1488 (95.3)	1003 (93.0)	739 (91.6)	623 (90.4)	512 (88.6)
Female	528 (2.8)	73 (4.7)	75 (7.0)	49 (6.1)	50 (7.3)	48 (8.3)
Transgender (1)	--	--	--	19 (2.4)	16 (2.3)	18 (3.1)
Total	19123 (100)	1561 (100)	1078 (100)	807 (100)	689 (100)	578 (100)

Gender	Year of Diagnosis					
	2000 No. (%)	2001 No. (%)	2002 No. (%)	2003 No. (%)	2004 No. (%)	2005 No. (%)
Male	476 (86.9)	443 (88.8)	425 (89.9)	432 (88.2)	338 (88.3)	160 (88.4)
Female	51 (9.3)	43 (8.6)	32 (6.8)	37 (7.6)	32 (8.4)	20 (11.0)
Transgender (1)	21 (3.8)	13 (2.6)	16 (3.4)	21 (4.3)	13 (3.4)	1 (0.6)
Total	548 (100)	499 (100)	473 (100)	490 (100)	383 (100)	181 (100)

* Residents of San Francisco at time of initial AIDS diagnosis.

(1) Transgender information was collected since September 1996. Data prior to this are incomplete.

(2) Includes persons who have had heterosexual contact with a person with AIDS or with a person who is at risk for HIV.

(3) Includes persons for whom risk information is incomplete (due to death, refusal to be interviewed or loss to follow-up), cases still under investigation, or interviewed patients who offered no plausible risk for HIV.

Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report
Summary of San Francisco Residents with AIDS*
Reported as of 09/30/2005

Table 3. AIDS Cases by Transmission Category and Race/Ethnicity, San Francisco, 1980-2005

Transmission Category (1)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/Pacific Islander No. (%)	Native American No. (%)
Adult/Adolescent					
Gay or bisexual male	15464 (80.9)	1469 (44.4)	2230 (73.9)	621 (77.4)	64 (47.1)
Injection drug user (IDU)	679 (3.6)	983 (29.7)	220 (7.3)	40 (5.0)	20 (14.7)
Gay or bisexual male IDU	2704 (14.1)	604 (18.3)	385 (12.8)	58 (7.2)	47 (34.6)
Lesbian or bisexual IDU	20 (0.1)	14 (0.4)	5 (0.2)	2 (0.2)	1 (0.7)
Hemophiliac	8 (0.0)	2 (0.1)	6 (0.2)	1 (0.1)	0 (0.0)
Heterosexual (2)	108 (0.6)	130 (3.9)	73 (2.4)	29 (3.6)	2 (1.5)
Transfusion recipient	69 (0.4)	26 (0.8)	26 (0.9)	22 (2.7)	0 (0.0)
None of the above/Other (3)	59 (0.3)	67 (2.0)	61 (2.0)	24 (3.0)	1 (0.7)
Pediatric (0-12 years) (4)					
	8 (0.0)	13 (0.4)	10 (0.3)	5 (0.6)	1 (0.7)
Total	19119 (100)	3308 (100)	3016 (100)	802 (100)	136 (100)

Table 4. AIDS Cases by Transmission Category and Asian/Pacific Islander Ethnicity, San Francisco, 1980-2005

Transmission Category (1)	Chinese No. (%)	Japanese No. (%)	Filipino No. (%)	Southeast Asian No. (%)	Korean No. (%)	Pacific Islander No. (%)	Other No. (%)
Adult/Adolescent							
Gay or bisexual male	149 (79.7)	80 (84.2)	207 (80.5)	55 (74.3)	9 (75.0)	42 (65.6)	79 (69.9)
Injection drug user (IDU)	7 (3.7)	0 (0.0)	9 (3.5)	3 (4.1)	2 (16.7)	9 (14.1)	10 (8.8)
Gay or bisexual male IDU	7 (3.7)	11 (11.6)	18 (7.0)	6 (8.1)	1 (8.3)	6 (9.4)	9 (8.0)
Lesbian or bisexual IDU	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	2 (3.1)	0 (0.0)
Hemophiliac	0 (0.0)	1 (1.1)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (2)	4 (2.1)	1 (1.1)	11 (4.3)	4 (5.4)	0 (0.0)	4 (6.3)	5 (4.4)
Transfusion recipient	11 (5.9)	1 (1.1)	7 (2.7)	3 (4.1)	0 (0.0)	0 (0.0)	0 (0.0)
None of the above/Other (3)	7 (3.7)	1 (1.1)	4 (1.6)	3 (4.1)	0 (0.0)	1 (1.6)	8 (7.1)
Pediatric (0-12 years) (4)							
	2 (1.1)	0 (0.0)	1 (0.4)	0 (0.0)	0 (0.0)	0 (0.0)	2 (1.8)
Total	187 (100)	95 (100)	257 (100)	74 (100)	12 (100)	64 (100)	113 (100)

* Residents of San Francisco at time of initial AIDS diagnosis.

(1) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.

(2) Includes persons who have had heterosexual contact with a person with AIDS or with a person who is at risk for HIV.

(3) Includes persons for whom risk information is incomplete (due to death, refusal to be interviewed or loss to follow-up), cases still under investigation, or interviewed patients who offered no plausible risk for HIV.

(4) Includes children who have hemophilia or other coagulation disorder, have received a blood transfusion, or who have acquired their infection from an infected mother during the perinatal period.

Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report
 Summary of San Francisco Residents with AIDS*
 Reported as of 09/30/2005

Table 5. AIDS Cases by Transmission Category and Year of Diagnosis, San Francisco, 1980-2005

Transmission Category (1)	Year of Diagnosis					
	< 1995 No. (%)	1995 No. (%)	1996 No. (%)	1997 No. (%)	1998 No. (%)	1999 No. (%)
Adult/Adolescent						
Gay or bisexual male	15022 (78.6)	1120 (71.7)	724 (67.2)	556 (68.9)	446 (64.7)	361 (62.5)
Injection drug user (IDU)	1001 (5.2)	151 (9.7)	144 (13.4)	108 (13.4)	92 (13.4)	83 (14.4)
Gay or bisexual male IDU	2667 (13.9)	242 (15.5)	166 (15.4)	110 (13.6)	126 (18.3)	99 (17.1)
Lesbian or bisexual IDU	26 (0.1)	4 (0.3)	0 (0.0)	2 (0.2)	2 (0.3)	1 (0.2)
Hemophiliac	12 (0.1)	4 (0.3)	1 (0.1)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (2)	163 (0.9)	20 (1.3)	25 (2.3)	16 (2.0)	14 (2.0)	17 (2.9)
Transfusion recipient	128 (0.7)	6 (0.4)	4 (0.4)	2 (0.2)	0 (0.0)	1 (0.2)
None of the above/Other (3)	76 (0.4)	11 (0.7)	12 (1.1)	12 (1.5)	9 (1.3)	15 (2.6)
Pediatric (0-12 years) (4)	28 (0.1)	3 (0.2)	2 (0.2)	1 (0.1)	0 (0.0)	1 (0.2)
Total	19123 (100)	1561 (100)	1078 (100)	807 (100)	689 (100)	578 (100)

Transmission Category (1)	Year of Diagnosis					
	2000 No. (%)	2001 No. (%)	2002 No. (%)	2003 No. (%)	2004 No. (%)	2005 No. (%)
Adult/Adolescent						
Gay or bisexual male	337 (61.5)	309 (61.9)	307 (64.9)	320 (65.3)	255 (66.6)	109 (60.2)
Injection drug user (IDU)	87 (15.9)	68 (13.6)	58 (12.3)	80 (16.3)	47 (12.3)	25 (13.8)
Gay or bisexual male IDU	93 (17.0)	79 (15.8)	82 (17.3)	59 (12.0)	57 (14.9)	24 (13.3)
Lesbian or bisexual IDU	3 (0.5)	3 (0.6)	0 (0.0)	1 (0.2)	0 (0.0)	0 (0.0)
Hemophiliac	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (2)	18 (3.3)	16 (3.2)	14 (3.0)	16 (3.3)	13 (3.4)	10 (5.5)
Transfusion recipient	0 (0.0)	1 (0.2)	1 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)
None of the above/Other (3)	9 (1.6)	23 (4.6)	10 (2.1)	13 (2.7)	11 (2.9)	13 (7.2)
Pediatric (0-12 years) (4)	1 (0.2)	0 (0.0)	1 (0.2)	1 (0.2)	0 (0.0)	0 (0.0)
Total	548 (100)	499 (100)	473 (100)	490 (100)	383 (100)	181 (100)

* Residents of San Francisco at time of initial AIDS diagnosis.

- (1) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
- (2) Includes persons who have had heterosexual contact with a person with AIDS or with a person who is at risk for HIV.
- (3) Includes persons for whom risk information is incomplete (due to death, refusal to be interviewed or loss to follow-up), cases still under investigation, or interviewed patients who offered no plausible risk for HIV.
- (4) Includes children who have hemophilia or other coagulation disorder, have received a blood transfusion, or who have acquired their infection from an infected mother during the perinatal period.

Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report
Summary of San Francisco Residents with AIDS*
Reported as of 09/30/2005

Table 6. AIDS Cases by Gender, Age Group and Race/Ethnicity, San Francisco, 1980-2005

Male						
Age (years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/Pacific Islander No. (%)	Native American No. (%)	Total (1) No. (%)
0 - 12	4 (0.0)	4 (0.1)	5 (0.2)	4 (0.6)	1 (0.8)	19 (0.1)
13 - 19	9 (0.0)	0 (0.0)	14 (0.5)	1 (0.1)	2 (1.7)	26 (0.1)
20 - 24	290 (1.6)	62 (2.3)	88 (3.2)	21 (2.9)	4 (3.3)	466 (1.9)
25 - 29	1607 (8.6)	245 (9.0)	400 (14.4)	81 (11.3)	22 (18.3)	2357 (9.4)
30 - 39	8535 (45.7)	1141 (41.9)	1361 (49.0)	326 (45.6)	61 (50.8)	11433 (45.6)
40 - 49	6045 (32.4)	877 (32.2)	666 (24.0)	213 (29.8)	24 (20.0)	7833 (31.3)
50 - 59	1761 (9.4)	315 (11.6)	195 (7.0)	52 (7.3)	5 (4.2)	2331 (9.3)
60 +	431 (2.3)	82 (3.0)	51 (1.8)	17 (2.4)	1 (0.8)	582 (2.3)
Male subtotal	18682 (100)	2726 (100)	2780 (100)	715 (100)	120 (100)	25047 (100)
Female						
Age (years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/Pacific Islander No. (%)	Native American No. (%)	Total (1) No. (%)
0 - 12	4 (1.2)	9 (1.9)	5 (3.6)	1 (1.7)	0 (0.0)	19 (1.8)
13 - 19	1 (0.3)	1 (0.2)	2 (1.4)	0 (0.0)	0 (0.0)	4 (0.4)
20 - 24	15 (4.4)	9 (1.9)	9 (6.4)	4 (6.8)	1 (8.3)	38 (3.7)
25 - 29	40 (11.6)	41 (8.6)	21 (15.0)	10 (16.9)	0 (0.0)	113 (10.9)
30 - 39	141 (41.0)	195 (40.7)	46 (32.9)	17 (28.8)	8 (66.7)	409 (39.4)
40 - 49	87 (25.3)	159 (33.2)	36 (25.7)	18 (30.5)	3 (25.0)	303 (29.2)
50 - 59	29 (8.4)	44 (9.2)	13 (9.3)	6 (10.2)	0 (0.0)	92 (8.9)
60 +	27 (7.8)	21 (4.4)	8 (5.7)	3 (5.1)	0 (0.0)	59 (5.7)
Female subtotal	344 (100)	479 (100)	140 (100)	59 (100)	12 (100)	1037 (100)
Transgender (2)						
Age (years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/Pacific Islander Native American No. (%)	Total (1) No. (%)	
13 - 29	21 (22.6)	24 (23.3)	25 (26.0)	9 (28.1)	79 (24.2)	
30 - 39	46 (49.5)	37 (35.9)	48 (50.0)	15 (46.9)	147 (45.1)	
40 +	26 (28.0)	42 (40.8)	23 (24.0)	8 (25.0)	100 (30.7)	
Transgender subtotal	93 (100)	103 (100)	96 (100)	32 (100)	326 (100)	

* Residents of San Francisco at time of initial AIDS diagnosis.

(1) Total includes persons with multiple or unknown race.

(2) Transgender information was collected since September 1996. Certain age or race/ethnic groups are combined for transgender cases because of small number.

Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report
 Summary of San Francisco Residents with AIDS*
 Reported as of 09/30/2005

Table 7. AIDS Cases by Race/Ethnicity and Year of Diagnosis, San Francisco, 1980-2005

Race/Ethnicity	Year of Diagnosis					
	< 1995 No. (%)	1995 No. (%)	1996 No. (%)	1997 No. (%)	1998 No. (%)	1999 No. (%)
White	14623 (76.5)	1076 (68.9)	673 (62.4)	516 (63.9)	442 (64.2)	340 (58.8)
African American	2014 (10.5)	227 (14.5)	194 (18.0)	146 (18.1)	140 (20.3)	104 (18.0)
Latino	1916 (10.0)	191 (12.2)	160 (14.8)	110 (13.6)	77 (11.2)	103 (17.8)
Asian/Pacific Islander	475 (2.5)	60 (3.8)	42 (3.9)	30 (3.7)	27 (3.9)	25 (4.3)
Native American	89 (0.5)	5 (0.3)	8 (0.7)	4 (0.5)	2 (0.3)	5 (0.9)
Total (1)	19123 (100)	1561 (100)	1078 (100)	807 (100)	689 (100)	578 (100)

Race/Ethnicity	Year of Diagnosis					
	2000 No. (%)	2001 No. (%)	2002 No. (%)	2003 No. (%)	2004 No. (%)	2005 No. (%)
White	318 (58.0)	290 (58.1)	283 (59.8)	246 (50.2)	215 (56.1)	97 (53.6)
African American	110 (20.1)	99 (19.8)	82 (17.3)	98 (20.0)	64 (16.7)	30 (16.6)
Latino	87 (15.9)	70 (14.0)	72 (15.2)	108 (22.0)	84 (21.9)	38 (21.0)
Asian/Pacific Islander	27 (4.9)	32 (6.4)	28 (5.9)	28 (5.7)	13 (3.4)	15 (8.3)
Native American	4 (0.7)	5 (1.0)	3 (0.6)	6 (1.2)	5 (1.3)	0 (0.0)
Total (1)	548 (100)	499 (100)	473 (100)	490 (100)	383 (100)	181 (100)

Table 8. AIDS Cases and Cumulative Rates per 100,000 by Race/Ethnicity and Gender, San Francisco

Race/Ethnicity	Male		Female		Total (2)	
	No.	(Rate)	No.	(Rate)	No.	(Rate)
White	18682	(10795.5)	344	(209.7)	19026	(5643.7)
African American	2726	(7313.2)	479	(1226.1)	3205	(4198.2)
Latino	2780	(5417.7)	140	(283.4)	2920	(2899.2)
Asian/Pacific Islander	715	(724.2)	59	(55.2)	774	(376.3)
Native American	120	(8733.6)	12	(951.6)	132	(5009.5)
Total (1)	25047	(6909.6)	1037	(286.9)	26084	(3603.0)

* Residents of San Francisco at time of initial AIDS diagnosis.

(1) Total includes persons with multiple or unknown race.

(2) Transgender cases were excluded because population size for transgender can not be determined for rates calculation.

Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report
 Summary of San Francisco Residents with AIDS*
 Reported as of 09/30/2005

Table 9. AIDS Incidence, Mortality, and Prevalence by Year, San Francisco, 1980-2005

Year	Number of Cases Reported per Year	Number of Cases Diagnosed per Year (1)	Number of Deaths Occurred per Year (1)	Number of Persons Living with AIDS (1)
1980	0	3	0	3
1981	21	26	8	21
1982	75	99	32	88
1983	197	274	111	251
1984	451	557	272	536
1985	673	859	531	864
1986	981	1236	807	1293
1987	1287	1629	876	2046
1988	1408	1762	1036	2772
1989	1585	2161	1273	3660
1990	1687	2048	1362	4346
1991	1685	2280	1495	5131
1992	1639	2327	1633	5825
1993	4275	2077	1584	6318
1994	1917	1785	1584	6519
1995	1636	1561	1475	6605
1996	1247	1078	982	6701
1997	1062	807	413	7095
1998	795	689	396	7388
1999	727	578	352	7614
2000	630	548	340	7822
2001	500	499	319	8002
2002	441	473	318	8157
2003	536	490	219	8428
2004	565	383	219	8592
2005	390	181	151	8622
Total	26410	26410	17788	

* Residents of San Francisco at time of initial AIDS diagnosis.

(1) Data in recent years is incomplete due to delay in cases/deaths reporting.

Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report
 Summary of San Francisco Residents with AIDS*
 Reported as of 09/30/2005

Table 10. Cases by Initial AIDS-Defining Condition, San Francisco, 1980-2005

Initial AIDS-Defining Condition	Def. (1) No.	Pres. (2) No.	Total No. (%)
Bacterial infections, recurrent, < 13 years [HIV+(3)]	8	N/A(4)	8 (0.0)
Candidiasis of bronchi, trachea, or lungs	24	N/A	24 (0.1)
Candidiasis of esophagus	290	217	507 (1.9)
Cervical cancer, invasive [HIV+]	5	N/A	5 (0.0)
Coccidioidomycosis, disseminated or extrapulmonary [HIV+]	7	N/A	7 (0.0)
Cryptococcosis, extrapulmonary	365	N/A	365 (1.4)
Cryptosporidiosis, intestinal (> 1 mo. duration)	328	N/A	328 (1.2)
Cytomegalovirus (except liver, spleen, lymph nodes), > 1 month of age	202	N/A	202 (0.8)
CMV retinitis with loss of vision [HIV+]	0	112	112 (0.4)
HIV encephalopathy [HIV+]	380	N/A	380 (1.4)
Herpes simplex: chronic (> 1 mo. duration), bronchitis, pneumonitis, esophagitis	96	N/A	96 (0.4)
Histoplasmosis, disseminated or extrapulmonary [HIV+]	25	N/A	25 (0.1)
Isosporiasis, intestinal (> 1 mo. duration) [HIV+]	21	N/A	21 (0.1)
Kaposi's sarcoma (5)	2516	273	2789 (10.6)
Lymphoid interstitial pneumonia/pulmonary lymphoid hyperplasia, < 13 years	4	0	4 (0.0)
Lymphoma, Burkitt's (non-Hodgkin's) [HIV+]	98	N/A	98 (0.4)
Lymphoma, immunoblastic (non-Hodgkin's) [HIV+]	409	N/A	409 (1.5)
Lymphoma, primary in brain (5)	39	N/A	39 (0.1)
Mycobacterium avium complex or M. kansasii, disseminated or extrapulmonary	254	3	257 (1.0)
Mycobacterium tuberculosis, pulmonary [HIV+]	191	13	204 (0.8)
Mycobacterium tuberculosis, disseminated or extrapulmonary [HIV+]	136	3	139 (0.5)
Mycobacterium other species, disseminated or extrapulmonary [HIV+]	16	11	27 (0.1)
Pneumocystis carinii pneumonia	5048	578	5626 (21.3)
Pneumonia, recurrent [HIV+]	217	23	240 (0.9)
Progressive multifocal leukoencephalopathy	50	N/A	50 (0.2)
Salmonella sepsis, recurrent [HIV+]	8	N/A	8 (0.0)
Toxoplasmosis of brain, > 1 month of age	44	199	243 (0.9)
Wasting syndrome [HIV+]	667	N/A	667 (2.5)
CD4 T lymphocyte count < 200 or percent < 14 [HIV+]	13526	N/A	13526 (51.2)
Any AIDS indicator condition and HIV-negative and CD4 count < 400 (6)	4	N/A	4 (0.0)
Total	24978	1432	26410 (100)

* Residents of San Francisco at time of initial AIDS diagnosis.

- (1) Indicator conditions diagnosed definitively (e.g. culture or biopsy proven).
- (2) Indicator conditions diagnosed presumptively in a person who has laboratory evidence of HIV infection.
- (3) [HIV+]: Indicator conditions that require laboratory evidence of HIV infection.
- (4) N/A: Conditions which require definitive diagnoses only.
- (5) Laboratory evidence of HIV infection in persons > 60 years of age.
- (6) In the absence of other causes of immunocompromise.

Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report
 Summary of San Francisco Residents with AIDS*
 Reported as of 09/30/2005

Table 11. Cumulative AIDS Indicator Conditions among Persons with AIDS, San Francisco, 1980-2005

AIDS Indicator Condition (1)	Total No. (%)
Bacterial infections, recurrent, < 13 years [HIV+(2)]	11 (0.0)
Candidiasis of bronchi, trachea, or lungs	130 (0.5)
Candidiasis of esophagus	2387 (9.0)
Cervical cancer, invasive [HIV+]	9 (0.0)
Coccidioidomycosis, disseminated or extrapulmonary [HIV+]	54 (0.2)
Cryptococcosis, extrapulmonary	1756 (6.6)
Cryptosporidiosis, intestinal (> 1 mo. duration)	1340 (5.1)
Cytomegalovirus (except liver, spleen, lymph nodes), > 1 month of age	2388 (9.0)
CMV retinitis with loss of vision [HIV+]	2442 (9.2)
HIV encephalopathy [HIV+]	2420 (9.2)
Herpes simplex: chronic (> 1 mo. duration), bronchitis, pneumonitis, esophagitis	404 (1.5)
Histoplasmosis, disseminated or extrapulmonary [HIV+]	151 (0.6)
Isosporiasis, intestinal (> 1 mo. duration) [HIV+]	66 (0.2)
Kaposi's sarcoma (3)	6560 (24.8)
Lymphoid interstitial pneumonia/pulmonary lymphoid hyperplasia, < 13 years	5 (0.0)
Lymphoma, Burkitt's (non-Hodgkin's) [HIV+]	514 (1.9)
Lymphoma, immunoblastic (non-Hodgkin's) [HIV+]	1088 (4.1)
Lymphoma, primary in brain (3)	377 (1.4)
Mycobacterium avium complex or M. kansasii, disseminated or extrapulmonary	4992 (18.9)
Mycobacterium tuberculosis, pulmonary [HIV+]	612 (2.3)
Mycobacterium tuberculosis, disseminated or extrapulmonary [HIV+]	446 (1.7)
Mycobacterium other species, disseminated or extrapulmonary [HIV+]	323 (1.2)
Pneumocystis carinii pneumonia	10679 (40.4)
Pneumonia, recurrent [HIV+]	968 (3.7)
Progressive multifocal leukoencephalopathy	299 (1.1)
Salmonella sepsis, recurrent [HIV+]	57 (0.2)
Toxoplasmosis of brain, > 1 month of age	1138 (4.3)
Wasting syndrome [HIV+]	4197 (15.9)

* Residents of San Francisco at time of initial AIDS diagnosis.

(1) Cases may have more than one condition.

(2) [HIV+]: Indicator conditions that require laboratory evidence of HIV infection.

(3) Laboratory evidence of HIV infection in persons > 60 years of age.

Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report
 Summary of San Francisco Residents with AIDS*
 Reported as of 09/30/2005

Table 12. Living Adult/Adolescent AIDS Cases (>12 years) by Transmission Category, San Francisco

Transmission Category	No.	(%)
Gay or bisexual male	6208	(72.1)
Heterosexual male injection drug user	493	(5.7)
Heterosexual female injection drug user	294	(3.4)
Gay or bisexual male injection drug user	1095	(12.7)
Lesbian or bisexual injection drug user	19	(0.2)
Transgender (1)	173	(2.0)
Hemophiliac	5	(0.1)
Heterosexual contact male (2)	46	(0.5)
Heterosexual contact female (2)	136	(1.6)
Transfusion recipient	22	(0.3)
None of the above/Other (3)	117	(1.4)
Total	8608	(100)

Table 13. Living AIDS Cases by Transmission Category and Race/Ethnicity, San Francisco

Transmission Category	White	African	Latino	Asian/Pacific	Native
	No. (%)	American No. (%)	No. (%)	Islander No. (%)	American No. (%)
Adult/Adolescent					
Gay or bisexual male	4527 (79.6)	493 (40.9)	951 (74.2)	285 (75.0)	23 (45.1)
Injection drug user (IDU)	292 (5.1)	379 (31.5)	87 (6.8)	19 (5.0)	10 (19.6)
Gay or bisexual male IDU	770 (13.5)	213 (17.7)	152 (11.9)	31 (8.2)	16 (31.4)
Lesbian or bisexual IDU	9 (0.2)	8 (0.7)	1 (0.1)	1 (0.3)	0 (0.0)
Hemophiliac	3 (0.1)	2 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)
Heterosexual (2)	50 (0.9)	64 (5.3)	50 (3.9)	18 (4.7)	1 (2.0)
Transfusion recipient	8 (0.1)	3 (0.2)	5 (0.4)	6 (1.6)	0 (0.0)
None of the above/Other (3)	28 (0.5)	39 (3.2)	31 (2.4)	17 (4.5)	0 (0.0)
Pediatric (0-12 years) (4)	0 (0.0)	4 (0.3)	5 (0.4)	3 (0.8)	1 (2.0)
Total	5687 (100)	1205 (100)	1282 (100)	380 (100)	51 (100)

* Residents of San Francisco at time of initial AIDS diagnosis.

(1) Transgender information was collected since September 1996. Data prior to this are incomplete.

(2) Includes persons who have had heterosexual contact with a person with AIDS or with a person who is at risk for HIV.

(3) Includes persons for whom risk information is incomplete (due to death, refusal to be interviewed or loss to follow-up), cases still under investigation, or interviewed patients who offered no plausible risk for HIV.

(4) Includes children who have hemophilia or other coagulation disorder, have received a blood transfusion, or who have acquired their infection from an infected mother during the perinatal period.

Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report
Summary of San Francisco Residents with AIDS*
Reported as of 09/30/2005

Table 14. Living AIDS Cases by Gender, Age Group and Race/Ethnicity, San Francisco

Male						
Age (years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/Pacific Islander No. (%)	Native American No. (%)	Total (1) No. (%)
0 - 12	0 (0.0)	1 (0.1)	1 (0.1)	2 (0.6)	1 (2.3)	6 (0.1)
13 - 19	4 (0.1)	0 (0.0)	7 (0.6)	1 (0.3)	2 (4.7)	14 (0.2)
20 - 24	101 (1.8)	32 (3.4)	44 (3.8)	15 (4.6)	3 (7.0)	196 (2.5)
25 - 29	439 (8.0)	77 (8.3)	184 (16.0)	39 (11.9)	7 (16.3)	747 (9.4)
30 - 39	2538 (46.2)	380 (40.8)	567 (49.4)	148 (45.0)	25 (58.1)	3665 (46.1)
40 - 49	1778 (32.4)	308 (33.0)	261 (22.7)	104 (31.6)	5 (11.6)	2459 (30.9)
50 - 59	538 (9.8)	114 (12.2)	68 (5.9)	14 (4.3)	0 (0.0)	735 (9.2)
60 +	90 (1.6)	20 (2.1)	16 (1.4)	6 (1.8)	0 (0.0)	132 (1.7)
Male subtotal	5488 (100)	932 (100)	1148 (100)	329 (100)	43 (100)	7954 (100)
Female						
Age (years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/Pacific Islander No. (%)	Native American No. (%)	Total (1) No. (%)
0 - 12	0 (0.0)	3 (1.4)	4 (4.9)	1 (3.1)	0 (0.0)	8 (1.6)
13 - 19	0 (0.0)	1 (0.5)	1 (1.2)	0 (0.0)	0 (0.0)	2 (0.4)
20 - 24	7 (4.5)	5 (2.3)	6 (7.4)	4 (12.5)	0 (0.0)	22 (4.4)
25 - 29	22 (14.2)	20 (9.1)	16 (19.8)	6 (18.8)	0 (0.0)	65 (13.1)
30 - 39	66 (42.6)	89 (40.5)	24 (29.6)	12 (37.5)	4 (80.0)	196 (39.6)
40 - 49	42 (27.1)	73 (33.2)	20 (24.7)	5 (15.6)	1 (20.0)	141 (28.5)
50 - 59	18 (11.6)	25 (11.4)	5 (6.2)	3 (9.4)	0 (0.0)	51 (10.3)
60 +	0 (0.0)	4 (1.8)	5 (6.2)	1 (3.1)	0 (0.0)	10 (2.0)
Female subtotal	155 (100)	220 (100)	81 (100)	32 (100)	5 (100)	495 (100)
Transgender (2)						
Age (years)	White No. (%)	African American No. (%)	Latino No. (%)	Asian/Pacific Islander Native American No. (%)	Total (1) No. (%)	
13 - 29	13 (29.5)	13 (24.5)	13 (24.5)	4 (18.2)	43 (24.9)	
30 +	31 (70.5)	40 (75.5)	40 (75.5)	18 (81.8)	130 (75.1)	
Transgender subtotal	44 (100)	53 (100)	53 (100)	22 (100)	173 (100)	

* Residents of San Francisco at time of initial AIDS diagnosis.

(1) Total includes persons with multiple or unknown race.

(2) Transgender information was collected since September 1996. Certain age or race/ethnic groups are combined for transgender cases because of small number.

Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report
 Summary of San Francisco Residents with AIDS*
 Reported as of 09/30/2005

12

Table 15. Living AIDS Cases by Initial AIDS-Defining Condition, San Francisco

Initial AIDS-Defining Condition	Def. (1) No.	Pres. (2) No.	Total No. (%)
Bacterial infections, recurrent, < 13 years [HIV+(3)]	7	N/A(4)	7 (0.1)
Candidiasis of bronchi, trachea, or lungs	5	N/A	5 (0.1)
Candidiasis of esophagus	49	55	104 (1.2)
Cervical cancer, invasive [HIV+]	2	N/A	2 (0.0)
Coccidioidomycosis, disseminated or extrapulmonary [HIV+]	1	N/A	1 (0.0)
Cryptococcosis, extrapulmonary	46	N/A	46 (0.5)
Cryptosporidiosis, intestinal (> 1 mo. duration)	114	N/A	114 (1.3)
Cytomegalovirus (except liver, spleen, lymph nodes), > 1 month of age	27	N/A	27 (0.3)
CMV retinitis with loss of vision [HIV+]	0	23	23 (0.3)
HIV encephalopathy [HIV+]	72	N/A	72 (0.8)
Herpes simplex: chronic (> 1 mo. duration), bronchitis, pneumonitis, esophagitis	15	N/A	15 (0.2)
Histoplasmosis, disseminated or extrapulmonary [HIV+]	8	N/A	8 (0.1)
Isosporiasis, intestinal (> 1 mo. duration) [HIV+]	8	N/A	8 (0.1)
Kaposi's sarcoma (5)	316	61	377 (4.4)
Lymphoid interstitial pneumonia/pulmonary lymphoid hyperplasia, < 13 years	2	0	2 (0.0)
Lymphoma, Burkitt's (non-Hodgkin's) [HIV+]	19	N/A	19 (0.2)
Lymphoma, immunoblastic (non-Hodgkin's) [HIV+]	54	N/A	54 (0.6)
Lymphoma, primary in brain (5)	3	N/A	3 (0.0)
Mycobacterium avium complex or M. kansasii, disseminated or extrapulmonary	19	0	19 (0.2)
Mycobacterium tuberculosis, pulmonary [HIV+]	78	5	83 (1.0)
Mycobacterium tuberculosis, disseminated or extrapulmonary [HIV+]	36	1	37 (0.4)
Mycobacterium other species, disseminated or extrapulmonary [HIV+]	2	2	4 (0.0)
Pneumocystis carinii pneumonia	421	137	558 (6.5)
Pneumonia, recurrent [HIV+]	94	8	102 (1.2)
Progressive multifocal leukoencephalopathy	1	N/A	1 (0.0)
Salmonella sepsis, recurrent [HIV+]	1	N/A	1 (0.0)
Toxoplasmosis of brain, > 1 month of age	10	26	36 (0.4)
Wasting syndrome [HIV+]	145	N/A	145 (1.7)
CD4 T lymphocyte count < 200 or percent < 14 [HIV+]	6748	N/A	6748 (78.3)
Any AIDS indicator condition and HIV-negative and CD4 count < 400 (6)	1	N/A	1 (0.0)
Total	8304	318	8622 (100)

* Residents of San Francisco at time of initial AIDS diagnosis.

(1) Indicator conditions diagnosed definitively (e.g. culture or biopsy proven).

(2) Indicator conditions diagnosed presumptively in a person who has laboratory evidence of HIV infection.

(3) [HIV+]: Indicator conditions that require laboratory evidence of HIV infection.

(4) N/A: Conditions which require definitive diagnoses only.

(5) Laboratory evidence of HIV infection in persons > 60 years of age.

(6) In the absence of other causes of immunocompromise.

Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report
 Summary of San Francisco Residents with AIDS*
 Reported as of 09/30/2005

Table 16. Cumulative AIDS Indicator Conditions among Persons Living with AIDS, San Francisco

AIDS Indicator Condition (1)	Total No. (%)
Bacterial infections, recurrent, < 13 years [HIV+(2)]	10 (0.1)
Candidiasis of bronchi, trachea, or lungs	22 (0.3)
Candidiasis of esophagus	416 (4.8)
Cervical cancer, invasive [HIV+]	3 (0.0)
Coccidioidomycosis, disseminated or extrapulmonary [HIV+]	10 (0.1)
Cryptococcosis, extrapulmonary	250 (2.9)
Cryptosporidiosis, intestinal (> 1 mo. duration)	318 (3.7)
Cytomegalovirus (except liver, spleen, lymph nodes), > 1 month of age	202 (2.3)
CMV retinitis with loss of vision [HIV+]	201 (2.3)
HIV encephalopathy [HIV+]	214 (2.5)
Herpes simplex: chronic (> 1 mo. duration), bronchitis, pneumonitis, esophagitis	72 (0.8)
Histoplasmosis, disseminated or extrapulmonary [HIV+]	26 (0.3)
Isosporiasis, intestinal (> 1 mo. duration) [HIV+]	16 (0.2)
Kaposi's sarcoma (3)	865 (10.0)
Lymphoid interstitial pneumonia/pulmonary lymphoid hyperplasia, < 13 years	2 (0.0)
Lymphoma, Burkitt's (non-Hodgkin's) [HIV+]	57 (0.7)
Lymphoma, immunoblastic (non-Hodgkin's) [HIV+]	119 (1.4)
Lymphoma, primary in brain (3)	16 (0.2)
Mycobacterium avium complex or M. kansasii, disseminated or extrapulmonary	305 (3.5)
Mycobacterium tuberculosis, pulmonary [HIV+]	190 (2.2)
Mycobacterium tuberculosis, disseminated or extrapulmonary [HIV+]	93 (1.1)
Mycobacterium other species, disseminated or extrapulmonary [HIV+]	43 (0.5)
Pneumocystis carinii pneumonia	1359 (15.8)
Pneumonia, recurrent [HIV+]	286 (3.3)
Progressive multifocal leukoencephalopathy	14 (0.2)
Salmonella sepsis, recurrent [HIV+]	3 (0.0)
Toxoplasmosis of brain, > 1 month of age	88 (1.0)
Wasting syndrome [HIV+]	656 (7.6)

* Residents of San Francisco at time of initial AIDS diagnosis.

(1) Cases may have more than one condition.

(2) [HIV+]: Indicator conditions that require laboratory evidence of HIV infection.

(3) Laboratory evidence of HIV infection in persons > 60 years of age.