

QUARTERLY AIDS SURVEILLANCE REPORT

San Francisco Department of Public Health AIDS Cases Reported Through June 2010

| Contents | Page |
|--|------|
| Surveillance Summary..... | 1 |
| Table 1: Adult/Adolescent AIDS Cases by Transmission Category..... | 2 |
| Table 2: AIDS Cases by Gender and Year of Diagnosis..... | 2 |
| Table 3: AIDS Cases by Transmission Category and Race/Ethnicity..... | 3 |
| Table 4: AIDS Cases by Transmission Category and Asian/Pacific Islander Ethnicity..... | 3 |
| Table 5: AIDS Cases by Transmission Category and Year of Diagnosis..... | 4 |
| Table 6: AIDS Cases by Gender, Age Group and Race/Ethnicity..... | 5 |
| Table 7: AIDS Cases by Race/Ethnicity and Year of Diagnosis..... | 6 |
| Table 8: AIDS Cases and Cumulative Rates per 100,000 by Race/Ethnicity and Gender..... | 6 |
| Table 9: AIDS Incidence, Mortality, and Prevalence by Year..... | 7 |
| Table 10: Cases by Initial AIDS-Defining Condition..... | 8 |
| Table 11: Cumulative AIDS Indicator Conditions among Persons with AIDS..... | 9 |
| Table 12: Living Adult/Adolescent AIDS Cases by Transmission Category..... | 10 |
| Table 13: Living AIDS Cases by Transmission Category and Race/Ethnicity..... | 10 |
| Table 14: Living AIDS Cases by Gender, Age Group and Race/Ethnicity..... | 11 |
| Table 15: Living AIDS Cases by Initial AIDS-Defining Condition..... | 12 |
| Table 16: Cumulative AIDS Indicator Conditions among Persons Living with AIDS..... | 13 |

The AIDS Surveillance Report is published quarterly by the
San Francisco Department of Public Health, HIV Epidemiology Section
25 Van Ness Avenue, Suite 500, San Francisco, CA 94102; Phone (415) 554-9050, FAX (415) 431-0353
Director of Health: Mitchell Katz, MD; Section Directors: Susan Scheer, PhD, MPH, Ling Hsu, MPH;
Program Coordinators: Maree Kay Parisi, Viva Delgado, MPH;
Epidemiologists: Miao-Jung Mia Chen, PhD, MPH, Anne Hirozawa, MPH, Alison Hughes, MPH,
Priscilla Lee Chu, MPH, Sharon Pipkin, MPH, Tara Schubert, MS, Annie Vu, MPH
The AIDS Surveillance Report is accessible via internet:
www.sfdph.org/dph/comupg/oprograms/hivepisecc/default.asp

City and County of San Francisco

Department of Public Health



Gavin Newsom

Mayor

Mitchell M. Katz, MD

Director

August 3, 2010

Dear Colleague:

We would like to inform you that we have made corrections to three previously published *Quarterly AIDS Surveillance Reports*. These reports contain data on AIDS cases reported through September 2009, December 2009 and March 2010. We have identified an error in the computer code used to summarize cumulative AIDS indicator conditions following a change in the case reporting data management system. This resulted in the numbers and percentages for certain AIDS indicator conditions being displayed incorrectly in Table 11 (Cumulative AIDS Indicator Conditions among Persons with AIDS) and Table 16 (Cumulative AIDS Indicator Conditions among Persons Living with AIDS). Those errors have been corrected and the revised reports have been posted online at www.sfdph.org/dph/comupg/oprograms/hivepisecc/default.asp.

If you have questions concerning our reports, or would like to have more information, please contact the HIV Epidemiology Section of the San Francisco Department of Public Health at (415) 554-9050.

Sincerely,

Handwritten signature of Ling Hsu in black ink.

Ling Hsu

Director, HIV Core Surveillance

Handwritten signature of Susan Scheer in black ink.

Susan Scheer

Director, HIV Epidemiology Section

AIDS Surveillance Summary

San Francisco (as of 06/30/2010)

| | |
|---------------------------------|--------|
| Cumulative cases ¹ : | 28,584 |
| Cumulative deaths: | 19,199 |


California (as of 03/31/2010)

| | |
|--------------------|---------|
| Cumulative cases: | 157,405 |
| Cumulative deaths: | 87,885 |

United States²

| | |
|---------------------------------------|-----------|
| Cumulative cases (as of 12/31/2008): | 1,106,391 |
| Cumulative deaths (as of 12/31/2007): | 597,499 |

San Francisco AIDS Incidence, Mortality, and Prevalence by Year, 1980-2010³



1. Includes SF residents diagnosed in SF and SF residents diagnosed in other jurisdictions. Excludes persons diagnosed in SF who resided in other jurisdictions at the time of their AIDS diagnosis.
2. The US numbers do not represent actual cases or deaths reported. Rather, these numbers are estimated and adjusted for reporting delays. For additional details, see www.cdc.gov/hiv/topics/surveillance/resources/reports.
3. Reporting for recent year is incomplete. See Table 9 for actual numbers per year.

**Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report
Summary of San Francisco Residents with AIDS*
Reported as of 06/30/2010**

Table 1. Adult/Adolescent AIDS Cases (>12 years) by Transmission Category, San Francisco, 1980-2010

| Transmission Category | No. | (%) |
|--|--------------|---------------|
| Gay or bisexual male | 20993 | (73.5) |
| Heterosexual male injection drug user | 1412 | (4.9) |
| Heterosexual female injection drug user | 701 | (2.5) |
| Gay or bisexual male injection drug user | 4110 | (14.4) |
| Lesbian or bisexual injection drug user | 62 | (0.2) |
| Transgender (1) | 408 | (1.4) |
| Hemophiliac | 16 | (0.1) |
| Heterosexual contact male (2) | 144 | (0.5) |
| Heterosexual contact female (2) | 309 | (1.1) |
| Transfusion recipient | 143 | (0.5) |
| Risk not reported/Other (3) | 248 | (0.9) |
| Total | 28546 | (100) |

Table 2. AIDS Cases by Gender and Year of Diagnosis, San Francisco, 1980-2010

| Gender | Year of Diagnosis | | | | | |
|-----------------|---------------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | < 2000 No. (%) | 2000 No. (%) | 2001 No. (%) | 2002 No. (%) | 2003 No. (%) | 2004 No. (%) |
| Male | 22774 (95.5) | 481 (86.5) | 455 (88.4) | 443 (89.1) | 497 (88.6) | 424 (88.3) |
| Female | 822 (3.4) | 52 (9.4) | 45 (8.7) | 36 (7.2) | 40 (7.1) | 39 (8.1) |
| Transgender (1) | 251 (1.1) | 23 (4.1) | 15 (2.9) | 18 (3.6) | 24 (4.3) | 17 (3.5) |
| Total | 23847 (100) | 556 (100) | 515 (100) | 497 (100) | 561 (100) | 480 (100) |

| Gender | Year of Diagnosis | | | | | |
|-----------------|-------------------|-------------------|-------------------|-------------------|-------------------|------------------|
| | 2005 No. (%) | 2006 No. (%) | 2007 No. (%) | 2008 No. (%) | 2009 No. (%) | 2010 No. (%) |
| Male | 419 (89.3) | 392 (90.3) | 393 (89.7) | 352 (87.6) | 267 (89.6) | 77 (88.5) |
| Female | 37 (7.9) | 30 (6.9) | 31 (7.1) | 41 (10.2) | 23 (7.7) | 6 (6.9) |
| Transgender (1) | 13 (2.8) | 12 (2.8) | 14 (3.2) | 9 (2.2) | 8 (2.7) | 4 (4.6) |
| Total | 469 (100) | 434 (100) | 438 (100) | 402 (100) | 298 (100) | 87 (100) |

* Residents of San Francisco at time of initial AIDS diagnosis.

(1) Transgender information was collected since September 1996. Data prior to this are incomplete.

(2) Includes persons who have had heterosexual contact with a person with HIV/AIDS or with a person who is at risk for HIV.

(3) Includes persons for whom risk information is incomplete (due to death, refusal to be interviewed or loss to follow-up), cases still under investigation, or interviewed patients who offered no plausible risk for HIV.

**Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report
Summary of San Francisco Residents with AIDS*
Reported as of 06/30/2010**

Table 3. AIDS Cases by Transmission Category and Race/Ethnicity, San Francisco, 1980-2010

| Transmission Category (1) | White No. (%) | African American No. (%) | Latino No. (%) | Asian/ Pacific Islander No. (%) | Native American No. (%) |
|-----------------------------------|---------------------|--------------------------------|--------------------|--|-------------------------------|
| Adult/Adolescent | | | | | |
| Gay or bisexual male | 16134 (79.9) | 1559 (43.2) | 2577 (73.6) | 697 (77.1) | 55 (43.3) |
| Injection drug user (IDU) | 744 (3.7) | 1039 (28.8) | 248 (7.1) | 39 (4.3) | 18 (14.2) |
| Gay or bisexual male IDU | 3006 (14.9) | 688 (19.0) | 461 (13.2) | 72 (8.0) | 46 (36.2) |
| Lesbian or bisexual IDU | 28 (0.1) | 23 (0.6) | 6 (0.2) | 2 (0.2) | 2 (1.6) |
| Hemophiliac | 8 (0.0) | 2 (0.1) | 5 (0.1) | 1 (0.1) | 0 (0.0) |
| Heterosexual (2) | 115 (0.6) | 186 (5.1) | 106 (3.0) | 44 (4.9) | 4 (3.1) |
| Transfusion recipient | 68 (0.3) | 23 (0.6) | 27 (0.8) | 19 (2.1) | 0 (0.0) |
| Risk not reported/Other (3) | 81 (0.4) | 79 (2.2) | 59 (1.7) | 25 (2.8) | 1 (0.8) |
| Pediatric (0-12 years) (4) | 8 (0.0) | 13 (0.4) | 10 (0.3) | 5 (0.6) | 1 (0.8) |
| Total | 20192 (100) | 3612 (100) | 3499 (100) | 904 (100) | 127 (100) |

Table 4. AIDS Cases by Transmission Category and Asian/Pacific Islander Ethnicity, San Francisco, 1980-2010

| Transmission Category (1) | Chinese No. (%) | Japanese No. (%) | Filipino No. (%) | Southeast Asian No. (%) | Korean No. (%) | Pacific Islander No. (%) |
|-----------------------------------|--------------------|---------------------|---------------------|-------------------------------|-------------------|--------------------------------|
| Adult/Adolescent | | | | | | |
| Gay or bisexual male | 154 (80.6) | 80 (82.5) | 224 (80.9) | 65 (73.0) | 9 (81.8) | 46 (65.7) |
| Injection drug user (IDU) | 6 (3.1) | 0 (0.0) | 8 (2.9) | 4 (4.5) | 1 (9.1) | 9 (12.9) |
| Gay or bisexual male IDU | 9 (4.7) | 12 (12.4) | 23 (8.3) | 7 (7.9) | 1 (9.1) | 8 (11.4) |
| Lesbian or bisexual IDU | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 2 (2.9) |
| Hemophiliac | 0 (0.0) | 1 (1.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) |
| Heterosexual (2) | 4 (2.1) | 2 (2.1) | 11 (4.0) | 5 (5.6) | 0 (0.0) | 4 (5.7) |
| Transfusion recipient | 9 (4.7) | 1 (1.0) | 6 (2.2) | 3 (3.4) | 0 (0.0) | 0 (0.0) |
| Risk not reported/Other (3) | 7 (3.7) | 1 (1.0) | 4 (1.4) | 5 (5.6) | 0 (0.0) | 1 (1.4) |
| Pediatric (0-12 years) (4) | 2 (1.0) | 0 (0.0) | 1 (0.4) | 0 (0.0) | 0 (0.0) | 0 (0.0) |
| Total | 191 (100) | 97 (100) | 277 (100) | 89 (100) | 11 (100) | 70 (100) |

* Residents of San Francisco at time of initial AIDS diagnosis.

- (1) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.
- (2) Includes persons who have had heterosexual contact with a person with HIV/AIDS or with a person who is at risk for HIV.
- (3) Includes persons for whom risk information is incomplete (due to death, refusal to be interviewed or loss to follow-up), cases still under investigation, or interviewed patients who offered no plausible risk for HIV.
- (4) Includes children who have hemophilia or other coagulation disorder, have received a blood transfusion, or who have acquired their infection from an infected mother during the perinatal period.

**Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report
Summary of San Francisco Residents with AIDS*
Reported as of 06/30/2010**

Table 5. AIDS Cases by Transmission Category and Year of Diagnosis, San Francisco, 1980-2010

| Transmission Category (1) | Year of Diagnosis | | | | | |
|-----------------------------|---------------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | < 2000 No. (%) | 2000 No. (%) | 2001 No. (%) | 2002 No. (%) | 2003 No. (%) | 2004 No. (%) |
| Adult/Adolescent | | | | | | |
| Gay or bisexual male | 18175 (76.2) | 337 (60.6) | 312 (60.6) | 308 (62.0) | 365 (65.1) | 316 (65.8) |
| Injection drug user (IDU) | 1561 (6.5) | 85 (15.3) | 69 (13.4) | 64 (12.9) | 77 (13.7) | 51 (10.6) |
| Gay or bisexual male IDU | 3505 (14.7) | 102 (18.3) | 91 (17.7) | 97 (19.5) | 84 (15.0) | 85 (17.7) |
| Lesbian or bisexual IDU | 38 (0.2) | 3 (0.5) | 3 (0.6) | 2 (0.4) | 3 (0.5) | 1 (0.2) |
| Hemophiliac | 15 (0.1) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) |
| Heterosexual (2) | 256 (1.1) | 19 (3.4) | 18 (3.5) | 15 (3.0) | 21 (3.7) | 14 (2.9) |
| Transfusion recipient | 141 (0.6) | 0 (0.0) | 1 (0.2) | 1 (0.2) | 0 (0.0) | 0 (0.0) |
| Risk not reported/Other (3) | 121 (0.5) | 9 (1.6) | 21 (4.1) | 9 (1.8) | 10 (1.8) | 13 (2.7) |
| Pediatric (0-12 years) (4) | 35 (0.1) | 1 (0.2) | 0 (0.0) | 1 (0.2) | 1 (0.2) | 0 (0.0) |
| Total | 23847 (100) | 556 (100) | 515 (100) | 497 (100) | 561 (100) | 480 (100) |

| Transmission Category (1) | Year of Diagnosis | | | | | |
|-----------------------------|-------------------|-------------------|-------------------|-------------------|-------------------|------------------|
| | 2005 No. (%) | 2006 No. (%) | 2007 No. (%) | 2008 No. (%) | 2009 No. (%) | 2010 No. (%) |
| Adult/Adolescent | | | | | | |
| Gay or bisexual male | 295 (62.9) | 286 (65.9) | 286 (65.3) | 252 (62.7) | 196 (65.8) | 43 (49.4) |
| Injection drug user (IDU) | 54 (11.5) | 41 (9.4) | 42 (9.6) | 33 (8.2) | 34 (11.4) | 8 (9.2) |
| Gay or bisexual male IDU | 85 (18.1) | 76 (17.5) | 70 (16.0) | 71 (17.7) | 40 (13.4) | 23 (26.4) |
| Lesbian or bisexual IDU | 3 (0.6) | 2 (0.5) | 0 (0.0) | 4 (1.0) | 2 (0.7) | 1 (1.1) |
| Hemophiliac | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 1 (0.3) | 0 (0.0) |
| Heterosexual (2) | 19 (4.1) | 20 (4.6) | 31 (7.1) | 24 (6.0) | 15 (5.0) | 5 (5.7) |
| Transfusion recipient | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) |
| Risk not reported/Other (3) | 13 (2.8) | 9 (2.1) | 9 (2.1) | 18 (4.5) | 10 (3.4) | 7 (8.0) |
| Pediatric (0-12 years) (4) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) |
| Total | 469 (100) | 434 (100) | 438 (100) | 402 (100) | 298 (100) | 87 (100) |

* Residents of San Francisco at time of initial AIDS diagnosis.

(1) Persons with more than one risk factor (other than the combinations listed in the tables) are tabulated only in the most likely transmission category.

(2) Includes persons who have had heterosexual contact with a person with HIV/AIDS or with a person who is at risk for HIV.

(3) Includes persons for whom risk information is incomplete (due to death, refusal to be interviewed or loss to follow-up), cases still under investigation, or interviewed patients who offered no plausible risk for HIV.

(4) Includes children who have hemophilia or other coagulation disorder, have received a blood transfusion, or who have acquired their infection from an infected mother during the perinatal period.

**Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report
Summary of San Francisco Residents with AIDS*
Reported as of 06/30/2010**

Table 6. AIDS Cases by Gender, Age Group and Race/Ethnicity, San Francisco, 1980-2010

| Male Age at Diagnosis (Years) | White No. (%) | African American No. (%) | Latino No. (%) | Asian/ Pacific Islander No. (%) | Native American No. (%) | Total (1) No. (%) |
|--|------------------|--------------------------------|-------------------|--|-------------------------------|----------------------|
| 0 - 12 | 4 (0.0) | 4 (0.1) | 5 (0.2) | 4 (0.5) | 1 (0.9) | 19 (0.1) |
| 13 - 19 | 12 (0.1) | 1 (0.0) | 13 (0.4) | 1 (0.1) | 2 (1.8) | 30 (0.1) |
| 20 - 24 | 306 (1.6) | 69 (2.3) | 119 (3.7) | 25 (3.1) | 5 (4.5) | 527 (2.0) |
| 25 - 29 | 1657 (8.4) | 247 (8.4) | 453 (14.1) | 83 (10.3) | 22 (20.0) | 2491 (9.2) |
| 30 - 39 | 8804 (44.7) | 1159 (39.4) | 1519 (47.4) | 358 (44.6) | 51 (46.4) | 11992 (44.5) |
| 40 - 49 | 6475 (32.9) | 981 (33.4) | 802 (25.0) | 247 (30.8) | 24 (21.8) | 8587 (31.8) |
| 50 - 59 | 1953 (9.9) | 376 (12.8) | 238 (7.4) | 64 (8.0) | 4 (3.6) | 2659 (9.9) |
| 60 + | 481 (2.4) | 104 (3.5) | 58 (1.8) | 21 (2.6) | 1 (0.9) | 669 (2.5) |
| Male subtotal | 19692 (100) | 2941 (100) | 3207 (100) | 803 (100) | 110 (100) | 26974 (100) |
| Female Age at Diagnosis (Years) | White No. (%) | African American No. (%) | Latino No. (%) | Asian/ Pacific Islander No. (%) | Native American No. (%) | Total (1) No. (%) |
| 0 - 12 | 4 (1.0) | 9 (1.7) | 5 (2.9) | 1 (1.6) | 0 (0.0) | 19 (1.6) |
| 13 - 19 | 1 (0.3) | 2 (0.4) | 2 (1.2) | 0 (0.0) | 0 (0.0) | 5 (0.4) |
| 20 - 24 | 15 (3.9) | 10 (1.8) | 10 (5.8) | 5 (7.8) | 1 (7.1) | 41 (3.4) |
| 25 - 29 | 41 (10.6) | 46 (8.4) | 24 (14.0) | 11 (17.2) | 1 (7.1) | 125 (10.4) |
| 30 - 39 | 154 (39.8) | 203 (37.2) | 59 (34.5) | 25 (39.1) | 8 (57.1) | 456 (37.9) |
| 40 - 49 | 103 (26.6) | 187 (34.3) | 43 (25.1) | 15 (23.4) | 4 (28.6) | 357 (29.7) |
| 50 - 59 | 40 (10.3) | 65 (11.9) | 15 (8.8) | 4 (6.3) | 0 (0.0) | 128 (10.6) |
| 60 + | 29 (7.5) | 23 (4.2) | 13 (7.6) | 3 (4.7) | 0 (0.0) | 71 (5.9) |
| Female subtotal | 387 (100) | 545 (100) | 171 (100) | 64 (100) | 14 (100) | 1202 (100) |
| Transgender (2) Age at Diagnosis (Years) | White No. (%) | African American No. (%) | Latino No. (%) | Asian/ Pacific Islander Native American No. (%) | Total (1) No. (%) | |
| 13 - 29 | 22 (19.5) | 27 (21.4) | 33 (27.3) | 15 (37.5) | 97 (23.8) | |
| 30 - 39 | 55 (48.7) | 47 (37.3) | 57 (47.1) | 17 (42.5) | 181 (44.4) | |
| 40 + | 36 (31.9) | 52 (41.3) | 31 (25.6) | 8 (20.0) | 130 (31.9) | |
| Transgender subtotal | 113 (100) | 126 (100) | 121 (100) | 40 (100) | 408 (100) | |

* Residents of San Francisco at time of initial AIDS diagnosis.

(1) Total includes persons with multiple or unknown race.

(2) Transgender information was collected since September 1996. Certain age or race/ethnic groups are combined for transgender cases because of small number.

**Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report
Summary of San Francisco Residents with AIDS*
Reported as of 06/30/2010**

Table 7. AIDS Cases by Race/Ethnicity and Year of Diagnosis, San Francisco, 1980-2010

| Race/Ethnicity | Year of Diagnosis | | | | | |
|------------------------|---------------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | < 2000 No. (%) | 2000 No. (%) | 2001 No. (%) | 2002 No. (%) | 2003 No. (%) | 2004 No. (%) |
| White | 17577 (73.7) | 317 (57.0) | 296 (57.5) | 293 (59.0) | 283 (50.4) | 261 (54.4) |
| African American | 2738 (11.5) | 113 (20.3) | 99 (19.2) | 89 (17.9) | 105 (18.7) | 78 (16.3) |
| Latino | 2640 (11.1) | 93 (16.7) | 74 (14.4) | 76 (15.3) | 126 (22.5) | 110 (22.9) |
| Asian/Pacific Islander | 619 (2.6) | 25 (4.5) | 34 (6.6) | 30 (6.0) | 35 (6.2) | 21 (4.4) |
| Native American | 92 (0.4) | 2 (0.4) | 5 (1.0) | 2 (0.4) | 6 (1.1) | 4 (0.8) |
| Total (1) | 23847 (100) | 556 (100) | 515 (100) | 497 (100) | 561 (100) | 480 (100) |

| Race/Ethnicity | Year of Diagnosis | | | | | |
|------------------------|-------------------|-------------------|-------------------|-------------------|-------------------|------------------|
| | 2005 No. (%) | 2006 No. (%) | 2007 No. (%) | 2008 No. (%) | 2009 No. (%) | 2010 No. (%) |
| White | 272 (58.0) | 242 (55.8) | 250 (57.1) | 209 (52.0) | 153 (51.3) | 39 (44.8) |
| African American | 81 (17.3) | 80 (18.4) | 79 (18.0) | 76 (18.9) | 59 (19.8) | 15 (17.2) |
| Latino | 86 (18.3) | 78 (18.0) | 73 (16.7) | 67 (16.7) | 57 (19.1) | 19 (21.8) |
| Asian/Pacific Islander | 24 (5.1) | 23 (5.3) | 28 (6.4) | 40 (10.0) | 17 (5.7) | 8 (9.2) |
| Native American | 3 (0.6) | 6 (1.4) | 0 (0.0) | 3 (0.7) | 3 (1.0) | 1 (1.1) |
| Total (1) | 469 (100) | 434 (100) | 438 (100) | 402 (100) | 298 (100) | 87 (100) |

Table 8. AIDS Cases and Cumulative Rates per 100,000 by Race/Ethnicity and Gender, San Francisco

| Race/Ethnicity | Male | Female | Total (2) |
|------------------------|------------------------|----------------------|------------------------|
| | No. (Rate) | No. (Rate) | No. (Rate) |
| White | 19692 (11379.2) | 387 (235.9) | 20079 (5956.1) |
| African American | 2941 (7890.0) | 545 (1395.0) | 3486 (4566.2) |
| Latino | 3207 (6249.9) | 171 (346.1) | 3378 (3354.0) |
| Asian/Pacific Islander | 803 (813.3) | 64 (59.8) | 867 (421.5) |
| Native American | 110 (8005.8) | 14 (1110.2) | 124 (4705.9) |
| Total (1) | 26974 (7441.2) | 1202 (332.5) | 28176 (3891.9) |

* Residents of San Francisco at time of initial AIDS diagnosis.

(1) Total includes persons with multiple or unknown race.

(2) Transgender cases were excluded because population size for transgender can not be determined for rates calculation.

**Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report
Summary of San Francisco Residents with AIDS*
Reported as of 06/30/2010**

Table 9. AIDS Incidence, Mortality, and Prevalence by Year, San Francisco, 1980-2010

| Year | Number of Cases Reported per Year | Number of Cases Diagnosed per Year (1) | Number of Deaths Occurred per Year (1) | Number of Persons Living with AIDS (1) |
|-------|---|--|--|--|
| 1980 | 0 | 3 | 0 | 3 |
| 1981 | 21 | 26 | 8 | 21 |
| 1982 | 75 | 99 | 32 | 88 |
| 1983 | 197 | 274 | 111 | 251 |
| 1984 | 451 | 557 | 273 | 535 |
| 1985 | 673 | 859 | 534 | 860 |
| 1986 | 981 | 1236 | 807 | 1289 |
| 1987 | 1287 | 1629 | 877 | 2041 |
| 1988 | 1408 | 1763 | 1038 | 2766 |
| 1989 | 1584 | 2161 | 1275 | 3652 |
| 1990 | 1686 | 2046 | 1364 | 4334 |
| 1991 | 1685 | 2284 | 1505 | 5113 |
| 1992 | 1638 | 2327 | 1641 | 5799 |
| 1993 | 4271 | 2073 | 1599 | 6273 |
| 1994 | 1915 | 1787 | 1592 | 6468 |
| 1995 | 1633 | 1564 | 1481 | 6551 |
| 1996 | 1242 | 1082 | 987 | 6646 |
| 1997 | 1060 | 804 | 422 | 7028 |
| 1998 | 794 | 695 | 401 | 7322 |
| 1999 | 724 | 578 | 356 | 7544 |
| 2000 | 626 | 556 | 348 | 7752 |
| 2001 | 497 | 515 | 322 | 7945 |
| 2002 | 440 | 497 | 323 | 8119 |
| 2003 | 533 | 561 | 302 | 8378 |
| 2004 | 557 | 480 | 305 | 8553 |
| 2005 | 500 | 469 | 312 | 8710 |
| 2006 | 424 | 434 | 288 | 8856 |
| 2007 | 537 | 438 | 269 | 9025 |
| 2008 | 560 | 402 | 224 | 9203 |
| 2009 | 357 | 298 | 148 | 9353 |
| 2010 | 228 | 87 | 55 | 9385 |
| Total | 28584 | 28584 | 19199 | |

* Residents of San Francisco at time of initial AIDS diagnosis.

(1) Data in recent years is incomplete due to delay in cases/deaths reporting.

**Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report
Summary of San Francisco Residents with AIDS*
Reported as of 06/30/2010**

Table 10. Cases by Initial AIDS-Defining Condition, San Francisco, 1980-2010

| Initial AIDS-Defining Condition | Def. (1) No. | Pres. (2) No. | Total No. (%) |
|--|-----------------|------------------|---------------------|
| Bacterial infections, recurrent, <13 years [HIV+(3)] | 8 | N/A(4) | 8 (0.0) |
| Candidiasis of bronchi, trachea, or lungs | 25 | N/A | 25 (0.1) |
| Candidiasis of esophagus | 303 | 226 | 529 (1.9) |
| Cervical cancer, invasive [HIV+] | 5 | N/A | 5 (0.0) |
| Coccidioidomycosis, disseminated or extrapulmonary [HIV+] | 8 | N/A | 8 (0.0) |
| Cryptococcosis, extrapulmonary | 380 | N/A | 380 (1.3) |
| Cryptosporidiosis, intestinal (>1 mo. duration) | 334 | N/A | 334 (1.2) |
| Cytomegalovirus (except liver, spleen, lymph nodes), >1 month of age | 204 | N/A | 204 (0.7) |
| CMV retinitis with loss of vision [HIV+] | 0 | 113 | 113 (0.4) |
| HIV encephalopathy [HIV+] | 392 | N/A | 392 (1.4) |
| Herpes simplex: chronic (>1 mo.), bronchitis, pneumonitis, esophagitis | 100 | N/A | 100 (0.3) |
| Histoplasmosis, disseminated or extrapulmonary [HIV+] | 26 | N/A | 26 (0.1) |
| Isosporiasis, intestinal (>1 mo. duration) [HIV+] | 21 | N/A | 21 (0.1) |
| Kaposi's sarcoma (5) | 2551 | 293 | 2844 (9.9) |
| Lymphoid interstitial pneumonia/pulmonary lymphoid hyperplasia, <13 years | 4 | 0 | 4 (0.0) |
| Lymphoma, Burkitt's (non-Hodgkin's) [HIV+] | 108 | N/A | 108 (0.4) |
| Lymphoma, immunoblastic (non-Hodgkin's) [HIV+] | 420 | N/A | 420 (1.5) |
| Lymphoma, primary in brain (5) | 40 | N/A | 40 (0.1) |
| Mycobacterium avium complex or M. kansasii, disseminated or extrapulmonary | 257 | 3 | 260 (0.9) |
| Mycobacterium tuberculosis, pulmonary [HIV+] | 205 | 15 | 220 (0.8) |
| Mycobacterium tuberculosis, disseminated or extrapulmonary [HIV+] | 144 | 5 | 149 (0.5) |
| Mycobacterium other species, disseminated or extrapulmonary [HIV+] | 16 | 11 | 27 (0.1) |
| Pneumocystis carinii pneumonia | 5099 | 604 | 5703 (20.0) |
| Pneumonia, recurrent [HIV+] | 220 | 27 | 247 (0.9) |
| Progressive multifocal leukoencephalopathy | 50 | N/A | 50 (0.2) |
| Salmonella sepsis, recurrent [HIV+] | 8 | N/A | 8 (0.0) |
| Toxoplasmosis of brain, >1 month of age | 43 | 200 | 243 (0.9) |
| Wasting syndrome [HIV+] | 683 | N/A | 683 (2.4) |
| CD4 T lymphocyte count <200 or percent <14 [HIV+] | 15429 | N/A | 15429 (54.0) |
| Any AIDS indicator condition and HIV-negative and CD4 count <400 (6) | 4 | N/A | 4 (0.0) |
| Total | 27087 | 1497 | 28584 (100) |

* Residents of San Francisco at time of initial AIDS diagnosis.

(1) Indicator conditions diagnosed definitively (e.g. culture or biopsy proven).

(2) Indicator conditions diagnosed presumptively in a person who has laboratory evidence of HIV infection.

(3) [HIV+]: Indicator conditions that require laboratory evidence of HIV infection.

(4) N/A: Conditions which require definitive diagnoses only.

(5) Laboratory evidence of HIV infection in persons > 60 years of age.

(6) In the absence of other causes of immunocompromise.

**Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report
Summary of San Francisco Residents with AIDS*
Reported as of 06/30/2010**

Table 11. Cumulative AIDS Indicator Conditions among Persons with AIDS, San Francisco, 1980-2010

| AIDS Indicator Condition (1) | Total No. (%) |
|--|------------------|
| Bacterial infections, recurrent, <13 years [HIV+(2)] | 11 (0.0) |
| Candidiasis of bronchi, trachea, or lungs | 138 (0.5) |
| Candidiasis of esophagus | 2556 (8.9) |
| Cervical cancer, invasive [HIV+] | 11 (0.0) |
| Coccidioidomycosis, disseminated or extrapulmonary [HIV+] | 62 (0.2) |
| Cryptococcosis, extrapulmonary | 1859 (6.5) |
| Cryptosporidiosis, intestinal (>1 mo. duration) | 1373 (4.8) |
| Cytomegalovirus (except liver, spleen, lymph nodes), >1 month of age | 2449 (8.6) |
| CMV retinitis with loss of vision [HIV+] | 2486 (8.7) |
| HIV encephalopathy [HIV+] | 2479 (8.7) |
| Herpes simplex: chronic (>1 mo.), bronchitis, pneumonitis, esophagitis | 436 (1.5) |
| Histoplasmosis, disseminated or extrapulmonary [HIV+] | 158 (0.6) |
| Isosporiasis, intestinal (>1 mo. duration) [HIV+] | 67 (0.2) |
| Kaposi's sarcoma (3) | 6708 (23.5) |
| Lymphoid interstitial pneumonia/pulmonary lymphoid hyperplasia, <13 years | 5 (0.0) |
| Lymphoma, Burkitt's (non-Hodgkin's) [HIV+] | 545 (1.9) |
| Lymphoma, immunoblastic (non-Hodgkin's) [HIV+] | 1131 (4.0) |
| Lymphoma, primary in brain (3) | 390 (1.4) |
| Mycobacterium avium complex or M. kansasii, disseminated or extrapulmonary | 5074 (17.8) |
| Mycobacterium tuberculosis, pulmonary [HIV+] | 654 (2.3) |
| Mycobacterium tuberculosis, disseminated or extrapulmonary [HIV+] | 478 (1.7) |
| Mycobacterium other species, disseminated or extrapulmonary [HIV+] | 332 (1.2) |
| Pneumocystis carinii pneumonia | 10986 (38.4) |
| Pneumonia, recurrent [HIV+] | 1005 (3.5) |
| Progressive multifocal leukoencephalopathy | 310 (1.1) |
| Salmonella sepsis, recurrent [HIV+] | 58 (0.2) |
| Toxoplasmosis of brain, >1 month of age | 1159 (4.1) |
| Wasting syndrome [HIV+] | 4331 (15.2) |

* Residents of San Francisco at time of initial AIDS diagnosis.

(1) Cases may have more than one condition.

(2) [HIV+]: Indicator conditions that require laboratory evidence of HIV infection.

(3) Laboratory evidence of HIV infection in persons > 60 years of age.

**Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report
Summary of San Francisco Residents with AIDS*
Reported as of 06/30/2010**

Table 12. Living Adult/Adolescent AIDS Cases (>12 years) by Transmission Category, San Francisco

| Transmission Category | No. | (%) |
|--|-------------|---------------|
| Gay or bisexual male | 6718 | (71.7) |
| Heterosexual male injection drug user | 459 | (4.9) |
| Heterosexual female injection drug user | 279 | (3.0) |
| Gay or bisexual male injection drug user | 1270 | (13.6) |
| Lesbian or bisexual injection drug user | 29 | (0.3) |
| Transgender (1) | 197 | (2.1) |
| Hemophiliac | 5 | (0.1) |
| Heterosexual contact male (2) | 91 | (1.0) |
| Heterosexual contact female (2) | 179 | (1.9) |
| Transfusion recipient | 20 | (0.2) |
| Risk not reported/Other (3) | 124 | (1.3) |
| Total | 9371 | (100) |

Table 13. Living AIDS Cases by Transmission Category and Race/Ethnicity, San Francisco

| Transmission Category | White No. (%) | African American No. (%) | Latino No. (%) | Asian/ Pacific Islander No. (%) | Native American No. (%) |
|-----------------------------------|--------------------|--------------------------------|--------------------|--|-------------------------------|
| Adult/Adolescent | | | | | |
| Gay or bisexual male | 4690 (78.9) | 551 (42.1) | 1143 (74.9) | 364 (76.0) | 25 (44.6) |
| Injection drug user (IDU) | 269 (4.5) | 348 (26.6) | 88 (5.8) | 17 (3.5) | 10 (17.9) |
| Gay or bisexual male IDU | 871 (14.6) | 240 (18.3) | 182 (11.9) | 42 (8.8) | 16 (28.6) |
| Lesbian or bisexual IDU | 13 (0.2) | 13 (1.0) | 1 (0.1) | 1 (0.2) | 1 (1.8) |
| Hemophiliac | 3 (0.1) | 2 (0.2) | 0 (0.0) | 0 (0.0) | 0 (0.0) |
| Heterosexual (2) | 54 (0.9) | 107 (8.2) | 75 (4.9) | 33 (6.9) | 3 (5.4) |
| Transfusion recipient | 7 (0.1) | 2 (0.2) | 5 (0.3) | 6 (1.3) | 0 (0.0) |
| Risk not reported/Other (3) | 41 (0.7) | 41 (3.1) | 28 (1.8) | 13 (2.7) | 0 (0.0) |
| Pediatric (0-12 years) (4) | 0 (0.0) | 4 (0.3) | 5 (0.3) | 3 (0.6) | 1 (1.8) |
| Total | 5948 (100) | 1308 (100) | 1527 (100) | 479 (100) | 56 (100) |

* Residents of San Francisco at time of initial AIDS diagnosis.

- (1) Transgender information was collected since September 1996. Data prior to this are incomplete.
- (2) Includes persons who have had heterosexual contact with a person with HIV/AIDS or with a person who is at risk for HIV.
- (3) Includes persons for whom risk information is incomplete (due to death, refusal to be interviewed or loss to follow-up), cases still under investigation, or interviewed patients who offered no plausible risk for HIV.
- (4) Includes children who have hemophilia or other coagulation disorder, have received a blood transfusion, or who have acquired their infection from an infected mother during the perinatal period.

**Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report
Summary of San Francisco Residents with AIDS*
Reported as of 06/30/2010**

Table 14. Living AIDS Cases by Gender, Race/Ethnicity and Age as of 06/30/2010, San Francisco

| Male Current Age (Years) | White No. (%) | African American No. (%) | Latino No. (%) | Asian/ Pacific Islander No. (%) | Native American No. (%) | Total (1) No. (%) |
|---|------------------|--------------------------------|-------------------|--|-------------------------------|----------------------|
| 0 - 12 | 0 (0.0) | 0 (0.0) | 1 (0.1) | 0 (0.0) | 0 (0.0) | 2 (0.0) |
| 13 - 19 | 0 (0.0) | 1 (0.1) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 1 (0.0) |
| 20 - 24 | 5 (0.1) | 6 (0.6) | 12 (0.9) | 4 (1.0) | 1 (2.1) | 29 (0.3) |
| 25 - 29 | 40 (0.7) | 12 (1.2) | 36 (2.6) | 6 (1.5) | 3 (6.4) | 100 (1.2) |
| 30 - 39 | 391 (6.8) | 81 (8.0) | 207 (15.1) | 71 (17.2) | 2 (4.3) | 769 (8.9) |
| 40 - 49 | 2008 (34.9) | 378 (37.3) | 614 (44.8) | 169 (40.9) | 27 (57.4) | 3218 (37.2) |
| 50 - 59 | 2287 (39.8) | 374 (36.9) | 375 (27.3) | 114 (27.6) | 12 (25.5) | 3176 (36.7) |
| 60 + | 1015 (17.7) | 161 (15.9) | 127 (9.3) | 49 (11.9) | 2 (4.3) | 1357 (15.7) |
| Male subtotal | 5746 (100) | 1013 (100) | 1372 (100) | 413 (100) | 47 (100) | 8652 (100) |
| Female Current Age (Years) | White No. (%) | African American No. (%) | Latino No. (%) | Asian/ Pacific Islander No. (%) | Native American No. (%) | Total (1) No. (%) |
| 0 - 12 | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) |
| 13 - 19 | 0 (0.0) | 2 (0.9) | 3 (3.2) | 1 (2.5) | 0 (0.0) | 6 (1.1) |
| 20 - 24 | 0 (0.0) | 2 (0.9) | 0 (0.0) | 1 (2.5) | 0 (0.0) | 3 (0.6) |
| 25 - 29 | 2 (1.3) | 6 (2.6) | 4 (4.3) | 3 (7.5) | 0 (0.0) | 15 (2.8) |
| 30 - 39 | 16 (10.3) | 28 (11.9) | 19 (20.4) | 9 (22.5) | 2 (28.6) | 76 (14.2) |
| 40 - 49 | 72 (46.5) | 69 (29.4) | 27 (29.0) | 14 (35.0) | 1 (14.3) | 185 (34.5) |
| 50 - 59 | 46 (29.7) | 98 (41.7) | 26 (28.0) | 8 (20.0) | 4 (57.1) | 184 (34.3) |
| 60 + | 19 (12.3) | 30 (12.8) | 14 (15.1) | 4 (10.0) | 0 (0.0) | 67 (12.5) |
| Female subtotal | 155 (100) | 235 (100) | 93 (100) | 40 (100) | 7 (100) | 536 (100) |
| Transgender (2) Current Age (Years) | White No. (%) | African American No. (%) | Latino No. (%) | Asian/ Pacific Islander Native American No. (%) | Total (1) No. (%) | |
| 13 - 39 | 9 (19.1) | 11 (18.3) | 18 (29.0) | 9 (32.1) | 47 (23.9) | |
| 40 + | 38 (80.9) | 49 (81.7) | 44 (71.0) | 19 (67.9) | 150 (76.1) | |
| Transgender subtotal | 47 (100) | 60 (100) | 62 (100) | 28 (100) | 197 (100) | |

* Residents of San Francisco at time of initial AIDS diagnosis.

(1) Total includes persons with multiple or unknown race.

(2) Transgender information was collected since September 1996. Certain age or race/ethnic groups are combined for transgender cases because of small number.

**Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report
Summary of San Francisco Residents with AIDS*
Reported as of 06/30/2010**

Table 15. Living AIDS Cases by Initial AIDS-Defining Condition, San Francisco

| Initial AIDS-Defining Condition | Def. (1) No. | Pres. (2) No. | Total No. (%) |
|--|-----------------|------------------|--------------------|
| Bacterial infections, recurrent, <13 years [HIV+(3)] | 7 | N/A(4) | 7 (0.1) |
| Candidiasis of bronchi, trachea, or lungs | 6 | N/A | 6 (0.1) |
| Candidiasis of esophagus | 53 | 52 | 105 (1.1) |
| Cervical cancer, invasive [HIV+] | 0 | N/A | 0 (0.0) |
| Coccidioidomycosis, disseminated or extrapulmonary [HIV+] | 1 | N/A | 1 (0.0) |
| Cryptococcosis, extrapulmonary | 47 | N/A | 47 (0.5) |
| Cryptosporidiosis, intestinal (>1 mo. duration) | 106 | N/A | 106 (1.1) |
| Cytomegalovirus (except liver, spleen, lymph nodes), >1 month of age | 22 | N/A | 22 (0.2) |
| CMV retinitis with loss of vision [HIV+] | 0 | 21 | 21 (0.2) |
| HIV encephalopathy [HIV+] | 58 | N/A | 58 (0.6) |
| Herpes simplex: chronic (>1 mo.), bronchitis, pneumonitis, esophagitis | 15 | N/A | 15 (0.2) |
| Histoplasmosis, disseminated or extrapulmonary [HIV+] | 9 | N/A | 9 (0.1) |
| Isosporiasis, intestinal (>1 mo. duration) [HIV+] | 7 | N/A | 7 (0.1) |
| Kaposi's sarcoma (5) | 318 | 66 | 384 (4.1) |
| Lymphoid interstitial pneumonia/pulmonary lymphoid hyperplasia, <13 years | 2 | 0 | 2 (0.0) |
| Lymphoma, Burkitt's (non-Hodgkin's) [HIV+] | 23 | N/A | 23 (0.2) |
| Lymphoma, immunoblastic (non-Hodgkin's) [HIV+] | 47 | N/A | 47 (0.5) |
| Lymphoma, primary in brain (5) | 1 | N/A | 1 (0.0) |
| Mycobacterium avium complex or M. kansasii, disseminated or extrapulmonary | 18 | 0 | 18 (0.2) |
| Mycobacterium tuberculosis, pulmonary [HIV+] | 72 | 7 | 79 (0.8) |
| Mycobacterium tuberculosis, disseminated or extrapulmonary [HIV+] | 35 | 2 | 37 (0.4) |
| Mycobacterium other species, disseminated or extrapulmonary [HIV+] | 2 | 2 | 4 (0.0) |
| Pneumocystis carinii pneumonia | 409 | 142 | 551 (5.9) |
| Pneumonia, recurrent [HIV+] | 72 | 10 | 82 (0.9) |
| Progressive multifocal leukoencephalopathy | 0 | N/A | 0 (0.0) |
| Salmonella sepsis, recurrent [HIV+] | 1 | N/A | 1 (0.0) |
| Toxoplasmosis of brain, >1 month of age | 8 | 21 | 29 (0.3) |
| Wasting syndrome [HIV+] | 130 | N/A | 130 (1.4) |
| CD4 T lymphocyte count <200 or percent <14 [HIV+] | 7592 | N/A | 7592 (80.9) |
| Any AIDS indicator condition and HIV-negative and CD4 count <400 (6) | 1 | N/A | 1 (0.0) |
| Total | 9062 | 323 | 9385 (100) |

* Residents of San Francisco at time of initial AIDS diagnosis.

(1) Indicator conditions diagnosed definitively (e.g. culture or biopsy proven).

(2) Indicator conditions diagnosed presumptively in a person who has laboratory evidence of HIV infection.

(3) [HIV+]: Indicator conditions that require laboratory evidence of HIV infection.

(4) N/A: Conditions which require definitive diagnoses only.

(5) Laboratory evidence of HIV infection in persons > 60 years of age.

(6) In the absence of other causes of immunocompromise.

**Acquired Immunodeficiency Syndrome (AIDS) Quarterly Surveillance Report
Summary of San Francisco Residents with AIDS*
Reported as of 06/30/2010**

Table 16. Cumulative AIDS Indicator Conditions among Persons Living with AIDS, San Francisco

| AIDS Indicator Condition (1) | Total No. (%) |
|--|------------------|
| Bacterial infections, recurrent, <13 years [HIV+(2)] | 10 (0.1) |
| Candidiasis of bronchi, trachea, or lungs | 23 (0.2) |
| Candidiasis of esophagus | 439 (4.7) |
| Cervical cancer, invasive [HIV+] | 1 (0.0) |
| Coccidioidomycosis, disseminated or extrapulmonary [HIV+] | 10 (0.1) |
| Cryptococcosis, extrapulmonary | 269 (2.9) |
| Cryptosporidiosis, intestinal (>1 mo. duration) | 289 (3.1) |
| Cytomegalovirus (except liver, spleen, lymph nodes), >1 month of age | 205 (2.2) |
| CMV retinitis with loss of vision [HIV+] | 196 (2.1) |
| HIV encephalopathy [HIV+] | 187 (2.0) |
| Herpes simplex: chronic (>1 mo.), bronchitis, pneumonitis, esophagitis | 84 (0.9) |
| Histoplasmosis, disseminated or extrapulmonary [HIV+] | 26 (0.3) |
| Isosporiasis, intestinal (>1 mo. duration) [HIV+] | 15 (0.2) |
| Kaposi's sarcoma (3) | 872 (9.3) |
| Lymphoid interstitial pneumonia/pulmonary lymphoid hyperplasia, <13 years | 2 (0.0) |
| Lymphoma, Burkitt's (non-Hodgkin's) [HIV+] | 61 (0.6) |
| Lymphoma, immunoblastic (non-Hodgkin's) [HIV+] | 115 (1.2) |
| Lymphoma, primary in brain (3) | 13 (0.1) |
| Mycobacterium avium complex or M. kansasii, disseminated or extrapulmonary | 301 (3.2) |
| Mycobacterium tuberculosis, pulmonary [HIV+] | 181 (1.9) |
| Mycobacterium tuberculosis, disseminated or extrapulmonary [HIV+] | 98 (1.0) |
| Mycobacterium other species, disseminated or extrapulmonary [HIV+] | 38 (0.4) |
| Pneumocystis carinii pneumonia | 1392 (14.8) |
| Pneumonia, recurrent [HIV+] | 222 (2.4) |
| Progressive multifocal leukoencephalopathy | 13 (0.1) |
| Salmonella sepsis, recurrent [HIV+] | 4 (0.0) |
| Toxoplasmosis of brain, >1 month of age | 84 (0.9) |
| Wasting syndrome [HIV+] | 599 (6.4) |

* Residents of San Francisco at time of initial AIDS diagnosis.

(1) Cases may have more than one condition.

(2) [HIV+]: Indicator conditions that require laboratory evidence of HIV infection.

(3) Laboratory evidence of HIV infection in persons > 60 years of age.