



San Francisco Monthly STD Report

Data for June, 2016
Report prepared August 3, 2016

Table 1. STDs among residents, June, 2016.

	2016		2015	
	month	YTD	month	YTD
Gonorrhea	410	2,392	401	2,090
Male rectal gonorrhea	102	680	113	541
Chlamydia	669	3,916	586	3,396
Male rectal chlamydia	182	1059	148	757
Syphilis (adult total)	86	654	117	710
Primary & secondary	35	261	43	256
Early latent	42	295	56	354
Unknown latent	1	11	5	17
Late latent	8	87	13	83
Neurosyphilis	0	11	1	12
Congenital syphilis	0	0	0	0
PID	4	27	5	49




Figure 1. Monthly STD trends for San Francisco. Syphilis cases different scale.

Table 2. Selected STD cases and rates for San Francisco by age and race/ethnicity, 2016 through June only. Rates equal cases per 100,000 residents per year based on 2000 US Census data.

	(All races)		Asian/PI		African American		Hispanic		White	
	cases	rate	cases	rate	cases	rate	cases	rate	cases	rate
All ages										
Chlamydia	3,916	1,008.3	466	369.3	398	1,242.4	582	1,063.0	1,450	855.7
Gonorrhea	2,392	615.9	211	167.2	265	827.2	382	697.7	1,147	676.9
Early syphilis	556	143.2	62	49.1	56	174.8	121	221.0	279	164.7
Under 20 yrs										
Chlamydia	260	1,012.2	24	220.3	80	2,430.0	63	1,137.3	29	522.8
Gonorrhea	54	210.2	5	45.9	20	607.5	11	198.6	9	162.3
Early syphilis	2	7.8	0	0.0	1	30.4	1	18.1	0	0.0

Table 3. HIV testing among City Clinic patients, June, 2016.

	2016		2015	
	month	YTD	month	YTD
Tests	562	3,221	556	2,909
Antibody positive	8	34	4	27
Acute HIV infection	1	4	2	4

Note: All statistics are provisional until the annual report is released for the year. Morbidity is based on date of diagnosis. Totals for past months may change due to delays in reporting from labs and providers.




Figure 2. City Clinic visits by gender and orientation.




Figure 3. Average number of recent* sex partners for City Clinic visits by gender and sexual orientation. *Recall period is 3 months.

Outbreak of Meningococcal Disease involving Men who Have Sex With Men in Southern California; Recommendations for San Francisco

The San Francisco Department of Public Health (SFPDH) sent a Health Advisory on June 28, 2016 providing an update on (1) CDC's Advisory Committee on Immunization Practices expanded recommendation for immunization for invasive meningococcal disease (IMD) with quadrivalent meningococcal vaccine (MenACWY; Menactra® or Menveo®) of all HIV-positive persons over 2 years of age due to their higher risk and (2) an outbreak of IMD during May-June 2016 including primarily men who have sex with men (MSM) in Los Angeles and Orange Counties. Both the California Department of Public Health and SFPDH additionally are recommending vaccination for MSM or Transwomen who are not HIV-infected but who have other risk factors for meningococcal disease, including:

- Regularly have close or intimate contact with multiple partners, or who seek partners through the use of online websites or digital phone applications;
- Regularly visit crowded venues such as bars, parties, etc.;
- Smoke cigarettes, marijuana, hookahs, or illegal drugs or spend time in smoky settings.

While there is currently no outbreak of IMD among MSM in San Francisco, please take every opportunity to vaccinate individuals at risk for IMD. MSM with multiple sex partners should also be screened every three months for HIV and STDs, including rectal gonorrhea and chlamydia if they report receptive anal sex, and syphilis. HIV PrEP may also be indicated for HIV-uninfected persons.

Additional details on the new ACIP vaccination recommendation to prevent IMD: <http://www.sfdcp.org/meningococcal.html>

Additional information on PrEP: <http://sfcityclinic.org/services/prep.asp>. DS/TQN/SP

Provider STD Reporting: 415-487-5555, 415-431-4628 (fax)

www.sfcityclinic.org