2015 ZSFG JOINT COMMISSION/CMS CORE MEASURE RESULTS

| Measure | Measure Name | Q1 2015 | Q2 2015 | Q3 2015 | Q4 2015 | UHC Median (Q3 2015) |
|--|---|---------------------------|---------------------------|---------------------------|---------------------------|---|
| STROKE | | | | | | |
| STK-1** | Venous Thromboembolism (VTE) Prophylaxis | 100% | 100% | 100% | 95% | 100% |
| STK-2** | Discharged on Antithrombotic Therapy | 100% | 100% | 100% | 100% | 100% |
| STK-3** | Anticoagulation Therapy for Atrial Fibrillation/Flutter | 100% | 100% | 100% | 100% | 100% |
| STK-4 | Thrombolytic Therapy | 100% | 100% | 100% | 100% | 100% |
| STK-5** | Antithrombotic Therapy by End of Hospital Day 2 | 100% | 100% | 100% | 100% | 100% |
| STK-6** | Discharged on Statin Medication | 100% | 100% | 100% | 100% | 100% |
| STK-8** | Stroke Education | 100% | 100% | 100% | 100% | 100% |
| STK-10** | Assessed for Rehabilitation | 100% | 100% | 100% | 100% | 100% |
| PERINATAL | CARE | | | | | |
| PC-01 | Elective Delivery Prior to 39 Completed Weeks Gestation (lower = better) | 0% | 0% | 0% | 0% | 0% |
| PC-02 | Cesarean Section Rate (lower = better) | 8% | 17% | 18% | 22% | 23% |
| PC-03 | Antenatal Steroids Given as Appropriate | | 100% | no cases | no cases | 100% |
| PC-04 | Health Care-Associated Bloodstream Infections in Newborns (lower = better) | 0% | 0% | 0% | 0% | 0% |
| PC-05 | Exclusive Breast Milk Feeding During Hospital Stay | 70% | 79% | 73% | 67% | 55% |
| | ROMBOEMBOLISM (VTE) | , 0,0 | , 570 | , 370 | 0170 | 3370 |
| VTE-1** | Venous Thromboembolism Prophylaxis | 89% | 91% | 91% | 83% | 97% |
| VTE-2** | Intensive Care Unit Venous Thromboembolism Prophylaxis | 75% | 96% | 82% | 98% | 100% |
| VTE-2** | Venous Thromboembolism Patients with Anticoagulation Overlap Therapy | 100% | 97% | | 95% | 100% |
| VTE-5 | Venous Thromboembolism Warfarin Therapy Discharge Instructions | | | 95% | | |
| VTE-6 | Hospital Acquired Potentially-Preventable Venous Thromboembolism | 89% | 81% | 78% | 79% | 100% |
| | (lower = better) | 0% | 0% | 0% | 9% | 0% |
| IMMUNIZA | TION | | | | | |
| IMM-2 | Influenza Immunization | 71% | Not Flu Season | Not Flu Season | 77% | Not Flu Season |
| PAIN MAN | AGEMENT (Minutes) | | | | | |
| OP-21 | Median Time to Pain Management for Long Bone Fracture in the ED | 50 | 40 | 72 | 74 | 55 |
| EMERGENC | CY THROUGHPUT (Minutes) | | | | | |
| ED-1 | Median Time from ED Arrival to ED Departure for Admitted ED Patients | 506 | 469 | 422 | 487 | 352 |
| ED-2 | Median Time -Admit Decision Time to ED Departure Time for Admitted Patients | 215 | 222 | 190 | 211 | 137 |
| OP-18 | Median Time from ED Arrival to ED Departure for Discharged ED Patients | 257 | 229 | 238 | 227 | 190 |
| OP-20 | Median Time - ED Door to Diagnostic Evaluation by a Qualified Medical Personnel | 62 | 46 | 28 | 30 | 31 |
| | BASED INPATIENT PSYCHIATRY | 02 | 10 | 20 | | 51 |
| HBIPS-1 | Admission Screening Completed | 92% | 88% | 97% | 94% | 100% |
| HBIPS-2 | Hours of Physical Restraint Use (per 1000 patient hours) | 0.81 | 1.09 | 0.41 | 0.78 | 0 |
| | Hours of Seclusion Use (per 1000 patient hours) | 6 | 4 | 3 | 5 | 0 |
| | | U | 4 | 3 | 5 | 0 |
| | Patients discharged on multiple antipsychotic medications (lower=better) | 1 5 0/ | C0/ | 0.0/ | 1.20/ | 6 20/ |
| HBIPS-4** | Patients discharged on multiple antipsychotic medications (lower=better) Patients discharged on multiple antipsychotic medications with appropriate | 15% | 6% | 9% 5.0% | 12% | 6.2% |
| HBIPS-4** HBIPS-5 | Patients discharged on multiple antipsychotic medications (lower=better) Patients discharged on multiple antipsychotic medications with appropriate justification | 42% | 40% | 50% | 17% | 66.7% |
| HBIPS-4** HBIPS-5 | Patients discharged on multiple antipsychotic medications (lower=better) Patients discharged on multiple antipsychotic medications with appropriate justification Post discharge continuing care plan created | | | | | |
| HBIPS-4** HBIPS-5 HBIPS-6** HBIPS-7** | Patients discharged on multiple antipsychotic medications (lower=better) Patients discharged on multiple antipsychotic medications with appropriate justification Post discharge continuing care plan created Post discharge continuing care plan transmitted to next level of care provider upon discharge | 42% 100% 84% | 40% 100% 52% | 50% 100% 90% | 17% 100% 84% | 66.7% 91.3% 84.0% |
| HBIPS-4** HBIPS-5 HBIPS-6** HBIPS-7** | Patients discharged on multiple antipsychotic medications (lower=better) Patients discharged on multiple antipsychotic medications with appropriate justification Post discharge continuing care plan created Post discharge continuing care plan transmitted to next level of care | 42% 100% | 40% 100% | 50% 100% | 17% 100% | 66.7% 91.3% |
| HBIPS-4** HBIPS-5 HBIPS-6** HBIPS-7** SUB-1 | Patients discharged on multiple antipsychotic medications (lower=better) Patients discharged on multiple antipsychotic medications with appropriate justification Post discharge continuing care plan created Post discharge continuing care plan transmitted to next level of care provider upon discharge | 42% 100% 84% 91% | 40% 100% 52% 97% | 50% 100% 90% 94% | 17% 100% 84% 99% | 66.7% 91.3% 84.0% 96% |
| HBIPS-3 HBIPS-4** HBIPS-5 HBIPS-6** HBIPS-7** SUB-1 TOB-1 TOB-2 | Patients discharged on multiple antipsychotic medications (lower=better) Patients discharged on multiple antipsychotic medications with appropriate justification Post discharge continuing care plan created Post discharge continuing care plan transmitted to next level of care provider upon discharge Alcohol Use Screening | 42% 100% 84% | 40% 100% 52% | 50% 100% 90% | 17% 100% 84% | 66.7% 91.3% 84.0% 96% no UHC Median |

** = Retired Q1 2016